

National Parks and Wildlife Service

Conservation Objectives

Clew Bay Complex SAC 001482



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*
*Department of
Arts, Heritage and the Gaeltacht*

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

001482 Clew Bay Complex SAC

QI	Description
1013	Geyer's whorl snail <i>Vertigo geyeri</i>
1140	Mudflats and sandflats not covered by seawater at low tide
1150	* Coastal lagoons
1160	Large shallow inlets and bays
1210	Annual vegetation of drift lines
1220	Perennial vegetation of stony banks
1330	Atlantic salt meadows (<i>Glauco-Puccinellietalia maritimae</i>)
1355	Otter <i>Lutra lutra</i>
1365	Common seal <i>Phoca vitulina</i>
2110	Embryonic shifting dunes
2120	Shifting dunes along the shoreline with <i>Ammophila arenaria</i> ("white dunes")

Supporting documents, relevant reports & publications (listed by date)

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

Title:	Monitoring and Assessment of Irish Lagoons for the purpose of the EU Water Framework Directive
Year:	in prep
Author:	Roden, C.M.; Oliver, G.
Series:	Unpublished report to the EPA
Title:	Clew Bay Complex SAC (001482): Conservation objectives supporting document - marine habitats and species [Version 1]
Year:	2011
Author:	NPWS
Series:	Unpublished Report to NPWS
Title:	Clew Bay Complex SAC (001482): Conservation objectives supporting document - coastal habitats [Version 1]
Year:	2011
Author:	NPWS
Series:	Unpublished Report to NPWS
Title:	Otter tracking study of Roaringwater Bay
Year:	2010
Author:	De Jongh, A.; O'Neill, L.
Series:	Unpublished Draft Report to NPWS
Title:	Subtidal benthic surveys (Clew Bay)
Year:	2009
Author:	Aquafact
Series:	Unpublished Report to NPWS
Title:	Saltmarsh Monitoring Report 2007-2008
Year:	2009
Author:	McCorry, M.; Ryle, T.
Series:	Unpublished Report to NPWS
Title:	Clew Bay baseline intertidal survey
Year:	2009
Author:	RPS
Series:	Unpublished Report to NPWS
Title:	Coastal Monitoring Project 2004-2006
Year:	2009
Author:	Ryle, T.; Murray, A.; Connolly, C.; Swann, M.
Series:	Unpublished Report to NPWS
Title:	The phytosociology and conservation value of Irish sand dunes
Year:	2008
Author:	Gaynor, K.
Series:	Unpublished PhD thesis, National University of Ireland, Dublin

Title:	Saltmarsh Monitoring Report 2006
Year:	2007
Author:	McCorry, M.
Series:	Unpublished Report to NPWS
Title:	Inventory of Irish coastal lagoons
Year:	2007
Author:	Oliver, G.
Series:	Unpublished Report to NPWS
Title:	A Survey of Intertidal Mudflats and Sandflats in Ireland
Year:	2006
Author:	Aquafact
Series:	Unpublished Report to NPWS
Title:	Otter Survey of Ireland 2004/2005
Year:	2006
Author:	Bailey, M.; Rochford, J.
Series:	Irish Wildlife Manuals No. 23
Title:	Otters - ecology, behaviour and conservation
Year:	2006
Author:	Kruuk, H.
Series:	Oxford University Press
Title:	Survey of sensitive subtidal benthic marine communities
Year:	2006
Author:	MERC
Series:	Unpublished Report to NPWS
Title:	Harbour seal population assessment in the Republic of Ireland: August 2003
Year:	2004
Author:	Cronin, M.; Duck, C.; Ó Cadhla, O.; Nairn, R.; Strong, D.; O'Keeffe, C.
Series:	Irish Wildlife Manuals No. 11
Title:	Summary of National Parks & Wildlife Service surveys for common (harbour) seals (<i>Phoca vitulina</i>) and grey seals (<i>Halichoerus grypus</i>), 1978 to 2003
Year:	2004
Author:	Lyons, D.O.
Series:	Irish Wildlife Manuals No. 13
Title:	Broadscale mapping of candidate marine Special Area of Conservation. Clew Bay Complex, cSAC (001482)
Year:	2003
Author:	SSI; Aquafact
Series:	Unpublished Report to NPWS
Title:	A Survey of selected littoral and sublittoral sites in Clew Bay, Co. Mayo
Year:	1999
Author:	Aquafact
Series:	Unpublished Report to NPWS

Title:	National Shingle Beach Survey of Ireland 1999
Year:	1999
Author:	Moore, D.; Wilson, F.
Series:	Unpublished Report to NPWS
Title:	Aquatic vegetation of Irish coastal lagoons
Year:	1998
Author:	Hatch, P.; Healy, B.
Series:	Bulletin of the Irish Biogeographical Society. 21: 2-21
Title:	A survey of the vegetation of Irish coastal lagoons
Year:	1996
Author:	Hatch, P.
Series:	Unpublished Report to NPWS
Title:	The spatial organization of otters (<i>Lutra lutra</i>) in Shetland
Year:	1991
Author:	Kruuk, H.; Moorhouse, A.
Series:	J. Zool, 224: 41-57
Title:	Otter survey of Ireland
Year:	1982
Author:	Chapman, P.J.; Chapman, L.L.
Series:	Unpublished Report to Vincent Wildlife Trust
Title:	Lough Furnace, County Mayo; physical and chemical studies of an Irish saline lake, with reference to the biology of <i>Neomysis integer</i>
Year:	1977
Author:	Parker, M.M.
Series:	Unpublished PhD thesis, University of Dublin, Trinity College.

Spatial data sources

Year:	Interpolated 2011
Title:	Intertidal and subtidal surveys 1999, 2006, 2009; broadscale mapping 2003
GIS operations:	Polygon feature classes from marine community types base data sub-divided based on interpolation of marine survey data; expert opinion used as necessary to resolve any issues arising
Used for:	Marine community types, 1140 (maps 2 & 4)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	High Water Mark (HWM) polyline feature class converted into polygon feature class; clipped to SAC boundary
Used for:	1160, 1365 (maps 3 & 9)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	High water mark (HWM) and low water mark (LWM) polyline feature classes converted into polygon feature classes and combined; Saltmarsh and Sand Dune CO datasets erased out if applicable
Used for:	Marine community types base data (map 4)
Year:	Revision 2011
Title:	Inventory of Irish Coastal Lagoons. Version 3
GIS operations:	Clipped to SAC boundary
Used for:	1150 (map 5)
Year:	Revision 2010
Title:	Saltmarsh Monitoring Project 2007-2008. Version 1
GIS operations:	QIs selected; clipped to SAC boundary; overlapping regions with Sand Dune CO data investigated and resolved with expert opinion used
Used for:	1330 (map 6)
Year:	2009
Title:	Coastal Monitoring Project 2004-2006. Version 1
GIS operations:	QIs selected; clipped to SAC boundary; overlapping regions with Saltmarsh CO data investigated and resolved with expert opinion used
Used for:	1210, 2110, 2120 (map 7)
Year:	2005
Title:	OSi Discovery series vector data
GIS operations:	Creation of an 80m buffer on the marine side of the high water mark (HWM); creation of a 10m buffer on the terrestrial side of the HWM; combination of 80m and 10m HWM buffer datasets; creation of a 10m buffer on the landward side of the river banks data; creation of a 20m buffer applied to river centerline and stream data; combination of 10m river banks and 20m river and stream centerline buffer datasets; combined river and stream buffer dataset clipped to HWM; combination of HWM buffer dataset with river and stream buffer dataset; overlapping regions investigated and resolved; resulting dataset clipped to SAC boundary; expert opinion used as necessary to resolve any issues arising
Used for:	1355 (map 8)

Year:	2011
Title:	NPWS rare and threatened species database
GIS operations:	Dataset created from spatial references in database records; expert opinion used as necessary to resolve any issues arising
Used for:	1365 (map 9)

1013 Geyer's whorl snail *Vertigo geyeri*

The status of Geyer's whorl snail as a qualifying Annex II species for Clew Bay Complex SAC is currently under review. The outcome of this review will determine whether a site-specific conservation objective is set for this species.

1140 Mudflats and sandflats not covered by seawater at low tide

To maintain the favourable conservation condition of Mudflats and sandflats not covered by seawater at low tide in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 2	Habitat area was estimated using OSI data as 1277ha. See marine supporting document for further details
Community distribution	Hectares	The following sediment communities should be maintained in a natural condition: Intertidal sandy mud with <i>Tubificoides benedii</i> and <i>Pygospio elegans</i> community complex; Sandy mud with polychaetes and bivalves community complex; and Fine sand dominated by <i>Nephtys cirrosa</i> community. See map 4	The likely area of sediment communities was derived from a combination of intertidal and subtidal surveys undertaken in 1999, 2006 and 2009. See marine supporting document for further details

1150 * Coastal lagoons

To maintain the favourable conservation condition of Lagoons in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 5 for mapped lagoons	The main lagoon is Furnace Lough. Claggan Lagoon has also been mapped, however, further information is required on this lagoon. NB there maybe other lagoons within the SAC. The following targets and notes concentrate on the largest lagoon, Furnace Lough
Habitat area	Hectares	Area stable, subject to slight natural variation. Favourable reference area of surveyed lagoons is 163.3ha. Furnace Lough- 162.1ha; Claggan Lagoon- 1.2ha. See map 5	Areas calculated from spatial data derived from Oliver, 2007. NB there maybe other lagoons within the SAC
Salinity regime	Practical salinity units (psu)	Maintain current spatial and temporal variation in salinity regime	Furnace Lough is a natural, deep (up to 21m), stratified lagoon with natural periodic overturns and anoxia. It has permanent open connection to the sea through which seawater enters when tides exceed MHWN though this connection is somewhat constricted by weirs. There are major freshwater inputs at the northern end from the large Lough Feeagh/Burrishoole catchment area. The surface layer is oligohaline to mesohaline (0.5-12.0 psu) for most of the time but salinity varies from north (fresh water) to south (high salinity) and summer to winter. The waters are sharply stratified, a permanent halocline runs from 1-3m down to 8m, below which the water is of constant salinity (approx. 20psu), anaerobic and stagnant (Parker, 1977). See Oliver (2007) and Roden and Oliver (in prep.) for further information
Hydrological regime	Metres	Maintain current annual water level fluctuations	This is to ensure maintenance of the current communities of the lagoon margins and the current hydrological functioning of the lagoon itself, especially the salinity regime
Hydrological regime	Discharge (m ³ /second)	Maintain/restore freshwater discharge regime	There is evidence that the original hydrological regime in the Burrishoole catchment has been impacted due to overgrazing and afforestation resulting in changes to run-off regimes with associated increased siltation and eutrophication. The extent to which these changes have impacted on Lough Furnace is unclear but needs further study

1150 * Coastal lagoons

To maintain the favourable conservation condition of Lagoons in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Barrier	Weir function	Maintain current weir structure at Furnace Lough to ensure maintenance of the current salinity regime	In Furnace Lough, input to and output of saline water is affected to an unknown degree by two weirs. The effect of the weirs needs to be quantified to determine their effect on the salinity regime of the lagoon. These weirs or some similar type structures are shown on the first edition of the 6" OS maps and therefore have been in place for over 170 years
Water quality: chlorophyll a	µg/L	Maintain annual median chlorophyll in Furnace Lough at less than 2.5µg/L	These limits are needed to ensure that excessive shading from phytoplankton does not reduce submergent macrophytes colonisation of the littoral zone the lagoon (J. Ryan, pers comm). The current median levels are less than the target but summer levels are elevated (Roden and Oliver, in prep.) and should be closely monitored
Water quality: Molybdate Reactive Phosphorus (MRP)	mg/L	Maintain annual median MRP in Furnace Lough at less than 0.01mg/L	These limits are needed to ensure that excessive shading from phytoplankton does not reduce submergent macrophytes colonisation of the littoral zone areas of the lagoon (J. Ryan, pers comm). The current median levels in Furnace Lough are 0.005mg/L (Roden and Oliver, in prep). It is possible that the target may be exceeded during periods of overturn. Collection of data on nutrient levels close to the halocline would be useful for the assessment of this possibility
Water quality: Dissolved Inorganic Nitrogen (DIN)	mg/L	Maintain annual median DIN (Dissolved inorganic nitrogen) in Furnace Lough at less than 0.15mg/L	These limits are needed to ensure that excessive shading from phytoplankton does not reduce submergent macrophytes colonisation of the littoral zone of the lagoon (J. Ryan, pers comm). The current median levels of DIN in Furnace Lough are less than 0.1mg/L (Roden and Oliver, in prep)
Water quality: Biological Oxygen Demand (BOD)	mg/L	Maintain annual median BOD (Biological Oxygen Demand) in Furnace Lough at less than 2.0mg/L	These limits are needed to ensure that excessive shading from phytoplankton does not reduce submergent macrophytes colonisation of the littoral zone of the lagoon (J. Ryan, pers comm). The current annual median levels of BOD in Furnace Lough are just below the target (Roden and Oliver, in prep) and should be closely monitored. The relationship between organic matter, mainly peat silt, input from L. Feeagh and BOD in the surface waters and anoxia in the deeper waters warrants further investigation

1150 * Coastal lagoons

To maintain the favourable conservation condition of Lagoons in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Depth of submergent macrophyte colonisation	Metres	Maintain/increase the depth of submergent macrophyte colonisation of the lagoon	Increased depth of colonisation increases both the extent and diversity of submergent macrophytes. In comparison with similar lagoons the extent of submergent macrophyte colonisation in Furnace Lough appears to be restricted probably due to high water colour. However data on the depth of colonisation and water colour and the relationship between them is lacking. It is also possible that anoxia may be a problem, at least in some areas. These issues need to be investigated
Typical plant species	Number and m ²	Maintain number and extent of listed lagoonal specialists, subject to natural variation	Species in Furnace Lough listed in Oliver (2007), Hatch (1996) and Hatch and Healy (1998). A very limited number of plant species are currently listed for the site based on a series of shallow water transects. A snorkelling survey of this complex lagoon is required establish if that list is fully representative of the flora of the lagoon
Typical animal species	Number	Maintain listed lagoon specialists, subject to natural variation	Species in Furnace Lough listed in Oliver (2007), which rated the aquatic fauna as of moderate-high conservation value based on its high diversity and the presence of rare and unexpected crustaceans
Negative indicator species	Number and % cover	Negative indicator species absent or under control	Eutrophication would favour phytoplankton blooms at the expense of submerged macrophytes

1160 Large shallow inlets and bays

To maintain the favourable conservation condition of Large shallow inlets and bays in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	The permanent habitat area is stable or increasing, subject to natural processes. See map 3	Habitat area was estimated using OSI data as 10189ha. See marine supporting document for further details.
Community extent	Hectares	Maintain the natural extent of the <i>Zostera</i> dominated and maërl dominated communities. See map 4	The likely extent of the <i>Zostera</i> dominated and maërl dominated communities was derived from the acoustic survey and the dive survey undertaken in 2006. See marine supporting document for further details
Shoot density	Shoots per m ²	Maintain the high quality of <i>Zostera</i> dominated community	2006 diver observation and underwater viewer. See marine supporting document for further details
Community structure	Biological composition	Maintain the high quality of maërl dominated communities	Area established from an acoustic mapping survey 2003 and a 2006 diver observation and underwater viewer. See marine supporting document for further details
Community distribution	Hectares	The following communities should be maintained in a natural condition: Sandy mud with polychaetes and bivalves community complex; Fine sand dominated by <i>Nephtys cirrosa</i> community; Intertidal sandy mud with <i>Tubificoides benedii</i> and <i>Pygospio elegans</i> community complex; Shingle; and Reef. See map 4	The likely area of sediment communities was derived from a combination of acoustic mapping survey in 2003, intertidal data from 1999, 2006 and 2009 and subtidal data obtained in 1999 and 2009. See marine supporting document for further details

1210 Annual vegetation of drift lines

To maintain the favourable conservation condition of Annual vegetation of driftlines in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Bartraw - 0.04ha and Rosmurrevagh - 0.08ha. See map 7	Current area unknown. Two sub-sites (Bartraw and Rosmurrevagh) were mapped during the Coastal Monitoring Project (Ryle et al., 2009), giving a total estimated area of 0.12ha. NB further unsurveyed areas maybe present in the site. Habitat is very difficult to measure in view of its dynamic nature which means that it can appear and disappear within a site from year to year. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes	Current distribution unknown. Majority of habitat found at Bartraw and Rosmurrevagh, although there may be additional patches distributed throughout the site. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Accumulation of organic matter in tidal litter is essential for trapping sand and initiating dune formation. Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities with typical species: <i>Cakile maritima</i> , <i>Honckenya peploides</i> , <i>Salsola kali</i> and <i>Atriplex</i> spp.	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details

1220 Perennial vegetation of stony banks

To maintain the favourable conservation condition of Perennial vegetation of stony banks in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession	Current area unknown, but Clew Bay is considered to have the largest shingle reserves in the country. It was recorded from Clew Bay Complex, Bartraw and Rosmurrevagh during the National Shingle Beach Survey (Moore and Wilson, 1999), but the extent was not mapped. The Coastal Monitoring Project mapped 0.48ha of this habitat at Bartraw and 0.01ha at Rosmurrevagh (Ryle et al., 2009). The extent is considerably greater than this figure, as substantial shingle deposits are known to occur in association with many of the drumlins in Clew Bay. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes	Distribution unknown at present, although the habitat has been recorded at Clew Bay Complex (Moore and Wilson, 1999), as well as Bartraw and Rosmurrevagh (Moore and Wilson, 1999; Ryle et al., 2009). See coastal habitats supporting document for further details
Physical structure: Functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Site represents the only known example of incipient gravel barrier formation in the country. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Moore and Wilson (1999) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain the presence of species-poor communities with typical species: <i>Honckenya peploides</i> , <i>Beta vulgaris</i> ssp. <i>maritima</i> , <i>Crithmum maritimum</i> , <i>Tripleurospermum maritimum</i> , <i>Glaucium flavum</i> and <i>Silene uniflora</i>	Based on data from Moore and Wilson (1999) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Moore and Wilson (1999) and Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. See coastal habitats supporting document for further details

Conservation objectives for: Clew Bay Complex SAC [001482]

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To restore the favourable conservation condition of Atlantic salt meadows in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Mallaranny - 19.76ha, Tooreen - 1.06ha, Rosmurrevagh - 6.40ha, Tierna - 0.39ha, Rockfleet Castle - 0.37ha, Rosharnagh East - 0.03ha, Caraholly - 0.36ha, Kiladangan - 0.96ha, Annagh Island - 5.23ha, Bartraw - 0.38ha. See map 6	Based on data from the Saltmarsh Monitoring Project (McCorry, 2007). Ten sub-sites were mapped (34.94ha) and additional areas of potential saltmarsh (3.92ha) were identified for an examination of aerial photographs, giving a total estimated area of 38.86ha. NB further unsurveyed areas maybe present within the site. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 6 for known distribution	Based on data from McCorry (2007). See coastal habitats supporting document for further details
Physical structure: sediment supply	Presence/absence of physical barriers	Maintain/restore natural circulation of sediments and organic matter, without any physical obstructions	See coastal habitats backing document for further details
Physical structure: creeks and pans	Occurrence	Maintain creek and pan structure, subject to natural processes, including erosion and succession	Based on data from McCorry (2007). The efficiency of sediment circulation throughout a saltmarsh depends on the creek pattern. See coastal habitats supporting document for further details
Physical structure: flooding regime	Hectares flooded; frequency	Maintain natural tidal regime	See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from McCorry and Ryle (2009). See coastal habitats supporting document for further details
Vegetation structure: vegetation height	Centimetres	Maintain structural variation within sward	Based on data from McCorry (2007). See coastal habitats supporting document for further details
Vegetation structure: vegetation cover	Percentage cover at a representative sample of monitoring stops	Maintain more than 90% area outside creeks vegetated.	Based on data from McCorry (2007). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative sample of monitoring stops	Maintain range of sub-communities with typical species listed in Saltmarsh Monitoring Project (McCorry & Ryle, 2009)	Based on data from McCorry (2007). See coastal habitats supporting document for further details

1330 Atlantic salt meadows (*Glauco-Puccinellietalia maritimae*)

To restore the favourable conservation condition of Atlantic salt meadows in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation structure: negative indicator species - <i>Spartina anglica</i>	Hectares	No significant expansion of <i>Spartina</i> . No new sites for this species and an annual spread of less than 1% where it is already known to occur	Based on data from McCorry (2007). See coastal habitats supporting document for further details

1355 Otter *Lutra lutra*

To restore the favourable conservation condition of Otter in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. FCS target, based on 1980/81 survey findings, is 88% in SACs. Current range in west estimated at 70% (Bailey and Rochford, 2006)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 233.1ha above high water mark (HWM); 47.3ha along river banks/ around ponds	No field survey. Areas mapped to include 10m terrestrial buffer along shoreline (above HWM and along river banks) identified as critical for otters (NPWS, 2007)
Extent of marine habitat	Hectares	No significant decline. Area mapped and calculated as 2426.7ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (HWM) (NPWS, 2007; Kruuk, 2006)
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 10.2km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake/lagoon) habitat	Hectares	No significant decline. Area mapped and calculated as 141.3ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk, 2006; Kruuk and Moorhouse, 1991)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006) and wrasse and rockling in coastal waters (Kingston et al., 1999)
Barriers to connectivity	Number	No significant increase. For guidance, see map 8	Otters will regularly commute across stretches of open water up to 500m. e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed

1365 Common seal *Phoca vitulina*

To maintain the favourable conservation condition of Harbour seal in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Access to suitable habitat	Number of artificial barriers	Species range within the site should not be restricted by artificial barriers to site use	See marine supporting document for further details
Breeding behaviour	Breeding sites	The breeding sites should be maintained in a natural condition. See map 9	Attribute and target based on background knowledge of Irish breeding populations, review of data from Lyons (2004) and unpublished National Parks and Wildlife Service records. See marine supporting document for further details
Moulting behaviour	Moult haul-out sites	The moult haul-out sites should be maintained in a natural condition. See map 9	Attribute and target based on background knowledge of Irish populations, review of data from Lyons (2004), Cronin et al. (2004) and unpublished National Parks and Wildlife Service records. See marine supporting document for further details
Resting behaviour	Resting haul-out sites	The resting haul-out sites should be maintained in a natural condition. See map 9	Attribute and target based on background knowledge of Irish populations, review of data from Lyons (2004) and unpublished National Parks and Wildlife Service records. See marine supporting document for further details
Disturbance	Level of impact	Human activities should occur at levels that do not adversely affect the harbour seal population at the site	See marine supporting document for further details

2110 Embryonic shifting dunes

To restore the favourable conservation condition of Embryonic shifting dunes in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes, including erosion and succession. For sub-sites mapped: Bartraw - 0.02ha and Rosmurrevagh - 1.38ha. See map 7	Current area unknown. Two sub-sites (Bartraw and Rosmurrevagh) were mapped during the Coastal Monitoring Project (Ryle et al., 2009), giving a total estimated area of 1.40ha. NB further unsurveyed areas maybe present in the site. Habitat is very difficult to measure in view of its dynamic nature. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 7 for known distribution	Mobile dunes are well developed at Rosmurrevagh, while those at Bartraw have been compromised by the installation of coastal protection works. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: plant health of foredune grasses	Percentage cover	More than 95% of <i>Elytrigia</i> and/or <i>Leymus</i> should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover	Maintain the presence of species-poor communities with typical species: <i>Elytrigia juncea</i> and/or <i>Leymus arenarius</i>	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See coastal habitats supporting document for further details

2120 Shifting dunes along the shoreline with *Ammophila arenaria* ("white dunes")

To restore the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:


Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes including erosion and succession. For sub-sites mapped: Bartraw - 0.18ha and Rosmurrevagh - 0.36ha. See map 7	Current area unknown. Two sub-sites (Bartraw and Rosmurrevagh) were mapped during the Coastal Monitoring Project (Ryle et al., 2009), giving a total estimated area of 0.54ha. NB further unsurveyed areas may be present in the site. Habitat is very difficult to measure in view of its dynamic nature. See coastal habitats supporting document for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 7 for known distribution	Mobile dunes are well developed at Rosmurrevagh, while those at Bartraw have been compromised by the installation of coastal protection works. See coastal habitats supporting document for further details
Physical structure: functionality and sediment supply	Presence/absence of physical barriers	Maintain the natural circulation of sediment and organic matter, without any physical obstructions	Dunes are naturally dynamic systems that require continuous supply and circulation of sand. <i>Ammophila</i> reproduces vegetatively and requires constant accretion of fresh sand to maintain active growth encouraging further accretion. Physical barriers can lead to fossilisation or over-stabilisation of dunes, as well as beach starvation resulting in increased rates of erosion. See coastal habitats supporting document for further details
Vegetation structure: zonation	Occurrence	Maintain the range of coastal habitats including transitional zones, subject to natural processes including erosion and succession	Based on data from Gaynor (2008) and Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: plant health of dune grasses	Percentage cover	More than 95% of <i>Ammophila</i> and/or <i>Leymus</i> should be healthy (i.e. green plant parts above ground and flowering heads present)	Based on data from Ryle et al. (2009). See coastal habitats supporting document for further details
Vegetation composition: typical species and sub-communities	Percentage cover at a representative number of monitoring stops	Maintain the presence of species-poor communities dominated by <i>Ammophila arenaria</i> and/or <i>Leymus arenarius</i>	Based on data from Gaynor (2008) and Ryle et al. (2009). See coastal habitats supporting document for further details

2120 Shifting dunes along the shoreline with *Ammophila arenaria* ("white dunes")

To restore the favourable conservation condition of Shifting dunes along the shoreline with *Ammophila arenaria* in Clew Bay Complex SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Vegetation composition: negative indicator species	Percentage cover	Negative indicator species (including non-natives) to represent less than 5% cover	Based on data from Ryle et al. (2009). Negative indicators include non-native species, species indicative of changes in nutrient status and species not considered characteristic of the habitat. Sea-buckthorn (<i>Hippophae rhamnoides</i>) should be absent or effectively controlled. See coastal habitats supporting document for further details






Legend

SAC 001482


1140 Mudflats and sandflats not covered by sea water at low tide

OSi Discovery Series County Boundary



Legend

- SAC 001482
- 1160 Large shallow inlets and bays
- OSI Discovery Series County Boundary



*An Roinn
Ealaíon, Oidhreachta agus Gaeltachta*
*Department of
Arts, Heritage and the Gaeltacht*

MAP 3:
CLEW BAY COMPLEX
CONSERVATION OBJECTIVES
LARGE SHALLOW INLETS AND BAYS

Map to be read in conjunction with the NPWS Conservation Objectives Document.

COUNTY MAYO

0

1

2

3 km


N

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision. Reproduced from Ordnance Survey material by permission of the Government (Permit number EN 0059208).
Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithniú a déanamh ar theorainneacha na gceantar comharthaíthe. Macasamhail d'ábhar na Suirbhéarachta Ordonáis le chead ón Rialtas (Ceadúnas Uimh. EN 0059208)

SITE CODE

SAC 001482
Version 1.05

Map Version 1
Date: June 2011



Legend

SAC 001482

OSi Discovery Series County Boundary

Marine Community Types

Fine sand dominated by *Nephtys cirrosa* community

Intertidal sandy mud with *Tubificoides benedii* and *Pygospio elegans* community complex


Maerl dominated community


Reef


Sandy mud with polychaetes and bivalves community complex

Shingle

Zostera community







Legend

SAC 001482

OSi Discovery Series County Boundary

Coastal Habitats

1210 Annual vegetation of drift lines (Qualifying Interest)


1220 Perennial vegetation of stony banks (Qualifying Interest)


2110 Embryonic shifting dunes (Qualifying Interest)

2120 Shifting dunes along the shoreline with *Ammophila arenaria* ('white dunes') (Qualifying Interest)

21AO Machairs (*in Ireland)

2130 *Fixed coastal dunes with herbaceous vegetation ('grey dunes')







An Roinn
Ealaíon, Oidhreachta agus Gaeltachta

Department of
Arts, Heritage and the Gaeltacht

**Produced by: National Parks and Wildlife Service,
Department of Arts, Heritage and the Gaeltacht,
7 Ely Place, Dublin 2, Ireland.
Web: www.npws.ie
E-mail: natureconservation@environ.ie**

Citation:

NPWS (2011) Conservation Objectives: Clew Bay Complex SAC 001482. Version 1.0. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht.

Series Editors: Rebecca Jeffrey & Naomi Kingston
ISSN 2009-4086