# **National Parks and Wildlife Service**

**Conservation Objectives Series** 

## Croaghaun/Slievemore SAC 001955



An Roinn Tithíochta, Rialtais Áitiúil agus Oidhreachta Department of Housing, Local Government and Heritage National Parks and Wildlife Service, Department of Housing, Local Government and Heritage,

90 King Street North, Dublin 7, D07 N7CV, Ireland.

Web: www.npws.ie E-mail: natureconservation@housing.gov.ie

Citation:

NPWS (2021) Conservation Objectives: Croaghaun/Slievemore SAC 001955. Version 1. National Parks and Wildlife Service, Department of Housing, Local Government and Heritage.

### Series Editors: Rebecca Jeffrey and Christina Campbell ISSN 2009-4086

#### Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance
- exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

• population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and

• the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and

• there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

#### **Notes/Guidelines:**

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.

2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.

3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.

4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.

5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

#### **Qualifying Interests**

#### \* indicates a priority habitat under the Habitats Directive

001955	S Croaghaun/Slievemore SAC		
4010	Northern Atlantic wet heaths with Erica tetralix		
4030	European dry heaths		
4060	Alpine and Boreal heaths		
7130	Blanket bogs (* if active bog)		
8110	Siliceous scree of the montane to snow levels (Androsacetalia alpinae and		

- 8110 Siliceous scree of the montane to snow levels (Androsacetalia alpinae and Galeopsietalia ladani)
- 8220 Siliceous rocky slopes with chasmophytic vegetation

Please note that this SAC is adjacent to Achill Head SAC (002268). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent site as appropriate.

### Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

#### **NPWS Documents**

Year :	2009		
Title :	Ireland Red List No. 2: Non-marine molluscs		
Author :	Byrne, A.; Moorkens, E.A.; Anderson, R.; Killeen, I.J.; Regan, E.C.		
Series :	Ireland Red List series, NPWS		
Year :	2010		
Title :	Ireland Red List No. 4: Butterflies		
Author :	Regan, E.C.; Nelson, B.; Aldwell, B.; Bertrand, C.; Bond, K.; Harding, J.; Nash, D.; Nixon, D.; Wilson, C.J.		
Series :	Ireland Red List series, NPWS		
Year :	2012		
Title :	Ireland Red List No. 8: Bryophytes		
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.		
Series :	Ireland Red List series, NPWS		
Year :	2014		
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0		
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.		
Series :	Irish Wildlife Manuals, No. 79		
Year :	2014		
Title :	National Survey of Upland Habitats (Phase 1, 2010-2011), Site Report No. 6: Croaghaun/Slievemore cSAC (001955), Co. Mayo (Revision)		
Author :	Roche, J.R.; Perrin, P.M.; Barron, S.J.; Daly, O.H.		
Series :	Unpublished report to NPWS		
Year :	2016		
Title :	Ireland Red List No. 10: Vascular Plants		
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.		
Series :	Ireland Red Lists series, NPWS		
Year :	2016		
Title :	Survey of Flora Protection Order Bryophytes 2016		
Author :	Hodd, R.L.		
Series :	Unpublished report to NPWS		
Year :	2021		
Title :	Croaghaun/Slievemore SAC (site code: 1955) Conservation objectives supporting document - upland habitats V1		
Author :	NPWS		
Series :	Conservation objectives supporting document		

#### **Other References**

Year :	2012
Title :	Rare and threatened bryophytes of Ireland
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	National Museums Northern Ireland

Year :	2017
Title :	Irish Vegetation Classification: Technical Progress Report No. 3
Author :	Perrin, P.
Series :	Report submitted to National Biodiversity Data Centre

Page 6 of 18

## Spatial data sources

Year :	2011
Title :	National Survey of Upland Habitats
GIS Operations :	Habitat dataset for site clipped to SAC boundary. Relevant QI selected and exported to new dataset. Expert opinion used as necessary to resolve any issues arising
Used For :	4010, 4030, 4060, 7130, 8110, 8220 (maps 3, 4, 5, 6, 7, 8)

#### 4010 Northern Atlantic wet heaths with Erica tetralix

To restore the favourable conservation condition of Northern Atlantic wet heaths with *Erica tetralix* in Croaghaun/Slievemore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes	Croaghaun/Slievemore SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 and Perrin et al., 2014). Northern Atlantic wet heaths with <i>Erica tetralix</i> was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 1,411.8ha; it was recorded as the most extensive Annex I habitat, covering 40.4% of the SAC. Roche et al. (2014) report obvious losses of the habitat since 1995 of approximately 0.55ha. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Croaghaun/Slievemore SAC conservation objectives supporting document for upland habitats where a summary of the mapping methodology and a brief discussion of restoration potential are also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 3	Wet heath was recorded throughout Croaghaun/Slievemore SAC by Roche et al. (2014), notably at the lower slopes of Slievemore, a large swathe running along the saddle between Slievemore and Croaghaun to the upper slopes of Croaghaun, and, on the western side of Croaghaun, wet heath also dominates the slopes north-west of Keem Bay. The habitat also occurs at the low-lying land at Saddle Head. See Roche et al. (2014) for further information. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	See the uplands supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded six different wet heath communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on the vegetation communities associated with this habitat is presented in Perrin et al. (2014). See also the Irish Vegetation Classification (Perrin, 2017; www.biodiversityireland.ie/projects/ivc-classification- explorer/)
Vegetation composition: cross-leaved heath	Occurrence within 20m of a representative number of 2m x 2m monitoring stops	Cross-leaved heath ( <i>Erica tetralix</i> ) present within a 20m radius of each monitoring stop	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is presented. Further details can be found in the uplands supporting document
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of <i>Cladonia</i> and <i>Sphagnum</i> species, <i>Racomitrium lanuginosum</i> and pleurocarpous mosses at least 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: ericoid species and crowberry	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of ericoid species and crowberry ( <i>Empetrum</i> <i>nigrum</i> ) at least 15%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details

Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrubs less than 75%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is presented. See the uplands supporting document for further details
Vegetation composition: non- native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details. The non-native moss <i>Campylopus</i> <i>introflexus</i> and rhododendron ( <i>Rhododendron</i> <i>ponticum</i> ) were recorded by Roche et al. (2014) within this habitat in the SAC
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken ( <i>Pteridium aquilinum</i> ) less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush ( <i>Juncus effusus</i> ) less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids, crowberry ( <i>Empetrum nigrum</i> ) and bog-myrtle ( <i>Myrica gale</i> ) showing signs of browsing	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas for this habitat is presented. See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: drainage	Percentage cover in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species on the Flora (Protection) Order, 2015 and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH. Roche et al. (2014) recorded hepatic mats of the <i>Calluna</i> <i>vulgaris-Herbertus aduncus</i> community within this habitat during the NSUH. See the uplands supporting document for further details

#### 4030 European dry heaths

To restore the favourable conservation condition of European dry heaths in Croaghaun/Slievemore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Croaghaun/Slievemore SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 and Perrin et al., 2014). European dry heaths was mapped in detail and the total area of the qualifying habitat as stated by Roche et al. (2014) is 301.7ha, covering 8.6% of the SAC. Roche et al. (2014) report no net loss of area of the habitat in the SAC since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Croaghaun/Slievemore SAC conservation objectives supporting document for upland habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 4	Dry heath was recorded by Roche et al. (2014) throughout Croaghaun/Slievemore SAC, notably around the summit of Slievemore, on the very high and steep western slopes of Tonacroaghaun and at the low-lying land at Saddle Head. See Roche et al. (2014) for further information. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	See the uplands supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded four different dry heath communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on the vegetation communities associated with this habitat is presented in Perrin et al. (2014). See also the Irish Vegetation Classification (Perrin, 2017; www.biodiversityireland.ie/projects/ivc-classification- explorer/)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop at least three, excluding <i>Campylopus</i> and <i>Polytrichum</i> mosses	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: number of positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species present at each monitoring stop at least two	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat, which is composed of dwarf shrubs, is presented. See the uplands supporting document for further details
Vegetation composition: cover of positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 50% for siliceous dry heath and 50- 75% for calcareous dry heath	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat, which is composed of dwarf shrubs, is presented. See the uplands supporting document for further details
Vegetation composition: dwarf shrub composition	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of dwarf shrub cover composed collectively of bog-myrtle ( <i>Myrica gale</i> ), creeping willow ( <i>Salix repens</i> ) and western gorse ( <i>Ulex gallii</i> ) is less than 50%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details

Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species is presented. See the uplands supporting document for further details
Vegetation composition: non- native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details. The non-native moss <i>Campylopus</i> <i>introflexus</i> and rhododendron ( <i>Rhododendron</i> <i>ponticum</i> ) were recorded by Roche et al. (2014) within this habitat in the SAC
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 20%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: bracken	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of bracken ( <i>Pteridium aquilinum</i> ) less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: soft rush	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of soft rush ( <i>Juncus effusus</i> ) less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: senescent ling	Percentage cover at a representative number of 2m x 2m monitoring stops	Senescent proportion of ling ( <i>Calluna vulgaris</i> ) cover less than 50%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids showing signs of browsing	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas is presented. See the uplands supporting document for further details
Vegetation structure: growth phases of ling	Percentage cover in local vicinity of a representative number of monitoring stops	Outside sensitive areas, all growth phases of ling ( <i>Calluna vulgaris</i> ) should occur throughout, with at least 10% of cover in the mature phase	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas is presented. See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species on the Flora (Protection) Order, 2015 (FPO) and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC, including the FPO listed Northern Atlantic hepatic mat species <i>Adelanthus</i> <i>lindenbergianus, Scapania ornithopodioides</i> and <i>S.</i> <i>nimbosa</i> , all Vulnerable in Ireland (Lockhart et al., 2012), and the Near Threatened <i>Mastigophora</i> <i>woodsii</i> (Lockhart et al., 2012), all of which were subsequently recorded in association with the habitat in the SAC by Hodd (2016). Roche et al. (2014) recorded hepatic mats of the <i>Calluna</i> <i>vulgaris-Herbertus aduncus</i> community within this habitat and <i>M. woodsii</i> was recorded during the NSUH. Any new records should also be considered within this attribute. See the uplands supporting document for further details

#### 4060 Alpine and Boreal heaths

To restore the favourable conservation condition of Alpine and Boreal heaths in Croaghaun/Slievemore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Croaghaun/Slievemore SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 and Perrin et al., 2014). Alpine and Boreal heath was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 268.2ha, covering 7.7% of the SAC. Roche et al. (2014) report no obvious significant losses of area since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Croaghaun/Slievemore SAC conservation objectives supporting document for upland habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 5	Alpine and Boreal heath was recorded by Roche et al. (2014) particularly around the summit of Slievemore on the eastern side of Croaghaun. See Roche et al. (2014) for further information. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	See the uplands supporting document for further details
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded four Alpine and Boreal heath communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on the vegetation communities associated with this habitat is presented in Perrin et al. (2014). See also the Irish Vegetation Classification (Perrin, 2017; www.biodiversityireland.ie/projects/ivc-classification- explorer/)
Vegetation composition: lichens and bryophytes	Number of species at a representative number of 2m x 2m monitoring stops	Number of bryophyte or non-crustose lichen species present at each monitoring stop at least three	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: positive indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of positive indicator species at least 66%	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is presented. See the uplands supporting document for further details
Vegetation composition: dwarf shrub species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of dwarf shrub species at least 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 10%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species is presented. See the uplands supporting document for further details
Vegetation composition: non- native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details. No non-native species were recorded within this habitat by Roche et al. (2014)
Vegetation structure: signs of grazing	Percentage of leaves grazed at a representative number of 2m x 2m monitoring stops	Less than 10% collectively of the live leaves of specific graminoids showing signs of grazing	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details, including the list of specific graminoids

Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Less than 33% collectively of the last complete growing season's shoots of ericoids and crowberry ( <i>Empetrum nigrum</i> ) showing signs of browsing	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning within the habitat	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat and no decline in status of hepatic mats associated with this habitat	This includes species on the Flora (Protection) Order, 2015 (FPO) and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC, including the FPO listed Northern Atlantic hepatic mat species <i>Scapania</i> <i>ornithopodioides</i> and <i>S. nimbosa</i> , both Vulnerable in Ireland (Lockhart et al., 2012), and the Near Threatened <i>Mastigophora woodsii</i> (Lockhart et al., 2012), all of which were subsequently recorded in association with the habitat in the SAC by Hodd (2016). Hepatic mats of the <i>Calluna vulgaris</i> - <i>Herbertus aduncus</i> community and the Near Threatened dwarf willow ( <i>Salix herbacea</i> ) (Wyse Jackson et al., 2016) were recorded within this habitat during the NSUH (Roche et al., 2014). New records should also be considered within this attribute. See the uplands supporting document for further details

Page 13 of 18

#### 7130 Blanket bogs (\* if active bog)

To restore the favourable conservation condition of Blanket bogs (\* if active bog) in Croaghaun/Slievemore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area increasing, subject to natural processes	Croaghaun/Slievemore SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 and Perrin et al., 2014). Roche et al. (2014) state that the total area of blanket bog is 218.8ha (6.3% of the SAC). The comprises 212.5ha of active blanket bog and 6.3ha of inactive blanket bog. Roche et al. (2014) also report obvious losses of habitat since 1995 of approximately 3.6ha. However, this is almost certainly an underestimate, as chronic losses due to erosion since 1995 cannot be quantified (32.3ha were mapped as eroding blanket bog by Roche et al., 2014). It should be noted also that further restoration of blanket bog would be required in order to fulfil the targets for peat formation and hydrology presented below. See the Croaghaun/Slievemore SAC conservation objectives supporting document for upland habitats for further details
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 6	Areas of blanket bog were recorded by Roche et al. (2014) throughout Croaghaun/Slievemore SAC. See Roche et al. (2014) for further information. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	See the uplands supporting document for further details
Ecosystem function: peat formation	Active blanket bog as a proportion of the total area of Annex I blanket bog habitat	At least 99% of the total Annex I blanket bog area is active	From the areas given by Roche et al. (2014) above, 97.1% of the Annex I blanket bog habitat is currently active. See the uplands supporting document for further details
Ecosystem function: hydrology	Flow direction, water levels, occurrence of drains and erosion gullies	Natural hydrology unaffected by drains and erosion	Further details and a brief discussion of restoration potential is presented in the uplands supporting document
Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	Roche et al. (2014) recorded seven different active blanket bog communities within this SAC. Data on the abundance of these communities is reproduced in the uplands supporting document. Further information on the vegetation communities associated with this habitat is presented in Perrin et al. (2014). See also the Irish Vegetation Classification (Perrin, 2017; www.biodiversityireland.ie/projects/ivc-classification- explorer/)
Vegetation composition: positive indicator species	Number of species at a representative number of 2m x 2m monitoring stops	Number of positive indicator species at each monitoring stop at least seven	Attribute and target based on Perrin et al. (2014), where the list of positive indicator species for this habitat is presented. See the uplands supporting document for further details
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes or lichens, excluding <i>Sphagnum fallax</i> , at least 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: potential dominant species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of each of the potential dominant species less than 75%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details, including the list of potentially dominant species
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Total cover of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species is presented. See the uplands supporting document for further details

17	Dec 2021	1
	DUULULI	

Page 14 of 18

Vegetation composition: non- native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details. The non-native moss <i>Campylopus</i> <i>introflexus</i> was recorded within this habitat in the SAC by Roche et al. (2014)
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: <i>Sphagnum</i> condition	Condition at a representative number of 2m x 2m monitoring stops	Less than 10% of the <i>Sphagnum</i> cover is crushed, broken and/or pulled up	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: signs of browsing	Percentage of shoots browsed at a representative number of 2m x 2m monitoring stops	Last complete growing season's shoots of ericoids, crowberry ( <i>Empetrum</i> <i>nigrum</i> ) and bog-myrtle ( <i>Myrica gale</i> ) showing signs of browsing collectively less than 33%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: burning	Occurrence in local vicinity of a representative number of monitoring stops	No signs of burning in sensitive areas, into the moss, liverwort or lichen layer or exposure of peat surface due to burning	Attribute and target based on Perrin et al. (2014), where the list of sensitive areas is presented. See the uplands supporting document for further details
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: drainage	Percentage area in local vicinity of a representative number of monitoring stops	Area showing signs of drainage from heavy trampling, tracking or ditches less than 10%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Physical structure: erosion	Percentage area in local vicinity of a representative number of monitoring stops	Less than 5% of the greater bog mosaic comprises erosion gullies and eroded areas	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species on the Flora (Protection) Order, 2015 and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH. No relevant species were recorded in this habitat during the NSUH; however, new records should be considered within this attribute. See the uplands supporting document for further details

## 8110 Siliceous scree of the montane to snow levels (Androsacetalia alpinae and Galeopsietalia ladani)

To maintain the favourable conservation condition of Siliceous scree of the montane to snow levels (Androsacetalia alpinae and Galeopsietalia ladani) in Croaghaun/Slievemore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Croaghaun/Slievemore SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 and Perrin et al., 2014). Siliceous scree of the montane to snow levels (Androsacetalia alpinae and Galeopsietalia ladani) was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 69.8ha, covering 2.1% of the SAC. Roche et al. (2014) report no significant losses of area of the habitat in the SAC since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Croaghaun/Slievemore SAC conservation objectives supporting document for upland habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 7	Siliceous scree was recorded by Roche et al. (2014) throughout the SAC on steeper ground. See Roche et al. (2014) for further information. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	See the uplands supporting document for further details
Vegetation composition: lichens and bryophytes	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of bryophytes and non-crustose lichen species at least 5%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of vegetation composed of negative indicator species less than 1%	Attribute and target based on Perrin et al. (2014), where the list of negative indicator species for this habitat is presented. See the uplands supporting document for further details
Vegetation composition: non- native species	Percentage cover at a representative number of 2m x 2m monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details. No non-native species were recorded within this habitat by Roche et al. (2014)
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	At least one positive indicator species present in vicinity of each monitoring stop in block scree	Attribute and target based on Perrin et al. (2014). The list of positive indicator species for this habitat is also presented in Perrin et al. (2014) and is the same as for 8220 Siliceous rocky slopes. Further details can be found in the uplands supporting document
Vegetation composition: grass species and dwarf shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of grass species and dwarf shrubs less than 20%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken ( <i>Pteridium aquilinum</i> ), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: grazing and browsing	Percentage of leaves/ shoots grazed/browsed at a representative number of 2m x 2m monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details

Indicators of local Occurr distinctiveness popula	n x 2m monitoring	scree running or vehicles less than 10%	details
	rrence and lation size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species on the Flora (Protection) Order, 2015 and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC and added any new records collected during the NSUH. Any new records should be considered within this attribute. See the uplands supporting document for further details

#### 8220 Siliceous rocky slopes with chasmophytic vegetation

To maintain the favourable conservation condition of Siliceous rocky slopes with chasmophytic vegetation in Croaghaun/Slievemore SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Croaghaun/Slievemore SAC was surveyed as part of the National Survey of Upland Habitats (NSUH; see Roche et al., 2014 and Perrin et al., 2014). Siliceous rocky slopes with chasmophytic vegetation was mapped in detail for the SAC and the total area of the qualifying habitat stated by Roche et al. (2014) is 38ha, covering 1.2% of the SAC. Roche et al. (2014) report no significant losses of area of the habitat in the SAC since 1995. Further information can be found in Roche et al. (2014). Further details on this and the following attributes can be found in the Croaghaun/Slievemore SAC conservation objectives supporting document for upland habitats where a summary of the mapping methodology is also presented
Habitat distribution	Occurrence	No decline, subject to natural processes. See map 8	Siliceous rocky slopes habitat was recorded by Roche et al. (2014) on steeper ground throughout the SAC. See Roche et al. (2014) for further information. A summary of the mapping methodology is presented in the uplands supporting document
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	See the uplands supporting document for further details
Vegetation composition: positive indicator species	Number of species in local vicinity of a representative number of monitoring stops	At least one positive indicator species present in vicinity of each monitoring stop	Attribute and target based on Perrin et al. (2014). The list of positive indicator species for this habitat is also presented in Perrin et al. (2014) and is the same as for 8110 Siliceous screes. Further details can be found in the uplands supporting document
Vegetation composition: non- native species	Percentage cover in local vicinity of a representative number of monitoring stops	Proportion of vegetation composed of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation composition: bracken, native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of bracken ( <i>Pteridium aquilinum</i> ), native trees and shrubs less than 25%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Vegetation structure: grazing and browsing	Percentage of leaves/ shoots grazed/browsed in local vicinity of a representative number of monitoring stops	Live leaves of forbs and shoots of dwarf shrubs showing signs of grazing or browsing collectively less than 50%	Attribute and target based on Perrin et al. (2014). See the uplands supporting document for further details
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat	This includes species on the Flora (Protection) Order, 2015 and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.). Roche et al. (2014) compiled and mapped existing rare and notable plant records for the SAC, including the FPO listed and Vulnerable liverwort <i>Acrobolbus wilsonii</i> (Lockhart et al., 2012) which was subsequently refound by Hodd (2016) in association with the habitat in the SAC. No relevant species were recorded in this habitat during the NSUH; however, any new records should be considered within this attribute. See the uplands supporting document for further details

Page 18 of 18















