National Parks and Wildlife Service

Conservation Objectives Series

Kildun Souterrain SAC 002320



An Roinn Cultúir, Oidhreachta agus Gaeltachta Department of Culture, Heritage and the Gaeltacht National Parks and Wildlife Service, Department of Culture, Heritage and the Gaeltacht,

90 King Street North, Dublin 7, D07 N7CV, Ireland.

Web: www.npws.ie E-mail: nature.conservation@chg.gov.ie

Citation:

NPWS (2018) Conservation Objectives: Kildun Souterrain SAC 002320. Version 1. National Parks and Wildlife Service, Department of Culture, Heritage and the Gaeltacht.

> Series Editor: Rebecca Jeffrey ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance
- exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

• population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and

• the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and

• there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.

2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.

3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.

4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.

5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

1303 Lesser Horseshoe Bat Rhinolophus hipposideros

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2018
Title :	Conservation objectives supporting document – lesser horseshoe bat (<i>Rhinolophus hipposideros</i>)
Author :	NPWS
Series :	Conservation objectives supporting document

Other References

Year :	2007								
Title :	Protecting and managing underground sites for bats								
Author :	Mitchell-Jones, A.J.; Bihari, Z.; Masing, M.; Rodrigues, L.								
Series :	EUROBATS Publication Series No. 2								
Year :	2008								
Title :	The lesser horseshoe bat conservation handbook								
Author :	Schofield, H.W.								
Series :	The Vincent Wildlife Trust								
Year :	2009								
Title :	Importance of night roosts for bat conservation: roosting behaviour of the lesser horseshoe bat <i>Rhinolophus hipposideros</i>								
Author :	Knight, T.; Jones, G.								
Series :	Endangered Species Research, 8: 79-86								

Spatial data sources

Year :	2018						
Title :	NPWS lesser horseshoe bat database						
GIS Operations :	Roost identified, clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising						
Used For :	1303 (map 2)						
Year :	2007						
Year : Title :	2007 Forest Inventory and Planning System (FIPS)						

Conservation Objectives for : Kildun Souterrain SAC [002320]


1303 Lesser Horseshoe Bat *Rhinolophus hipposideros*

l

To restore the favourable conservation condition of Lesser Horseshoe Bat in Kildun Souterrain SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes				
Population per roost	Number	Minimum number of 50 bats for the winter roost (roost id. 676 in NPWS database). See map 2	A figure of 100 bats for summer roosts and 50 bats for winter roosts was set as a minimum qualifying standard (MQS) when SACs were being selected for lesser horseshoe bat (<i>Rhinolophus hipposideros</i>). NPWS conduct annual counts at each qualifying roost. Qualified means from the 2013-2017 winter data have been calculated whereby the year with the highest maximum count and the year with the lowest maximum count were removed and the mea of the remaining years was calculated. This mean is usually set as the target figure for the roost. However, in the case of the winter roost (roost id. 676 in NPWS database) in Kildun Souterrain SAC, where a mean of 15 bats was recorded (2013- 2017), the target is instead set at the MQS of 50 bats. See the conservation objectives supporting document for lesser horseshoe bat (NPWS, 2018) for further information on all attributes and targets				
Winter roosts	Condition	No decline	Kildun Souterrain SAC has been selected for lesser horseshoe bat because of the presence of one internationally important winter roost (roost id. 676 in NPWS database). Damage or disturbance to the roost or to the habitat immediately surrounding it will lead to a decline in its condition (Mitchell-Jones et al., 2007)				
Auxiliary roosts	Number and condition	No decline	Lesser horseshoe bat populations will use a variety of roosts during the year besides the main summer maternity and winter hibernation roosts. Such additional roosts within the SAC may be important as night roosts, satellite roosts, etc. Night roosts ar also considered an integral part of core foraging areas and require protection (Knight and Jones, 2009). In addition, in response to weather conditions for example, bats may use different seasonal roosts from year to year; this is particular noticeable in winter. A database of all known lesses horseshoe bat roosts is available on the National Biodiversity Data Centre website. NB further unrecorded roosts may also be present within this SAC				
Extent of potential Hectares foraging habitat		No significant decline within 2.5km of qualifying roost	Lesser horseshoe bats normally forage in woodlands/scrub within 2.5km of their roosts (Schofield, 2008). See map 2 which shows a 2.5km zone around the above roost and identifies potentia foraging grounds				
Linear features Kilometres		No significant loss within 2.5km of qualifying roost. See map 2	This species follows commuting routes from its roos to its foraging grounds. Lesser horseshoe bats will not cross open ground. Consequently, linear features such as hedgerows, treelines and stone walls provide vital connectivity for this species within 2.5km around each roost (Schofield, 2008)				
Light pollution Lux		No significant increase in artificial light intensity adjacent to named roost or along commuting routes within 2.5km of the roost. See map 2	Lesser horseshoe bats are very sensitive to light pollution and will avoid brightly lit areas. Inappropriate lighting around roosts may cause abandonment; lighting along commuting routes ma cause preferred foraging areas to be abandoned, thus increasing energetic costs for bats (Schofield, 2008)				

Version 1



9	94	9	91	9	\sim	S			1	9	91	9
			9					4			91	9
					9			91				9
		91			91			91			91	9.0-
												6
91		9				91	1	91				91
91										Y L Y L F L		9 4 9 4 9 4
9 L	9 4 9 A					9 $_{1}$						9
91		91				91		27				
		9	9	9	9							9
		1			9 1						91	9
		¥	4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4							9	9	9
		Ł			91	9		1			91	9.4
		1	11									9
Ł	4 h					9 1						9
91						91		91			91	9
												9
91			9		91	9 1		91				91
			9									9 L
9 1		9			9	9 1		9 L			9 <u>1</u>	9
						91		9				1
						91		91				T
91		91			9. 1	9 1		91			91	9

SAC 002320 CO. MAYO 😽



