

National Parks and Wildlife Service

Conservation Objectives Series

Lough Derravaragh SPA 004043



NPWS

An tSeirbhís Páirceanna
Náisiúnta agus Fiadhúlra
National Parks and Wildlife
Service

**National Parks and Wildlife Service,
Department of Housing, Local Government and Heritage,
90 King Street North, Dublin 7, D07 N7CV, Ireland.
Web: www.npws.ie
E-mail: natureconservation@npws.gov.ie**

Citation:

**NPWS (2024) Conservation Objectives: Lough Derravaragh SPA 004043. Version
1. National Parks and Wildlife Service, Department of Housing, Local
Government and Heritage.**

**Series Editors: Maria Long and Colin Heaslip
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

004043	Lough Derravaragh SPA
--------	-----------------------

A038	Whooper Swan <i>Cygnus cygnus</i>
------	-----------------------------------

A059	Pochard <i>Aythya ferina</i>
------	------------------------------

A061	Tufted Duck <i>Aythya fuligula</i>
------	------------------------------------

A125	Coot <i>Fulica atra</i>
------	-------------------------

A999	Wetlands
------	----------

Please note that this SPA is adjacent to Garriskil Bog SAC (000679). See map 2. The conservation objectives for this site should be used in conjunction with those for the adjacent site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2013
Title :	A review of the SPA network of sites in the Republic of Ireland
Author :	NPWS
Series :	Unpublished Report
Year :	2019
Title :	Irish wetland bird survey: waterbird status and distribution 2009/10-2015/16
Author :	Lewis, L. J.; Burke, B.; Fitzgerald, N.; Tierney, T. D.; Kelly, S.
Series :	Irish Wildlife Manuals No. 106

Other References

Year :	1995
Title :	Impacts of hunting disturbance on waterbirds - a review
Author :	Madsen, J.; Fox, A.D.
Series :	Wildlife Biology 1(4):193-207
Year :	2016
Title :	Assessing connectivity with Special Protection Areas (SPAs)
Author :	Scottish Natural Heritage
Series :	Guidance Series Version 3 - June 2016
Year :	2018
Title :	Habitat and species mediated short and long-term distributional changes in waterbird abundance linked to variation in European winter weather
Author :	Pavón-Jordán, D.; Clausen, P.; Dagys, M.; Devos, K.; Encarnação, V.; Fox, A.D.; Frost, T.; et al.
Series :	Diversity and Distributions, 1-15
Year :	2021
Title :	Population size, breeding success and habitat use of whooper swan <i>Cygnus cygnus</i> and Bewick's Swan <i>Cygnus columbianus bewickii</i> in Ireland: results of the 2020 international swan census
Author :	Burke, B.; McElwaine, J.G.; Fitzgerald, N.; Kelly, S.B.A.; McCulloch, N.; Walsh, A.J.; Lewis, L.J.
Series :	Irish Birds 43:57-70
Year :	2022
Title :	Irish wetland bird survey: I-WeBS national and site trends report 1994/95 – 2019/20
Author :	Kennedy, J.; Burke, B.; Fitzgerald, N.; Kelly, S.B.A.; Walsh, A.J.; Lewis, L.J.
Series :	https://birdwatchireland.ie/app/uploads/2022/04/iwebs_trends_report.html

Conservation Objectives for : Lough Derravaragh SPA [004043]

A038

Whooper Swan *Cygnus cygnus*

To restore the Favourable conservation condition of Whooper Swan in Lough Derravaragh SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Winter population trend	Percentage change in number of individuals	Long term winter population trend is stable or increasing	The national population of Whooper Swan wintering in Ireland has increased in the long term, with a 40% population increase from 1991 - 2015 (Lewis et al., 2019). During the baseline assessments to inform SPA designation, 102 Whooper Swan were estimated to be using this SPA (5 year mean of peak counts for baseline period 1995/96 - 1999/2000; see NPWS, 2013). A population of 50 Whooper Swan was estimated to be using the Lough Derravaragh SPA in recent years (5 year mean of peak counts from the Irish Wetland Bird Survey, I-WeBS, monitoring for the period 2017/18 - 2021/22). Note that the 2020/21 peak count was derived from the 2020 International Swan Census (Burke et al., 2021). This represents an estimated population decrease of 51% since the baseline period which is contrary to the national trend
Winter spatial distribution	Hectares, time and intensity of use	Sufficient number of locations, area, and availability (in terms of timing and intensity of use) of suitable habitat to support the population target	Distribution encapsulates the number of locations and area of potentially suitable habitat for the wintering population and its availability for use. The suitability and availability of habitat areas is likely to vary throughout the season, for example, due to variation in land management practices or the abundance of resources available (due to natural variation and other factors). This will affect the spatio-temporal patterns of use of the habitats by the wintering population
Disturbance at wintering site	Intensity, frequency, timing and duration	The intensity, frequency, timing and duration of disturbance occurs at levels that do not significantly impact the achievement of targets for population trend and spatial distribution	The impact of any significant disturbance (direct or indirect) to the wintering population will ultimately affect the achievement of targets for population trend and/or spatial distribution. Disturbance contributes to increased energetic expenditure which can result in increased likelihood of winter mortality or reduced fitness (if energy expenditure is greater than energy gain), which can negatively impact population trends (see, for example, Madsen and Fox, 1995). Factors such as intensity, frequency, timing and duration of a (direct or indirect) disturbance source must be taken into account to determine the potential impact upon the targets for population trend and spatial distribution
Barriers to connectivity and site use	Number, location, shape and hectares	Barriers do not significantly impact the wintering population's access to the SPA or other ecologically important sites outside the SPA	Barriers limiting the population's access to this SPA or ecologically important sites outside the SPA will ultimately affect the achievement of targets for population trend and/or spatial distribution. Factors such as the number, location, shape and area of potential barriers must be taken into account to determine their potential impact. Access to ecologically important sites outside the SPA must also be considered as a single SPA may not satisfy all the ecological requirements of the wintering population, and it may require access to other SPAs or sites for certain activities, such as foraging when preferred foraging areas are unavailable due to disturbance, extensive flooding, or other factors

Forage spatial distribution, extent and abundance	Location, hectares, and forage biomass	Sufficient number of locations, area of suitable habitat and available forage biomass to support the population target	This species feeds on a wide range of aquatic and terrestrial vegetation. Key forage materials include: leaves, with significant consumption of grasses; seeds, including spilled grain; roots; tubers, including potatoes; shoots, including those from winter wheat and other cereals. Key foraging habitats are grasslands (including wet grassland, semi-improved grassland, and intensive grassland), arable stubble, winter cereals, rivers, lakes, turloughs and other wetland habitats. In general, the foraging distance of wintering Whooper Swan from night roosts is estimated to be less than 5km (Scottish Natural Heritage, 2016), although this will vary depending on site and landscape
Roost spatial distribution and extent	Location and hectares of roosting habitat	Sufficient number of locations, area and availability of suitable roosting habitat to support the population target	Overnight roosting habitat consists primarily of permanent waterbodies, such as rivers, lakes, turloughs, lagoons and other open waterbodies. Roosting is a critical ecological requirement for the wintering population. Daytime roosting is also a common behaviour, where birds minimise activity levels to conserve energy, while benefitting from the vigilance of other flock members. A lack of sufficient and suitable roosting habitats can result in increased mortality risk, whether indirectly (e.g. via increased energy expenditure travelling to/from roost sites) or directly (e.g. via increased predation risk), or reduction in site use; this would ultimately affect the achievement of targets for population trend and/or spatial distribution
Supporting habitat: area and quality	Hectares and quality	Sufficient area of utilisable habitat available in ecologically important sites outside the SPA	The wintering population can make extensive use of suitable habitats in important areas outside the SPA for foraging and roosting. The extent, availability and quality of these supporting habitats may be of importance for the resilience of the SPA population. Suitable supporting habitats include those highlighted in the attributes for foraging and roosting habitat

Conservation Objectives for : Lough Derravaragh SPA [004043]

A059 Pochard *Aythya ferina*

To restore the Favourable conservation condition of Pochard in Lough Derravaragh SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Winter population trend	Percentage change in number of individuals	Long term winter population trend is stable or increasing	The national population of wintering Pochard in Ireland has decreased by 79% from 1994/95 - 2019/20, as monitored via I-WeBS (Kennedy et al., 2022). During the baseline assessments to inform SPA designation, 3,129 Pochard were estimated to be using this SPA (5 year mean of peak counts for baseline period 1995/96 - 1999/00; see NPWS, 2013). A population of 111 Pochard was estimated to be using the Lough Derravaragh SPA in recent years (5 year mean of peak counts from I-WeBS monitoring for the period 2017/18 - 2021/22). This represents an estimated population decrease of 97% since the baseline period which is greater than the national trend. Population declines are likely linked, at least in part, to distribution shifts driven by climate change (see Pavón-Jordan et al., 2018)
Winter spatial distribution	Hectares, time and intensity of use	Sufficient number of locations, area, and availability (in terms of timing and intensity of use) of suitable habitat to support the population target	Distribution encapsulates the number of locations and area of potentially suitable habitat for the wintering population and its availability for use. The suitability and availability of habitat areas is likely to vary throughout the season, for example, due to variation in land management practices or the abundance of resources available (due to natural variation and other factors). This will affect the spatio-temporal patterns of use of the habitats by the wintering population
Disturbance at wintering site	Intensity, frequency, timing and duration	The intensity, frequency, timing and duration of disturbance occurs at levels that do not significantly impact the achievement of targets for population trend and spatial distribution	The impact of any significant disturbance (direct or indirect) to the wintering population will ultimately affect the achievement of targets for population trend and/or spatial distribution. Disturbance contributes to increased energetic expenditure which can result in increased likelihood of winter mortality or reduced fitness (if energy expenditure is greater than energy gain), which can negatively impact population trends (see, for example, Madsen and Fox, 1995). Factors such as intensity, frequency, timing and duration of a (direct or indirect) disturbance source must be taken into account to determine the potential impact upon the targets for population trend and spatial distribution
Barriers to connectivity and site use	Number, location, shape and hectares	Barriers do not significantly impact the wintering population's access to the SPA or other ecologically important sites outside the SPA	Barriers limiting the population's access to this SPA or ecologically important sites outside the SPA will ultimately affect the achievement of targets for population trend and/or spatial distribution. Factors such as the number, location, shape and area of potential barriers must be taken into account to determine their potential impact
Forage spatial distribution, extent and abundance	Location, hectares, and forage biomass	Sufficient number of locations, area of suitable habitat and available forage biomass to support the population target	This species is omnivorous and forages primarily in open freshwater or brackish waterbodies. The species prefers shallow areas where it feeds by diving (to depths of 1-2.5m, typically) but also dabbles at the surface. Diet is predominantly aquatic plants (taken when diving or at the surface) but also preys upon molluscs, fish, and insects (such as chironomid larvae). Utilised habitats include lakes, rivers and flood-waters, reservoirs, estuaries, and (less so) coastal areas. In winter, forages alone or as part of large aggregations

Roost spatial distribution and extent	Location and hectares of roosting habitat	Sufficient number of locations, area and availability of suitable roosting habitat to support the population target	Roosting is a critical ecological requirement for the wintering population. When roosting overnight, Pochard utilise open waterbodies (see foraging habitats). Daytime roosting is also a common behaviour, where birds minimise activity levels to conserve energy, while benefitting from the vigilance of other flock members. A lack of sufficient and suitable roosting habitats can result in increased mortality risk, whether indirectly (e.g. via increased energy expenditure travelling to/from roost sites) or directly (e.g. via increased predation risk), or reduction in site use; this would ultimately affect the achievement of targets for population trend and/or spatial distribution
---------------------------------------	---	---	---

Conservation Objectives for : Lough Derravaragh SPA [004043]

A061 Tufted Duck *Aythya fuligula*

To restore the Favourable conservation condition of Tufted Duck in Lough Derravaragh SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Winter population trend	Percentage change in number of individuals	Long term winter population trend is stable or increasing	The national population of wintering Tufted Duck in Ireland has decreased by 18% from 1994/95 - 2019/20, as monitored via I-WeBS (Kennedy et al., 2022). During the baseline assessments to inform SPA designation, 1,073 Tufted Duck were estimated to be using this SPA (5 year mean of peak counts for baseline period 1995/96 - 1999/00; see NPWS, 2013). A population of 167 Tufted Duck was estimated to be using the Lough Derravaragh SPA in recent years (5 year mean of peak counts from I-WeBS monitoring for the period 2017/18 - 2021/22). This represents an estimated population decrease of 85% since the baseline period which is significantly greater than the national trend. Population declines are likely linked, at least in part, to distribution shifts driven by climate change (see Pavón-Jordan et al., 2018)
Winter spatial distribution	Hectares, time and intensity of use	Sufficient number of locations, area, and availability (in terms of timing and intensity of use) of suitable habitat to support the population target	Distribution encapsulates the number of locations and area of potentially suitable habitat for the wintering population and its availability for use. The suitability and availability of habitat areas is likely to vary throughout the season, for example, due to variation in land management practices or the abundance of resources available (due to natural variation and other factors). This will affect the spatio-temporal patterns of use of the habitats by the wintering population
Disturbance at wintering site	Intensity, frequency, timing and duration	The intensity, frequency, timing and duration of disturbance occurs at levels that do not significantly impact the achievement of targets for population trend and spatial distribution	The impact of any significant disturbance (direct or indirect) to the wintering population will ultimately affect the achievement of targets for population trend and/or spatial distribution. Disturbance contributes to increased energetic expenditure which can result in increased likelihood of winter mortality or reduced fitness (if energy expenditure is greater than energy gain), which can negatively impact population trends (see, for example, Madsen and Fox, 1995). Factors such as intensity, frequency, timing and duration of a (direct or indirect) disturbance source must be taken into account to determine the potential impact upon the targets for population trend and spatial distribution
Barriers to connectivity and site use	Number, location, shape and hectares	Barriers do not significantly impact the wintering population's access to the SPA or other ecologically important sites outside the SPA	Barriers limiting the population's access to this SPA or ecologically important sites outside the SPA will ultimately affect the achievement of targets for population trend and/or spatial distribution. Factors such as the number, location, shape and area of potential barriers must be taken into account to determine their potential impact
Forage spatial distribution, extent and abundance	Location, hectares, and forage biomass	Sufficient number of locations, area of suitable habitat and available forage biomass to support the population target	This species is omnivorous and forages primarily in open freshwater or brackish waterbodies. Molluscs are the main food source, and hence the species prefers shallow areas (to c.15m depth), but will also consume fish, insects, amphibians and various plant materials (leaves, shoots, tubers, seeds). Tufted Duck feed primarily by diving, but to a lesser extent will also feed at the surface of waterbodies, wade in shallows, and forage onshore (e.g. for cereal grain). Utilised habitats include lakes, rivers, ponds, reservoirs, marshes, estuaries, lagoons, and (less so) coastal areas. In winter, individuals can forage alone or as part of large aggregations

Roost spatial distribution and extent	Location and hectares of roosting habitat	Sufficient number of locations, area and availability of suitable roosting habitat to support the population target	When roosting overnight, the species uses a range of waterbodies, as noted for foraging habitat. Roosting is a critical ecological requirement for the wintering population. Daytime roosting is also a common behaviour, where birds minimise activity levels to conserve energy, while benefitting from the vigilance of other flock members. A lack of sufficient and suitable roosting habitats can result in increased mortality risk, whether indirectly (e.g. via increased energy expenditure travelling to/from roost sites) or directly (e.g. via increased predation risk), or reduction in site use; this would ultimately affect the achievement of targets for population trend and/or spatial distribution
---------------------------------------	---	---	---

Conservation Objectives for : Lough Derravaragh SPA [004043]

A125

Coot *Fulica atra*

To restore the Favourable conservation condition of Coot in Lough Derravaragh SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Winter population trend	Percentage change in number of individuals	Long term winter population trend is stable or increasing	The national population of wintering Coot in Ireland has decreased by 23% from 1994/95 - 2019/20, as monitored via I-WeBS (Kennedy et al., 2022). During the baseline assessments to inform SPA designation, 1,358 Coot were estimated to be using this SPA (5 year mean of peak counts for baseline period 1995/96 - 1999/00; see NPWS, 2013). A population of 702 Coot was estimated to be using the Lough Derravaragh SPA in recent years (5 year mean of peak counts from I-WeBS monitoring for the period 2017/18 - 2021/22). This represents an estimated population decrease of 49% since the baseline period which is greater than the national trend. Population declines are likely linked, at least in part, to distribution shifts driven by climate change (see Pavón-Jordan et al., 2018)
Winter spatial distribution	Hectares, time and intensity of use	Sufficient number of locations, area, and availability (in terms of timing and intensity of use) of suitable habitat to support the population target	Distribution encapsulates the number of locations and area of potentially suitable habitat for the wintering population and its availability for use. The suitability and availability of habitat areas is likely to vary throughout the season, for example, due to variation in land management practices or the abundance of resources available (due to natural variation and other factors). This will affect the spatio-temporal patterns of use of the habitats by the wintering population
Disturbance at wintering site	Intensity, frequency, timing and duration	The intensity, frequency, timing and duration of disturbance occurs at levels that do not significantly impact the achievement of targets for population trend and spatial distribution	The impact of any significant disturbance (direct or indirect) to the wintering population will ultimately affect the achievement of targets for population trend and/or spatial distribution. Disturbance contributes to increased energetic expenditure which can result in increased likelihood of winter mortality or reduced fitness (if energy expenditure is greater than energy gain), which can negatively impact population trends (see, for example, Madsen and Fox, 1995). Factors such as intensity, frequency, timing and duration of a (direct or indirect) disturbance source must be taken into account to determine the potential impact upon the targets for population trend and spatial distribution
Barriers to connectivity and site use	Number, location, shape and hectares	Barriers do not significantly impact the wintering population's access to the SPA or other ecologically important sites outside the SPA	Barriers limiting the population's access to this SPA or ecologically important sites outside the SPA will ultimately affect the achievement of targets for population trend and/or spatial distribution. Factors such as the number, location, shape and area of potential barriers must be taken into account to determine their potential impact
Forage spatial distribution, extent and abundance	Location, hectares, and forage biomass	Sufficient number of locations, area of suitable habitat and available forage biomass to support the population target	This species is omnivorous; plants dominant the diet but it will also take invertebrate and vertebrate prey. It forages primarily in waterbodies, rarely foraging far from them. The species feeds at the surface and sub-surface of waterbodies by up-ending and diving. It prefers shallow, open, slow moving waterbodies with marginal, floating, emergent or bottom vegetation. Foraging habitats utilised by Coot include rivers, canals, lakes, reservoirs, ponds, lagoons, estuaries, drainage channels and flooded lands. In winter, individuals can forage alone or as part of large aggregations

Roost spatial distribution and extent	Location and hectares of roosting habitat	Sufficient number of locations, area and availability of suitable roosting habitat to support the population target	When roosting overnight, Coot use a range of waterbodies, as noted for foraging habitat. Roosting is a critical ecological requirement for the wintering population. Daytime roosting is also a common behaviour, where birds minimise activity levels to conserve energy, while benefitting from the vigilance of other flock members. A lack of sufficient and suitable roosting habitats can result in increased mortality risk, whether indirectly (e.g. via increased energy expenditure travelling to/from roost sites) or directly (e.g. via increased predation risk), or reduction in site use; this would ultimately affect the achievement of targets for population trend and/or spatial distribution
---------------------------------------	---	---	---


Conservation Objectives for : Lough Derravaragh SPA [004043]


A999

Wetlands

To maintain the Favourable conservation condition of Wetland habitats in Lough Derravaragh SPA as a resource for the regularly-occurring migratory waterbirds that utilise these areas. This is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Wetland habitat area	Hectares	No significant loss to wetland habitat within the SPA, other than that occurring from natural patterns of variation	Any significant loss to the wetland habitat within the SPA would likely negatively impact the regularly-occurring migratory waterbirds that utilise this wetland habitat. Such loss of wetland habitat would likely reduce the diversity and abundance of waterbird species that the wetland can support. This, in turn, could negatively impact the Conservation Objectives for waterbird species listed as Special Conservation Interests in the SPA or other regularly-occurring migratory waterbird species
Wetland habitat quality and functioning	Quality and function of the wetland habitat	No significant impact on the quality or functioning of the wetland habitat within the SPA, other than that occurring from natural patterns of variation	Any significant impact on the quality, functioning and accessibility of the wetland habitat within the SPA would likely negatively impact the regularly-occurring migratory waterbirds that utilise this wetland habitat. Impacts on wetland quality, functioning and accessibility would likely reduce the diversity and abundance of waterbird species that the wetland can support. This, in turn, could negatively impact the Conservation Objectives for waterbird species listed as Special Conservation Interests in the SPA or other regularly-occurring migratory waterbird species





Legend

Lough Derravaragh SPA 004043

Garriskil Bog SAC 000679