ISSN 2009-4086

National Parks and Wildlife Service

Conservation Objectives Series

Carrowmore Lake SPA 004052



National Parks and Wildlife Service, Department of Housing, Local Government and Heritage,

90 King Street North, Dublin 7, D07 N7CV, Ireland.

Web: www.npws.ie E-mail: natureconservation@npws.gov.ie

Citation:

NPWS (2025) Conservation Objectives: Carrowmore Lake SPA 004052. Version 1. National Parks and Wildlife Service, Department of Housing, Local Government and Heritage.

> Series Editors: Maria Long and Colin Heaslip ISSN 2009-4086

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance
- exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

• population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and

• the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and

• there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.

2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.

3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.

4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.

5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

* indicates a priority habitat under the Habitats Directive

004052	Carrowmore Lake SPA

A191 Sandwich Tern Sterna sandvicensis

Please note that this SPA overlaps with Carrowmore Lake Complex SAC (000476). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping site as appropriate.

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	2021				
Title :	Estimated foraging ranges of the breeding seabirds of Ireland's marine special protected area network				
Author :	Power, A.; McDonnell, P.; Tierney, T.D.				
Series :	Published NPWS report				
Year :	2022				
Title :	Lady's Island Lake Tern Report 2022				
Author :	Stubbings, E.; Büche, B.; Murray, T.; Newton, S.				
Series :	BirdWatch Ireland Seabird Conservation Report to NPWS				
Year :	2023				
Year : Title :	2023 Lady's Island Lake Tern Report 2023				
Year : Title : Author :	2023 Lady's Island Lake Tern Report 2023 Stubbings, E.; Büche, B.; Murray, T.; Newton, S.				
Year : Title : Author : Series :	2023 Lady's Island Lake Tern Report 2023 Stubbings, E.; Büche, B.; Murray, T.; Newton, S. BirdWatch Ireland Seabird Conservation Report to NPWS				
Year : Title : Author : Series : Year :	2023 Lady's Island Lake Tern Report 2023 Stubbings, E.; Büche, B.; Murray, T.; Newton, S. BirdWatch Ireland Seabird Conservation Report to NPWS 2024				
Year : Title : Author : Series : Year : Title :	2023 Lady's Island Lake Tern Report 2023 Stubbings, E.; Büche, B.; Murray, T.; Newton, S. BirdWatch Ireland Seabird Conservation Report to NPWS 2024 Lady's Island Lake Tern Report 2024				
Year : Title : Author : Series : Year : Title : Author :	2023 Lady's Island Lake Tern Report 2023 Stubbings, E.; Büche, B.; Murray, T.; Newton, S. BirdWatch Ireland Seabird Conservation Report to NPWS 2024 Lady's Island Lake Tern Report 2024 Stubbings, E.; Büche, B.; Doyle, H.; Burke, B.; Newton, S.				

Other References

Year :	1985		
Title :	The 1984 all Ireland tern survey		
Author :	Whilde, A.		
Series :	Irish Birds 3: 1-32		
Year :	1997		
Title :	The status and distribution of breeding sandwich, roseate, common, arctic and little terns in Ireland in 1995		
Author :	Hannon, C.; Berrow, S.D.; Newton, S.F.		
Series :	Irish Birds, 6: 1-22		
Year :	2003		
Title :	Implications for seaward extensions to existing breeding seabird colony Special Protection Areas		
Author :	McSorley, C.A.; Dean, B.J.; Webb, A.; Reid J.B.		
Series :	JNCC Report No. 329		
Year :	2004		
Title :	Seabird populations of Britain and Ireland		
Author :	Mitchell, P.I.; Newton, S.F.; Ratcliffe, N.; Dunn, T.E.		
Series :	Poyser, London		
Year :	2010		
Title :	How Representative is the Current Monitoring of Breeding Seabirds in the UK?		
Author :	Cook, A.S.C.P.; Robinson, R.A.		
Series :	BTO Research Report No. 573		
Year :	2019		
Title :	Desk-based revision of seabird foraging ranges used for HRA screening		
Author :	Woodward, I.; Thaxter, C.B.; Owen, E.; Cook, A.S.C.P.		
Series :	BTO Research Report No. 724		
Year :	2020		
Title :	Sandwich Tern (<i>Thalasseus sandvicensis</i>), version 1.0. In Birds of the World (S. M. Billerman, Editor)		
Author :	Shealer, D.; Liechty, J. S.; Pierce, A. R.; Pyle, P.; Patten., M. A.		
Series :	Cornell Lab of Ornithology, Ithaca, NY, USA		
Year :	2023		
Title :	Seabirds Count: a census of breeding seabirds in Britain and Ireland (2015-2021)		
Author :	Burnell, D.; Perkins, A.J.; Newton, S.F.; Bolton, M.; Tierney, T.D.; Dunn, T.E.		
Series :	Lynx Nature Books, Barcelona		

Conservation Objectives for : Carrowmore Lake SPA [004052]

A191 Sandwich Tern *Sterna sandvicensis*

To restore the Favourable conservation condition of Sandwich Tern in Carrowmore Lake SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Breeding population size	Number of Apparently Occupied Nests (AON)	Long term SPA population trend is stable or increasing	Sandwich Tern have bred at this SPA, on Derreens Island, since at least 1984 when the population was 164 pairs (Whilde, 1985). However, there were no breeding pairs in 1995 (Hannon et al., 1997). It was suspected that the colony had moved to Inishderry Island in Broadhaven Bay outside of the SPA (NPWS internal files). Three individuals were recorded in 2000, which indicates 1 - 2 pairs may have bred (Mitchell et al., 2004). Sandwich Tern did not breed in this SPA in 2017 and there are no records in subsequent years (Burnell et al., 2023). Some populations can show strong site fidelity but generally this species exhibits a distinct lack of fidelity to particular colony sites (Shealer et al., 2020) so if the breeding habitat is suitable it is possible that breeding Sandwich Tern may return. In contrast, the national population has increased by 39% between 1998 - 2002 and 2015 - 2021 (Burnell et al., 2023), primarily as a result of increases at wardened colonies
Productivity rate	Number of fledged young per breeding pair	Sufficient to maintain a stable or increasing population	There was no productivity data available for this species in this SPA. The breeding population at Lady's Island Lake SPA is one of the most well monitored in the country. During the period 2022 - 2024 the estimated productivity ranged from 0.13 to 0.72 chicks presumed fledged per AON with a mean of 0.49 (Stubbings et al., 2022, 2023 and 2024). Note the particularly low estimate of 0.13 in 2023 was caused by excessive predation at the study site and may not be representative of the SPA as a whole (Stubbings et al., 2023). Cook and Robinson (2010) undertook Population Viability Analyses (PVA) of a selection of breeding populations in the UK. Over their study period Sandwich Tern productivity at monitored nests was 0.66. Were this level to be maintained, Sandwich Tern populations would decline by 62% over 25 years. For the population to stabilise, breeding success would have to increase to 1.1 chicks per nest per year
Distribution: extent of available nesting options within the SPA	Numbers and spatial distribution	Sufficient availability of suitable nesting sites throughout the SPA to maintain a stable or increasing population	Distribution encapsulates the number of locations and area of potentially suitable nesting habitat for the breeding population and its availability for use. The suitability and availability of habitat across the SPA may vary through time. This will affect the spatio-temporal patterns of use of the habitats by Sandwich Tern. Typically, colonies are located in areas with bare sand, small pebbles, or short grass (Shealer et al., 2020). In Ireland, all known large colonies are situated on marine or inland islands of varying distances from the mainland/shore. Sandwich Tern have historically bred on Derreens Island, the largest island within the SPA, which is dominated by a grassy sward
Forage spatial distribution, extent, abundance and availability	Location, hectares, and forage biomass	Sufficient number of locations, area of suitable habitat and available forage biomass to support the population target	Sandwich Tern feed primarily along coastal marine areas. They are largely piscivorous. In north- temperate regions of Europe they primarily eat Clupeidae (herrings) and Ammodytidae (sandeels) families (Shealer et al., 2020). Based on several studies, Woodward et al. (2019) provide estimates (i.e. overall mean, mean of maximum distances across all studies, and maximum distance recorded) of Sandwich Tern foraging ranges from the nest site during the breeding season, which are 9km, 34km, and 80km respectively (see Power et al., 2021)

Disturbance at the breeding site	Intensity, frequency, timing and duration	Disturbance occurs at levels that do not significantly impact on birds at the breeding site	Disturbance events at the nest site/breeding colony level can result in a reduction of overall productivity and even lead to the abandonment of the breeding colony. The impact of any significant disturbance (direct or indirect) to the breeding population will ultimately affect the achievement of targets for population size and/or spatial distribution. Disturbance contributes to increased energetic expenditure, which can result in increased likelihood of mortality or reduced fitness (if energy expenditure is greater than energy gain) and, in turn, negatively impact population trends. Factors such as intensity, frequency, timing, and duration of a (direct or indirect) disturbance source must be taken into account to determine the potential impact upon the targets for population size and spatial distribution
Disturbance at areas ecologically connected to the colony	Intensity, frequency, timing and duration	Disturbance occurs at levels that do not significantly impact on breeding population	Seabird species can make extensive use of the waters adjacent to their breeding colonies for non site-specific maintenance behaviours (e.g. courtship, bathing, preening) as defined in McSorley et al. (2003). Additionally, some species may engage in maintenance behaviours outside of the breeding colony but not in the water. For example, terns may roost on rocky islets or beaches away from the breeding colony
Barriers to connectivity	Number, location, shape, and area (ha)	Barriers do not significantly impact the population's access to the SPA or other ecologically important sites outside the SPA	Seabirds, particularly during the breeding season, require regular access to waters ecologically connected to the colony in order to forage as well as to engage in other maintenance behaviours. Based on several studies, Woodward et al. (2019) provide estimates (i.e. overall mean, mean of maximum distances across all studies, and maximum distance recorded) of Sandwich Tern foraging ranges from the nest site during the breeding season, which are 9km, 34km, and 80km respectively (see Power et al., 2021)



