

National Parks and Wildlife Service

Conservation Objectives Series

The Gearagh SPA 004109



NPWS

An tSeirbhís Páirceanna
Náisiúnta agus Fiadhúlra
National Parks and Wildlife
Service

**National Parks and Wildlife Service,
Department of Housing, Local Government and Heritage,
90 King Street North, Dublin 7, D07 N7CV, Ireland.
Web: www.npws.ie
E-mail: natureconservation@npws.gov.ie**

Citation:

**NPWS (2025) Conservation Objectives: The Gearagh SPA 004109. Version 1.
National Parks and Wildlife Service, Department of Housing, Local Government
and Heritage.**

**Series Editors: Maria Long and Colin Heaslip
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

004109	The Gearagh SPA
<hr/>	
A050	Wigeon <i>Anas penelope</i>
A052	Teal <i>Anas crecca</i>
A053	Mallard <i>Anas platyrhynchos</i>
A125	Coot <i>Fulica atra</i>
A999	Wetlands

**Please note that this SPA overlaps with The Gearagh SAC (000108).
See map 2. The conservation objectives for this site should be used in
conjunction with those for the overlapping site as appropriate.**

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

Other References

Year :	1995
Title :	Impacts of hunting disturbance on waterbirds - a review
Author :	Madsen, J.; Fox, A.D.
Series :	Wildlife Biology 1(4):193-207
Year :	2019
Title :	Report under Article 12 of the Birds Directive Period 2013-2018
Author :	EEA
Series :	European Environment Agency. European Topic Centre on Biological Diversity. Pp 1-9. https://cdr.eionet.europa.eu/Converters/run_conversion?file=ie/eu/art12/envxztxxq/IE_birds_reports_20191031-130157.xml&conv=612&source=remote
Year :	2020
Title :	Green-winged Teal (<i>Anas crecca</i>), version 1.0. In Birds of the World (S. M. Billerman, Editor)
Author :	Johnson, K.; Carboneras, C.; Christie, D. A.; Kirwan, G. M.
Series :	Cornell Lab of Ornithology, Ithaca, NY, USA
Year :	2022
Title :	Irish wetland bird survey: I-WeBS national and site trends report 1994/95 – 2019/20
Author :	Kennedy, J.; Burke, B.; Fitzgerald, N.; Kelly, S.B.A.; Walsh, A.J.; Lewis, L.J.
Series :	https://birdwatchireland.ie/app/uploads/2022/04/iwebs_trends_report.html

Conservation Objectives for : The Gearagh SPA [004109]

A050 Wigeon *Anas penelope*

To restore the Favourable conservation condition of Wigeon in The Gearagh SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Winter population trend	Percentage change in number of individuals	Long term winter population trend is stable or increasing	The national population of wintering Wigeon in Ireland has declined by 18% from 1994/95 - 2019/20, as monitored via the Irish Wetland Bird Survey (I-WeBS) (Kennedy et al., 2022). The Gearagh SPA comprises a proportion of the I-WeBS monitoring site known as the Inishcarra Reservoirs. In a robust statistical analysis of I-WeBS data, Kennedy et al. (2022) determined that the population of Wigeon that used the Inishcarra Reservoirs decreased by 60% between winters 1995/96 - 2019/20. This trend is considered representative of the SPA given the ecological connectivity between the SPA and the wider Inishcarra Reservoirs site. This trend is significantly greater than the reported national population trend. In their analyses, Kennedy et al. (2022) used indices of the total count of a given species across the entire winter period, providing a more comprehensive assessment of change in the population that used the site compared to approaches such as a comparison of 5 year mean peak values
Winter spatial distribution	Hectares, time and intensity of use	Sufficient number of locations, area, and availability (in terms of timing and intensity of use) of suitable habitat to support the population target	Distribution encapsulates the number of locations and area of potentially suitable habitat for the wintering population and its availability for use. The suitability and availability of habitat areas are likely to vary throughout the season, for example, due to variation in land management practices or the abundance of resources available (due to natural variation and other factors). This will affect the spatio-temporal patterns of use of the habitats by the wintering population
Disturbance at wintering site	Intensity, frequency, timing and duration	Disturbance occurs at levels that do not significantly impact the achievement of targets for population trend and distribution	The impact of any significant disturbance (direct or indirect) to the wintering population will ultimately affect the achievement of targets for population trend and/or spatial distribution. Disturbance contributes to increased energetic expenditure which can result in increased likelihood of winter mortality or reduced fitness (if energy expenditure is greater than energy gain) and, in turn, negatively impact population trends (see, for example, Madsen and Fox, 1995). Factors such as intensity, frequency, timing and duration of a (direct or indirect) disturbance source must be taken into account to determine the potential impact upon the targets for population trend and spatial distribution
Barriers to connectivity and site use	Number, location, shape and hectares	Barriers do not significantly impact the wintering population's access to the SPA or other ecologically important sites outside the SPA	Barriers limiting the population's access to this SPA or ecologically important sites outside the SPA will ultimately affect the achievement of targets for population trend and/or spatial distribution. Factors such as the number, location, shape and area of potential barriers must be taken into account to determine their potential impact. Access to ecologically important sites outside the SPA must also be considered as a single SPA may not satisfy all the ecological requirements of the wintering population, and it may require access to other SPAs or sites for certain activities, such as foraging when preferred foraging areas are unavailable due to disturbance, extensive flooding, or other factors

Forage spatial distribution, extent and abundance	Location, hectares, and forage biomass	Sufficient number of locations, area of suitable habitat and available forage biomass to support the population target	This dabbling duck feeds primarily on aquatic vegetation, at surface-level in waterbodies or at ground level in wetland habitats. Key forage materials include leaves, stems, stolons, roots, rhizomes, and seeds (including cereals). Key wintering habitats are marshes, lagoons, estuaries, coastal bays, lakes, rivers and river floodplains, turloughs and other wetland habitats, as well as pastures
Roost spatial distribution and extent	Location and hectares of roosting habitat	Sufficient number of locations, area and availability of suitable roosting habitat to support the population target	Wigeon rely primarily on wetlands or waterbodies for roosting. Roosting is a critical ecological requirement for the wintering population. When roosting overnight, this species typically utilises a similar range of habitats as noted for foraging. Daytime roosting is also a common behaviour, where birds minimise activity levels to conserve energy, while benefitting from the vigilance of other flock members. A lack of sufficient and suitable roosting habitats can result in increased mortality risk, whether indirectly (e.g. via increased energy expenditure travelling to/from roost sites) or directly (e.g. via increased predation risk), or reduction in site use; this would ultimately affect the achievement of targets for population trend and/or spatial distribution
Supporting habitat: area and quality	Hectares and quality	Sufficient area of utilisable habitat available in ecologically important sites outside the SPA	The wintering population can make extensive use of suitable habitats in important areas outside the SPA for foraging and roosting. The extent, availability and quality of these supporting habitats may be of importance for the resilience of the SPA population. Suitable supporting habitats include those highlighted in the attributes for foraging and roosting habitat

Conservation Objectives for : The Gearagh SPA [004109]

A052

Teal *Anas crecca*

To restore the Favourable conservation condition of Teal in The Gearagh SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Winter population trend	Percentage change in number of individuals	Long term winter population trend is stable or increasing	The national population of wintering Teal in Ireland has increased by 19% from 1994/95 - 2019/20, as monitored via the Irish Wetland Bird Survey (I-WeBS) (Kennedy et al., 2022). The Gearagh SPA comprises a proportion of the I-WeBS monitoring site known as the Inishcarra Reservoirs. In a robust statistical analysis of I-WeBS data, Kennedy et al. (2022) determined that the population of Teal that used the Inishcarra Reservoirs decreased by 17% between winters 1995/96 - 2019/20. This trend is considered representative of the SPA given the ecological connectivity between the SPA and the wider Inishcarra Reservoirs site. This result is similar to the reported national population trend. In their analyses, Kennedy et al. (2022) used indices of the total count of a given species across the entire winter period, providing a more comprehensive assessment of change in the population that used the site compared to approaches such as a comparison of 5 year mean peak values
Winter spatial distribution	Hectares, time and intensity of use	Sufficient number of locations, area, and availability (in terms of timing and intensity of use) of suitable habitat to support the population target	Distribution encapsulates the number of locations and area of potentially suitable habitat for the wintering population and its availability for use. The suitability and availability of habitat areas are likely to vary throughout the season, for example, due to variation in land management practices or the abundance of resources available (due to natural variation and other factors). This will affect the spatio-temporal patterns of use of the habitats by the wintering population
Disturbance at wintering site	Intensity, frequency, timing and duration	Disturbance occurs at levels that do not significantly impact the achievement of targets for population trend and distribution	The impact of any significant disturbance (direct or indirect) to the wintering population will ultimately affect the achievement of targets for population trend and/or spatial distribution. Disturbance contributes to increased energetic expenditure which can result in increased likelihood of winter mortality or reduced fitness (if energy expenditure is greater than energy gain) and, in turn, negatively impact population trends (see, for example, Madsen and Fox, 1995). Factors such as intensity, frequency, timing and duration of a (direct or indirect) disturbance source must be taken into account to determine the potential impact upon the targets for population trend and spatial distribution
Barriers to connectivity and site use	Number, location, shape and hectares	Barriers do not significantly impact the wintering population's access to the SPA or other ecologically important sites outside the SPA	Barriers limiting the population's access to this SPA or ecologically important sites outside the SPA will ultimately affect the achievement of targets for population trend and/or spatial distribution. Factors such as the number, location, shape and area of potential barriers must be taken into account to determine their potential impact. Access to ecologically important sites outside the SPA must also be considered as a single SPA may not satisfy all the ecological requirements of the wintering population, and it may require access to other SPAs or sites for certain activities, such as foraging when preferred foraging areas are unavailable due to disturbance, extensive flooding, or other factors

Forage spatial distribution, extent and abundance	Location, hectares, and forage biomass	Sufficient number of locations, area of suitable habitat and available forage biomass to support the population target	Teal utilise a wide range of foraging habitats and have a broad diet. Key food sources are: small seeds of sedges, grasses and aquatic vegetation; aquatic invertebrates, including larvae, such as molluscs and crustaceans; as well as algae (particularly <i>Enteromorpha</i> spp.) (Johnson et al., 2020). Key habitats include shallow water, between depths of 4cm (dabbling) - 24cm (upending), and can be widespread on wetlands with good cover, such as reedbeds. The species uses a wide variety of shallow areas within wetland habitats, both coastal and inland, including estuaries, lagoons, mudflats, marshes, floodplains, lakes, ponds, turloughs and agricultural areas. They feed by day where they are safe from disturbance
Roost spatial distribution and extent	Location and hectares of roosting habitat	Sufficient number of locations, area and availability of suitable roosting habitat to support the population target	Roosting is a critical ecological requirement for the wintering population. When roosting overnight, Teal primarily utilise permanent waterbodies, marshes, wide ditches, wet grassland and wetlands (see foraging habitats). Daytime roosting is also a common behaviour, where birds minimise activity levels to conserve energy, while benefitting from the vigilance of other flock members. A lack of sufficient and suitable roosting habitats can result in increased mortality risk, whether indirectly (e.g. via increased energy expenditure travelling to/from roost sites) or directly (e.g. via increased predation risk), or reduction in site use; this would ultimately affect the achievement of targets for population trend and/or spatial distribution
Supporting habitat: area and quality	Hectares and quality	Sufficient area of utilisable habitat available in ecologically important sites outside the SPA	The wintering population can make extensive use of suitable habitats in important areas outside the SPA for foraging and roosting. The extent, availability and quality of these supporting habitats may be of importance for the resilience of the SPA population. Suitable supporting habitats include those highlighted in the attributes for foraging and roosting habitat

Conservation Objectives for : The Gearagh SPA [004109]

A053 Mallard *Anas platyrhynchos*

To restore the Favourable conservation condition of Mallard in The Gearagh SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Winter population trend	Percentage change in number of individuals	Long term winter population trend is stable or increasing	The national population of wintering Mallard in Ireland has declined by 19% from 1994/95 - 2019/20, as monitored via the Irish Wetland Bird Survey (I-WeBS) (Kennedy et al., 2022). The Gearagh SPA comprises a proportion of the I-WeBS monitoring site known as the Inishcarra Reservoirs. In a robust statistical analysis of I-WeBS data, Kennedy et al. (2022) determined that the population of Mallard that used the Inishcarra Reservoirs decreased by 35% between winters 1995/96 - 2019/20. This trend is considered representative of the SPA given the ecological connectivity between the SPA and the wider Inishcarra Reservoirs site. This result is similar to the reported national population trend. In their analyses, Kennedy et al. (2022) used indices of the total count of a given species across the entire winter period, providing a more comprehensive assessment of change in the population that used the site compared to approaches such as a comparison of 5 year mean peak values
Winter spatial distribution	Hectares, time and intensity of use	Sufficient number of locations, area, and availability (in terms of timing and intensity of use) of suitable habitat to support the population target	Distribution encapsulates the number of locations and area of potentially suitable habitat for the wintering population and its availability for use. The suitability and availability of habitat areas are likely to vary throughout the season, for example, due to variation in land management practices or the abundance of resources available (due to natural variation and other factors). This will affect the spatio-temporal patterns of use of the habitats by the wintering population
Disturbance at wintering site	Intensity, frequency, timing and duration	Disturbance occurs at levels that do not significantly impact the achievement of targets for population trend and distribution	The impact of any significant disturbance (direct or indirect) to the wintering population will ultimately affect the achievement of targets for population trend and/or spatial distribution. Disturbance contributes to increased energetic expenditure which can result in increased likelihood of winter mortality or reduced fitness (if energy expenditure is greater than energy gain) and, in turn, negatively impact population trends (see, for example, Madsen and Fox, 1995). Factors such as intensity, frequency, timing and duration of a (direct or indirect) disturbance source must be taken into account to determine the potential impact upon the targets for population trend and spatial distribution
Barriers to connectivity and site use	Number, location, shape and hectares	Barriers do not significantly impact the wintering population's access to the SPA or other ecologically important sites outside the SPA	Barriers limiting the population's access to this SPA or ecologically important sites outside the SPA will ultimately affect the achievement of targets for population trend and/or spatial distribution. Factors such as the number, location, shape and area of potential barriers must be taken into account to determine their potential impact. Access to ecologically important sites outside the SPA must also be considered as a single SPA may not satisfy all the ecological requirements of the wintering population, and it may require access to other SPAs or sites for certain activities, such as foraging when preferred foraging areas are unavailable due to disturbance, extensive flooding, or other factors

Forage spatial distribution, extent and abundance	Location, hectares, and forage biomass	Sufficient number of locations, area of suitable habitat and available forage biomass to support the population target	Foraging habitats include a range of wetlands, such as marshes, flooded areas, lakes, estuaries and lagoons, as well as grasslands. In winter, Mallard are primarily herbivorous, dabbling for roots, leaves, stems and seeds of plants in surface waters. Mallard will also consume aquatic insects, crustaceans, and molluscs, and, where adjacent to wetlands, they occasionally graze on grasslands or consume cereal grain
Roost spatial distribution and extent	Location and hectares of roosting habitat	Sufficient number of locations, area and availability of suitable roosting habitat to support the population target	Roosting is a critical ecological requirement for the wintering population. When roosting overnight, Mallard primarily utilise permanent waterbodies and wetland habitat (see foraging habitats). Daytime roosting is also a common behaviour, where birds minimise activity levels to conserve energy, while benefitting from the vigilance of other flock members. A lack of sufficient and suitable roosting habitats can result in increased mortality risk, whether indirectly (e.g. via increased energy expenditure travelling to/from roost sites) or directly (e.g. via increased predation risk), or reduction in site use; this would ultimately affect the achievement of targets for population trend and/or spatial distribution
Supporting habitat: area and quality	Hectares and quality	Sufficient area of utilisable habitat available in ecologically important sites outside the SPA	The wintering population can make extensive use of suitable habitats in important areas outside the SPA for foraging and roosting. The extent, availability and quality of these supporting habitats may be of importance for the resilience of the SPA population. Suitable supporting habitats include those highlighted in the attributes for foraging and roosting habitat

Conservation Objectives for : The Gearagh SPA [004109]

A125

Coot *Fulica atra*

To restore the Favourable conservation condition of Coot in The Gearagh SPA, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Winter population trend	Percentage change in number of individuals	Long term winter population trend is stable or increasing	The national population of wintering Coot in Ireland has declined by 23% from 1994/95 - 2019/20, as monitored via the Irish Wetland Bird Survey (I-WeBS) (Kennedy et al., 2022). The Gearagh SPA comprises a proportion of the I-WeBS monitoring site known as the Inishcarra Reservoirs. In a robust statistical analysis of I-WeBS data, Kennedy et al. (2022) determined that the population of Coot that used the Inishcarra Reservoirs decreased by 80% between winters 1995/96 - 2019/20. This trend is considered representative of the SPA given the ecological connectivity between the SPA and the wider Inishcarra Reservoirs site. This result is significantly greater than the reported national population trend. In their analyses, Kennedy et al. (2022) used indices of the total count of a given species across the entire winter period, providing a more comprehensive assessment of change in the population that used the site compared to approaches such as a comparison of 5 year mean peak values
Winter spatial distribution	Hectares, time and intensity of use	Sufficient number of locations, area, and availability (in terms of timing and intensity of use) of suitable habitat to support the population target	Distribution encapsulates the number of locations and area of potentially suitable habitat for the wintering population and its availability for use. The suitability and availability of habitat areas are likely to vary throughout the season, for example, due to variation in land management practices or the abundance of resources available (due to natural variation and other factors). This will affect the spatio-temporal patterns of use of the habitats by the wintering population
Disturbance at wintering site	Intensity, frequency, timing and duration	Disturbance occurs at levels that do not significantly impact the achievement of targets for population trend and distribution	The impact of any significant disturbance (direct or indirect) to the wintering population will ultimately affect the achievement of targets for population trend and/or spatial distribution. Disturbance contributes to increased energetic expenditure which can result in increased likelihood of winter mortality or reduced fitness (if energy expenditure is greater than energy gain) and, in turn, negatively impact population trends (see, for example, Madsen and Fox, 1995). Factors such as intensity, frequency, timing and duration of a (direct or indirect) disturbance source must be taken into account to determine the potential impact upon the targets for population trend and spatial distribution
Barriers to connectivity and site use	Number, location, shape and hectares	Barriers do not significantly impact the wintering population's access to the SPA or other ecologically important sites outside the SPA	Barriers limiting the population's access to this SPA or ecologically important sites outside the SPA will ultimately affect the achievement of targets for population trend and/or spatial distribution. Factors such as the number, location, shape and area of potential barriers must be taken into account to determine their potential impact
Forage spatial distribution, extent and abundance	Location, hectares, and forage biomass	Sufficient number of locations, area of suitable habitat and available forage biomass to support the population target	This species is omnivorous; plants dominate the diet but it will also take invertebrate and vertebrate prey. It forages primarily in waterbodies, rarely foraging far from them. The species feeds at the surface and sub-surface of waterbodies by upending and diving. Coot prefer shallow, open, slow moving waterbodies with marginal, floating, emergent or bottom vegetation. Foraging habitats utilised by Coot include rivers, canals, lakes, reservoirs, ponds, lagoons, estuaries, drainage channels and flooded lands. In winter, individual Coot can forage alone or as part of large aggregations

Roost spatial distribution and extent	Location and hectares of roosting habitat	Sufficient number of locations, area and availability of suitable roosting habitat to support the population target	When roosting overnight, Coot use a range of waterbodies, as noted for foraging habitat. Roosting is a critical ecological requirement for the wintering population. Daytime roosting is also a common behaviour, where birds minimise activity levels to conserve energy, while benefitting from the vigilance of other flock members. A lack of sufficient and suitable roosting habitats can result in increased mortality risk, whether indirectly (e.g. via increased energy expenditure travelling to/from roost sites) or directly (e.g. via increased predation risk), or reduction in site use; this would ultimately affect the achievement of targets for population trend and/or spatial distribution
---------------------------------------	---	---	---



Conservation Objectives for : The Gearagh SPA [004109]

A999

Wetlands

To maintain the Favourable conservation condition of Wetland habitats in The Gearagh SPA as a resource for the regularly-occurring migratory waterbirds that utilise these areas. This is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Wetland habitat area	Hectares	No significant loss to wetland habitat within the SPA, other than that occurring from natural patterns of variation	Any significant loss to the wetland habitat within the SPA would likely negatively impact the regularly-occurring migratory waterbirds that utilise this wetland habitat. Such loss of wetland habitat would likely reduce the diversity and abundance of waterbird species that the wetland can support. This, in turn, could negatively impact the Conservation Objectives for waterbird species listed as Special Conservation Interests in the SPA or other regularly-occurring migratory waterbird species
Wetland habitat quality and functioning	Quality and function of the wetland habitat	No significant impact on the quality or functioning of the wetland habitat within the SPA, other than that occurring from natural patterns of variation	Any significant impact on the quality, functioning and accessibility of the wetland habitat within the SPA would likely negatively impact the regularly-occurring migratory waterbirds that utilise this wetland habitat. Impacts on wetland quality, functioning and accessibility would likely reduce the diversity and abundance of waterbird species that the wetland can support. This, in turn, could negatively impact the Conservation Objectives for waterbird species listed as Special Conservation Interests in the SPA or other regularly-occurring migratory waterbird species



Legend

The Gearagh SPA 004109

