



Site Name: River Shannon Callows SAC

Site Code: 000216

The River Shannon Callows is a long and diverse site which consists of seasonally flooded, semi-natural, lowland wet grassland, along and beside the river between the towns of Athlone and Portumna. It is approximately 50 km long and averages about 0.75 km wide (reaching 1.5 km wide in places). Along much of its length the site is bordered by raised bogs (many, but not all, of which are subject to large-scale harvesting), esker ridges and limestone-bedrock hills. The soils grade from silty-alluvial to peat. This site has a common boundary, and is closely associated, with two other sites with similar habitats, River Suck Callows and Little Brosna Callows.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

- | |
|-------------------------------------|
| [6410] <i>Molinia</i> Meadows |
| [6510] Lowland Hay Meadows |
| [7230] Alkaline Fens |
| [8240] Limestone Pavement* |
| [91E0] Alluvial Forests* |
| [1355] Otter (<i>Lutra lutra</i>) |

The River Shannon Callows is mainly composed of lowland wet grassland. Different plant communities occur, depending on elevation, and therefore flooding patterns. Two habitats listed on Annex I of the E.U. Habitats Directive are well-represented within the site – *Molinia* meadows and lowland hay meadows. The former is characterised by the presence of the Meadow Thistle (*Cirsium dissectum*) and Purple Moor-grass (*Molinia caerulea*), while typical species in the latter include Meadow Fescue (*Festuca pratensis*), Rough Meadow-grass (*Poa trivialis*), Downy Oat-grass (*Avenula pubescens*), Common Knapweed (*Centaurea nigra*), Ribwort Plantain (*Plantago lanceolata*) and Common Sorrel (*Rumex acetosa*). In places these two habitats grade into one another.

Low-lying areas of the callows with more prolonged flooding are characterised by Floating Sweet-grass (*Glyceria fluitans*), Marsh Foxtail (*Alopecurus geniculatus*) and wetland herbs such as Yellow-cress (*Rorippa* spp.), Water Forget-me-not (*Myosotis scorpioides*) and Common Spike-rush (*Eleocharis palustris*). Most of the callows consist of a plant community characterised by Creeping Bent (*Agrostis stolonifera*), Brown Sedge (*Carex disticha*), Common Sedge (*Carex nigra*), and herbs such as Marsh-marigold (*Caltha palustris*) and Marsh Bedstraw (*Galium palustre*), while the more

elevated and peaty areas are characterised by low-growing sedges, particularly Yellow Sedge (*Carex flava* agg.) and Star Sedge (*Carex echinata*). All these communities are very diverse in their total number of plant species, and include the scarce species Meadow-rue (*Thalictrum flavum*), Summer Snowflake (*Leucojum aestivum*) and Marsh Stitchwort (*Stellaria palustris*).

A further two Annex I habitats, both listed with priority status, have a minor though important presence within the site. Alluvial forest occurs on a series of alluvial islands just below the ESB weir near Meelick. Several of the islands are dominated by well-grown woodland consisting mainly of Ash (*Fraxinus excelsior*) and Willows (*Salix* spp.). The islands are prone to regular flooding from the river.

At Clorhane, an area of limestone pavement represents the only known example in Co. Offaly. It is predominantly colonised by mature Hazel (*Corylus avellana*) woodland, with areas of open limestone and calcareous grassland interspersed. The open limestone pavement comprises bare or moss-covered rock, or rock with a very thin calcareous soil cover supporting a short grassy turf. The most notable plant in the grassy area is a substantial population of Green-winged Orchid (*Orchis morio*), which occurs with such species as Sweet Vernal-grass (*Anthoxanthum odoratum*), Quaking-grass (*Briza media*), sedges (*Carex caryophyllea*, *C. flacca*), Common Bird's-foot-trefoil (*Lotus corniculatus*), Common Knapweed (*Centaurea nigra*), and Ribwort Plantain (*Plantago lanceolata*). Ferns associated with the cracks in the pavement include *Asplenium trichomanes*, *A. ruta-muraria*, *A. adiantum-nigrum* and *Polypodium australe*. Bryophytes include *Grimmia apocarpa* and *Orthotrichum* cf. *anomalum*. Anthills are common within the open grassland. The Hazel wood is well-developed and has herbaceous species such as Primrose (*Primula vulgaris*), Common Dog-violet (*Viola riviniana*), Wood-sorrel (*Oxalis acetosella*) and Herb-Robert (*Geranium robertianum*). The wood is noted for its luxuriant growth of epiphytic mosses and liverworts, with such species as *Neckera crispa* and *Hylocomium brevirostre*. Yew (*Taxus baccata*) occurs in one area.

Other habitats of smaller area but also of importance within the site are lowland dry grassland, drainage ditches, freshwater marshes and reedbeds. The dry grassland areas, especially where they exist within hay meadows, are species-rich, and of two main types: calcareous grassland on glacial material, and dry grassland on levees of river alluvium. The former can contain many orchid species, Cowslip (*Primula veris*), abundant Adder's-tongue (*Ophioglossum vulgatum*) and Spring-sedge (*Carex caryophyllea*), and both contain an unusually wide variety of grasses, including False Oat-grass (*Arrhenatherum elatius*), Yellow Oat-grass (*Trisetum flavescens*), Meadow Foxtail (*Alopecurus pratense*), and Meadow Brome (*Bromus commutatus*). In places Summer Snowflake also occurs.

Good quality habitats on the edge of the callows included in the site are wet broadleaved semi-natural woodland dominated by both Downy Birch (*Betula pubescens*) and Alder (*Alnus glutinosa*), and dry broadleaved woodland dominated by Hazel. There are also areas of raised bog, fen on old cut-away bog with Black Bog-rush (*Schoenus nigricans*), and a 'petrifying stream' with associated species-rich

calcareous flush which supports Yellow Sedge (*Carex lepidocarpa*), Blunt-flowered Rush (*Juncus subnodulosus*) and Stoneworts (*Chara* spp.).

Immediately south of Portumna Bridge and south east of the town of Portumna the area of low-lying terrestrial land west of the river comprises a large area of the Annex I habitat alkaline fen. The fen comprises a complex of rich-fen plant communities. Sedges (*Carex lasiocarpa*, *Carex acutiformis*) and Bogbean (*Menyanthes trifoliata*) dominate parts of the fens while other small sedges are common throughout. The orchids Early Marsh Orchid (*Dactylorhiza incarnata*), Western Marsh Orchid (*D. majalis*) and Marsh Helleborine (*Epipactis palustris*) and the red-listed plant species Marsh Pea (*Lathyrus palustris*) have been recorded within the fen.

Two species which are legally protected under the Flora (Protection) Order, 2015, occur in the site - Opposite-leaved Pondweed (*Groenlandia densa*) in drainage ditches, and Meadow Barley (*Hordeum secalinum*) on dry alluvial grassland. This is one of only two known inland sites for Meadow Barley in Ireland. The Red Data Book plant Green-winged Orchid is known from dry calcareous grasslands within the site.

The site is of international importance for wintering waterfowl as numbers regularly exceed the 20,000 threshold (mean of 34,985 for five winters 1994/94-1998/99). Of particular note is an internationally important population of Whooper Swans (287). A further five species have populations of national importance (all figures are means for five winters 1995/96-1999/00): Mute Swan (349), Wigeon (2972), Golden Plover (4254), Lapwing (11578) and Black-tailed Godwit (388). Species which occur in numbers of regional or local importance include Bewick's Swan, Tufted Duck, Dunlin, Curlew and Redshank. The population of Dunlin is notable as it is one of the few regular inland flocks in Ireland. Small flocks of Greenland White-fronted Goose use the Shannon Callows; these are generally associated with larger flocks which occur on the adjacent Little Brosna Callows and River Suck Callows.

Shoveler (an estimated 12 pairs in 1987) and Black-tailed Godwit (Icelandic race) (one or two pairs in 1987) breed within this site. These species are listed in the Red Data Book as being threatened in Ireland. The scarce bird Quail is also known to breed within the area. The callows has at times held over 40% of the Irish population of the globally endangered Corncrake, although numbers have declined in recent years. A total of 66 calling birds were recorded in 1999, but numbers have dropped significantly since then. The total population of breeding waders (Lapwing, Redshank, Snipe and Curlew) in 1987 was one of three major concentrations in Ireland and Britain. The population of breeding Redshank in the site was estimated to be 10% of the Irish population, making it nationally significant. Also, the Annex I species Merlin and Hen Harrier are regularly reported hunting over the callows during the breeding season and in autumn and winter.

This site holds a population of Otter, a species listed on Annex II of the E.U. Habitats Directive, while the Irish Hare, which is listed in the Irish Red Data Book, is a common sight on the callows.

The Shannon Callows are used for summer dry-stock grazing (mostly cattle, with some sheep and a few horses), and permanent hay meadow. About 30 ha is a nature reserve owned by voluntary conservation bodies. The River Shannon is used increasingly for recreational purposes with coarse angling and boating accounting for much of the visitor numbers. Intermittent and scattered damage to the habitats has occurred due to over-deepening of drains and peat silt deposition, water-skiing, ploughing and neglect of hay meadow (or reversion to pasture). However, none of these damaging activities can yet be said to be having a serious impact. Threats to the quality of the site may come from the siting of boating marinas in areas away from centres of population, fertilising of botanically-rich fields, the use of herbicides, reversion of hay meadow to pasture, neglect of pasture and hay meadow, disturbance of birds by boaters, anglers, birdwatchers and the general tourist. The maintenance of generally high water levels in winter and spring benefits all aspects of the flora and fauna, but in this regard, summer flooding is a threat to breeding birds, and may cause neglect of farming.

The Shannon Callows has by far the largest area of lowland semi-natural grassland and associated aquatic habitats in Ireland, and one in which there is least disturbance of natural wetland processes. Botanically, it is extremely diverse with two legally protected species of plants and many scarce species. Excellent examples of two habitats listed on Annex I of the E.U. Habitats Directive occur within the site – *Molinia* meadows and lowland hay meadows with good examples of a further three Annex habitats (two with priority status). In winter the site is internationally important for numbers and species of waterfowl. In spring it feeds large numbers of birds on migration, and in summer it holds very large numbers of breeding waders, rare breeding birds and the endangered Corncrake, as well as a very wide variety of more common grassland and wetland birds. The presence of Otter, an Annex II species, adds further importance to the site.