

Site Name: Castlemaine Harbour SAC

Site Code: 000343

This is a large site located on the south-east corner of the Dingle Peninsula, Co. Kerry. It consists of the whole inner section of Dingle Bay, i.e. Castlemaine Harbour, the spits of Inch and White Strand/Rosbehy and a little of the coastline to the west. The River Maine, almost to Castlemaine, and much of the River Laune catchment, including the Gaddagh, Gweestion, Glanooragh, Cottoner's River and the River Loe, are also included within the site.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

[1130] Estuaries

[1140] Tidal Mudflats and Sandflats

[1210] Annual Vegetation of Drift Lines

[1220] Perennial Vegetation of Stony Banks

[1230] Vegetated sea cliffs of the Atlantic and Baltic coasts

[1310] Salicornia Mud

[1330] Atlantic Salt Meadows

[1410] Mediterranean Salt Meadows

[2110] Embryonic Shifting Dunes

[2120] Marram Dunes (White Dunes)

[2130] Fixed Dunes (Grey Dunes)*

[2170] Dunes with Creeping Willow

[2190] Humid Dune Slacks

[91E0] Alluvial Forests*

[1095] Sea Lamprey (Petromyzon marinus)

[1099] River Lamprey (Lampetra fluviatilis)

[1106] Atlantic Salmon (Salmo salar)

[1355] Otter (Lutra lutra)

[1395] Petalwort (Petalophyllum ralfsii)

Inch Spit holds a fine sand dune system. It is one of the largest and best remaining dune systems in the country. Fore dunes are found on the western side of Rosbehy and Inch. In these younger, mobile dunes, Marram (*Ammophila arenaria*) is common, with Groundsel (*Senecio vulgaris*), Sea Rocket (*Cakile maritima*) and Dandelion

(Taraxacum agg.) also present. Other characteristic species include Sand Couch (Elymus farctus), Lyme-grass (Leymus arenarius) and Sea Spurge (Euphorbia paralias). Fixed dune, a priority habitat under the E.U. Habitats Directive, is well-represented at the site, and in particular towards the tip of Inch Spit. Such areas support species such as Lady's Bedstraw (Galium verum), Common Bird's-foot-trefoil (Lotus corniculatus), Wild Thyme (Thymus praecox), Kidney Vetch (Anthyllis vulneraria), Wild Pansy (Viola tricolor), Biting Stonecrop (Sedum acre), Common Centuary (Centaurium erythraea), Thyme-leaved Sandwort (Arenaria serpyllifolia) and Common Whitlowgrass (Erophila verna), among others. There is also a rich lichen and bryophyte flora. The slightly damper conditions which prevail in dune slacks support Creeping Bent (Agrostis stolonifera), Crested Dog's-tail (Cynosurus cristatus), Glaucous Sedge (Carex flacca), Creeping Willow (Salix repens) and Jointed Rush (Juncus articulatus). The rare bryophyte Petalwort (Petalophyllum ralfsii), which is listed on Annex II of the E.U. Habitats Directive, has been recorded in this system. A smaller spit, with a similar diversity of dune types, occurs at Rosbehy on the southern shore, from where Yellow Centaury (Cicendia filiformis) and Knotted Pearlwort (Sagina nodosa) have been recorded from a dune slack along with other, more common, species.

The sand spits, and also the Coomore peninsula, are underlain by shingle and in places the shingle is exposed and supports a characteristic flora. Species present include Lyme-grass and Sea Sandwort (*Honkenya peploides*). Strandline communities are well-developed along Inch spit, with the exception of the north-western end where recreational pressure is high. Typical species of the strandline include Prickly Saltwort (*Salsola kali*), Sea Rocket, oraches (*Atriplex* spp.) and Sea Sandwort. Two Red Data Book plants, Sea Pea (*Lathyrus japonicus* subsp. *maritimus*) and Sea-kale (*Crambe maritima*), are found associated with the shingle and strandline communities.

The coastline is fringed in many places by saltmarsh. The vegetation here includes Thrift (*Armeria maritima*), Common Saltmarsh-grass (*Puccinellia maritima*), Sea Aster (*Aster tripolium*), Sea Rush (*Juncus maritimus*) and Sea Plantain (*Plantago maritima*). Upper saltmarsh communities extend inland, along estuarine channels, where they are mixed with freshwater communities. Sea Club-rush (*Scirpus maritimus*) and Common Reed (*Phragmites australis*) occur at these locations. Common Cord-grass (*Spartina anglica*) has colonised the lower part of the saltmarsh at Inch and extends out onto the open mudflat. Glassworts (*Salicornia* spp.) occur in association with saltmarsh.

West of Inch, cliffs of glacial drift occur, which support such plants as Ivy (*Hedera helix*), Red Fescue (*Festuca rubra*), Heather (*Calluna vulgaris*), Thrift, Sea Plantain, Sea Mayweed (*Matricaria maritima*), Kidney Vetch and Honeysuckle (*Lonicera periclymenum*). Along the cliff-tops there is coastal grassland with species such as Sweet Vernal-grass (*Anthoxanthum odoratum*), Cock's-foot (*Dactylis glomerata*) and Wood Sage (*Teucrium scorodonia*).

Much of the site consists of intertidal sand and mudflats, supporting a number of soft sediment communities, including beds of eelgrass (mostly *Zostera noltii*) in some

places. A subtidal mixed sediment community complex is also present in the channel between Rossbehy Point and Inch Point.

The rivers and their associated habitats also make up a considerable portion of the site. These associated habitats include wet grassland, woodland, scrub and bog/heath. In the valley up-river of Killorglin, is an interesting area of alluvial wet woodland, dominated by Alder (*Alnus glutinosa*) and willows (*Salix* spp.). The vegetation is quite diverse, and there are spectacular tussocks of Greater Tussock-sedge (*Carex paniculata*). Other species which occur include Ash (*Fraxinus excelsior*), Wild Angelica (*Angelica sylvestris*), Cuckooflower (*Cardamine pratensis*), Meadowsweet (*Filipendula ulmaria*), Common Nettle (*Urtica dioica*), Remote Sedge (*Carex remota*) and a range of bryophytes. While small in area, this is one of the few examples in Ireland of woodland on riverine alluvium dominated by native tree species.

Five plants listed in the Irish Red Data Book have been recorded at this site: Sea-kale, Corn Cockle (*Agrostemma githago*), Sea Pea, Pennyroyal (*Mentha pulegium*) and Irish Lady's-tresses (*Spiranthes romanzoffiana*). The three last-named are legally protected under the Flora (Protection) Order, 1999, as is the rare bryophyte, Petalwort. Other scarce species which occur here are Yellow Bartsia (*Parentucellia viscosa*), Laxflowered Sea-lavender (*Limonium humile*) and Blue-eyed-grass (*Sisyrinchium bermudiana*).

The vicinity of Castlemaine Harbour is also important as one of few areas in Ireland (all of which are in Co. Kerry) where the Natterjack Toad naturally occurs. This amphibian is listed in the Irish Red Data Book and on Annex IV of the E.U. Habitats Directive.

The site also supports a small colony of Common Seal, while two Lamprey species have been recorded in the Laune river catchment. The Laune catchment is used by Otter and is also an important Salmon system with nurseries, riffles pools and glides.

Castlemaine Harbour is a very important site for passage and wintering waterfowl. The following figures are derived from counts between 1994/5 and 1996/7. One species occurs here in internationally important numbers - Brent Goose (734) - with 16 species having populations of national importance: Cormorant (215), Shelduck (129), Pintail (167), Scaup (138), Wigeon (3,513), Red-breasted Merganser (51), Oystercatcher (1,539), Ringed Plover (330), Golden Plover (1,940), Grey Plover (122), Knot (347), Sanderling (207), Dunlin (1,360), Redshank (299), Greenshank (26) and Turnstone (296).

Castlemaine Harbour is of major ecological importance. It contains a range of coastal habitats of excellent quality, including many that are listed on Annex I of the E.U. Habitats Directive, and two which are listed with priority status (fixed dunes and alluvial forests). It also includes long stretches of river and stream which are excellent habitats for Salmon, Lamprey and Otter. Inch dunes are recognised as among the finest in the country, with particularly well-developed dune slacks. The

site supports internationally important waterfowl populations, rare plant species, the rare Natterjack Toad, as well as populations of several animal species that are listed on Annex II of the E.U. Habitats Directive. Part of the site is designated a Special Protection Area (SPA) and is listed as a site under the Ramsar Convention. Part of Castlemaine Harbour is a Statutory Nature Reserve, while Inch and Rosbehy are Wildfowl Sanctuaries.