



**Site Name: Raheenmore Bog SAC**

**Site Code: 000582**

This raised bog developed in a small basin in the catchment of two major river systems i.e. the Brosna and the Boyne. It is situated about 5 km from Daingean in Co. Offaly. The peat is very deep, up to 15 m in places. The bog has a well-developed hummock and hollow system.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (\* = priority; numbers in brackets are Natura 2000 codes):

[7110] Raised Bog (Active)*
[7120] Degraded Raised Bog
[7150] Rhynchosporion Vegetation

Active raised bog comprises areas of high bog that are wet and actively peat-forming, where the percentage cover of bog mosses (*Sphagnum* spp.) is high, and where some or all of the following features occur: hummocks, pools, wet flats, *Sphagnum* lawns, flushes and soaks. Degraded raised bog corresponds to those areas of high bog whose hydrology has been adversely affected by peat cutting, drainage and other land use activities, but which are capable of regeneration. The Rhynchosporion habitat occurs in wet depressions, pool edges and erosion channels where the vegetation includes White Beak-sedge (*Rhynchospora alba*) and/or Brown Beak-sedge (*R. fusca*), and at least some of the following associated species, Bog Asphodel (*Narthecium ossifragum*), sundews (*Drosera* spp.), Deergrass (*Scirpus cespitosus*) and Carnation Sedge (*Carex panicea*).

Raheenmore Bog contains a relatively large wet central core of active raised bog. The hummocks are often colonised by the bog mosses *S. imbricatum* and *S. fuscum*. Pools are well-represented, and it is the pool edges and wet lawns that the Rhynchosporion habitat is best developed. These areas are typically dominated by the bog moss *S. cuspidatum*. The associated vascular plant flora is species-poor, with Bogbean (*Menyanthes trifoliata*), White Beak-sedge, Bog Asphodel, Common Cottongrass (*Eriophorum angustifolium*) and Great Sundew (*Drosera anglica*) being the main species. In places, lawns of *Sphagnum magellanicum* have infilled the pools. Overall, the cover of *Sphagnum* moss on the bog is very good.

Degraded raised bog dominates the marginal areas of the uncut high bog where drainage effects are most pronounced. The vegetation of these degraded areas is still dominated by plant species typical of intact raised bog, though the vegetation tends to be less species-rich than in intact areas and the cover of *Sphagnum* is usually below

25%. The typical dominant species in degraded areas include Heather (*Calluna vulgaris*), Bog Asphodel, Cottongrasses (*Eriophorum* spp.), Deergrass, Cross-leaved Heath (*Erica tetralix*) and Carnation Sedge.

Of note at this site is the fact that, on the western side, mineral springs feeding the lagg zone still survive. A lagg zone is the transitional area at the edge of a bog, between the raised bog peat and the surrounding mineral soils. Conditions are often different here due to the fact that the water in the lagg zone is a mix of water coming from the bog as well as mineral-rich waters from outside. Lagg zones are uncommon features now, due to peat cutting and other land use activities which have removed or altered them significantly in most cases.

The high bog is surrounded by cutover bog. Some sections of old cutover have narrow strips of Downy Birch (*Betula pubescens*) woodland developing. Much of the rest of the cutover is now wet grassland, rich in rushes (*Juncus* spp.) and Purple Moor-grass (*Molinia caerulea*). Common Valerian (*Valeriana officinalis*), Meadowsweet (*Filipendula ulmaria*) and Brown Sedge (*Carex disticha*) can also be found in fields at the bog margins.

In 1959, the very rare Rannock-rush (*Scheuchzeria palustris*), found in its only Irish station in a nearby bog, was transplanted to Raheenmore Bog. However, it has not been recorded recently and may be now extinct.

Raheenmore Bog is within the breeding territory of a pair of Merlin, a scarce species in Ireland and one that is listed on Annex I of the E.U. Birds Directive. Other typical bogland birds which breed here include Red Grouse and Snipe. Red Grouse has declined in Ireland in recent years and is now a Red-listed species.

The structure of the bog habitat has been affected by drainage. This has resulted from peat-cutting along the margins of the bog which has led to the lowering of the water table within the adjoining, intact high bog areas. However, the prospects for the future functioning of the habitat are generally good, as the National Parks and Wildlife Service (NPWS) own much of the site and an extensive programme of drain blocking has taken place. Although the north-eastern section of the bog suffered from burning in the past, the majority of the site is relatively unaffected by this practice at present. Also, peat extraction has largely discontinued.

Raheenmore Bog is a classic example of a midland raised bog and the deepest remaining in Ireland. It is of high conservation importance as it contains good examples of the priority Annex I habitat active raised bog, and the non-priority habitats degraded raised bog and depressions on peat substrates (Rhynchosporion). Most of the site is owned by the NPWS and there has been considerable research and restoration work carried out on the site over the past 15 years. Of particular notes is that this is one of the few raised bogs where restoration of the lagg zone remains feasible.