

Site Name: Lough Ennell SAC

Site Code: 000685

Lough Ennell is a large, open, steep-sided lake, located 3 km south of Mullingar in Co. Westmeath. The lake bottom is of limestone with a marl deposit. The water is markedly alkaline and mesotrophic, possibly owing to effluents received from Mullingar town and to fertilizer inputs from farmland surrounding the lake. The River Brosna flows into the lake from the north at Butler's Bridge, and out from the south.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

[7230] Alkaline Fens

Lough Ennell supports a diverse aquatic flora. Seven stonewort species have been identified, including two Red Data Book species, *Chara denudata* and *C. tomentosa*.

Much of the lakeshore consists of dry, stony ground colonised by calcareous grassland. These areas were formerly part of the lake bed but are now exposed as a consequence of drainage. Species such as Mountain Everlasting (*Antennaria dioica*), Hairy Lady's-mantle (*Alchemilla filicaulis* subsp. *vestita*), Frog Orchid (*Coeloglossum viride*), Fairy Flax (*Linum catharticum*) and Yellow-wort (*Blackstonia perfoliata*) occur here.

Alkaline fen is also found on the lake shore, with species such as Grass-of-parnassus (*Parnassia palustris*), Marsh Pennywort (*Hydrocotyle vulgaris*) and Bottle Sedge (*Carex rostrata*). In wet marshy patches along the shore Marsh-marigold (*Caltha palustris*), Brookweed (*Samolus valerandi*) and Lesser Water-plantain (*Baldellia ranunculoides*) are common.

Reedbeds and species-poor swamp vegetation fringe the lake in places, particularly around the points of inflow and outflow, and on the eastern shore around Tudenham Park. Common Reed (*Phragmites australis*) is abundant here. Water-plantain (*Alisma plantago-aquatica*), Cowbane (*Cicuta virosa*), Frogbit (*Hydrocharis morsus-ranae*) and Tufted-sedge (*Carex elata*) also occur. The latter two species are of note in that they have restricted distributions in Ireland. The rare Fibrous Tussock-sedge (*Carex appropinquata*) has also been recorded from this site.

Mixed woodland of Beech (*Fagus sylvatica*), Ash (*Fraxinus excelsior*) and Downy Birch (*Betula pubescens*) fringes the lakeshore to the north-west. Bluebell (*Hyacinthoides non-*

scripta) and Lords-and-ladies (*Arum maculatum*) are among the woodland ground flora.

Yellow Archangel (*Lamiastrum galeobdolon*), a rare plant listed in the Red Data Book, has been recorded in the woods along the eastern shores of Lough Ennell. This is the only record for this species outside the south-east of Ireland.

The rare Myxomycete fungus, *Licea castanea*, has been recorded from woodland in the site. A species of blue-green alga (*Schizothrix fasciculata*), which forms little pebbles of lime that are cast up on the lakeshore, occurs in Lough Ennell and has not been recorded elsewhere in Ireland.

Scharff's Char (*Salvelinus scharffi*), a distinct race of char which was once found only in Lough Owel and Lough Ennell, is now thought to be extinct. Notable aquatic invertebrates recorded from the lake include *Tinodes maculicornis* (Order Trichoptera), *Metalyte fragilis* (Order Trichoptera), *Limnephilus nigriceps* (Order Trichoptera), *Picromerus bidens* (Order Heteroptera), *Monarthia humili* (Order Hemiptera) and *Donacia obscura* (Order Coleoptera).

This site shares an internationally important Greenland White-fronted Goose flock with Loughs Iron, Glen and Owel. The numbers of geese which visit Lough Ennell are lower than for the other lakes: 91 birds (3 year average peak). Nationally important bird populations which have been recorded on Lough Ennell are: Cormorant (average peak 149; absolute maximum 448); Mute Swan (average peak 424); Pochard (average peak 889; maximum 2,600 on 8/11/85); Tufted Duck (average peak 720) and Coot (average peak 639). All of these data were compiled from counts made over 3 seasons, 1984/85 - 1986/87. A single count of 522 Golden Plover was obtained in that period, constituting a regionally important population.

Lough Ennell is an important amenity area, much used for fishing, boating and camping. Sections of the shoreline are managed for visitor access and amenity. The chemical composition of effluent from the Mullingar sewage treatment plant has a significant impact on the water quality of Lough Ennell. The mid-1970s saw the introduction of treatment of the sewage to reduce phosphates, with a resulting improvement in water quality (according to data compiled during 1987-90). However, levels of planktonic algal growth in the lake water continue to fluctuate, in response to the variable efficiency of the phosphate removal facility at the sewage treatment plant and the re-mobilization of phosphate from the lake sediments.

Lough Ennell is of significance as a highly productive lake which supports a rich variety of lower plant and invertebrate species. Its lakeshore habitats, which include alkaline fen, a habitat listed on Annex I of the E.U. Habitats Directive, support a diverse flora. These habitats also provide important refuges for wildfowl.