

Site Name: Boyne Coast and Estuary SAC

Site Code: 001957

Boyne Coast and Estuary SAC is a coastal site which includes most of the tidal sections of the River Boyne, intertidal sand- and mudflats, saltmarshes, marginal grassland, and the stretch of coast from Bettystown to Termonfeckin that includes the Mornington and Baltray sand dune systems.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

- [1130] Estuaries
- [1140] Tidal Mudflats and Sandflats
- [1210] Annual vegetation of drift lines
- [1310] *Salicornia* Mud
- [1330] Atlantic Salt Meadows
- [2110] Embryonic Shifting Dunes
- [2120] Marram Dunes (White Dunes)
- [2130] Fixed Dunes (Grey Dunes)*

The Boyne River channel, which is navigable and dredged, is defined by training walls, these being breached in places. Intertidal flats occur on the sides of the channelled river. The sediments vary from fine muds in the sheltered areas to sandy muds or sands towards the river mouth. The linear stretches of intertidal flats to the north and south of the river mouth are mainly composed of sand. One or more species of eelgrass (*Zostera* spp.) occur in the estuary.

Parts of the intertidal areas are fringed by saltmarshes, most of which are of the Atlantic type, and dominated by Sea-purslane (*Halimione portulacoides*). Other species present include Common Saltmarsh-grass (*Puccinellia maritima*), Sea Plantain (*Plantago maritima*), Lax-flowered Sea-lavender (*Limonium humile*) and glassworts (*Salicornia* spp.). Common Cord-grass (*Spartina anglica*) occurs frequently on the flats and saltmarshes.

The two sand dune systems in the site, at Baltray and Mornington, are of conservation value, despite the restricted distribution of the intact areas and the high recreational pressure to which they are subjected. A gradient from embryonic dunes to Marram (*Ammophila arenaria*) dunes and then fixed dunes is shown at both systems.

The largest area of annual vegetation of drift lines within this SAC is located at Baltray, north of the estuary. The vegetation is highly representative of the habitat type, which is limited to a small number of highly specialised species that are capable of coping with harsh environmental conditions including high salinity, wind exposure, and unstable substrate and lack of soil moisture. Species present include oraches (*Atriplex* spp.), Sea Rocket (*Cakile maritima*), Prickly Saltwort (*Salsola kali*) and Sea Sandwort (*Honkenya peploides*). Embryonic dunes are particularly well-developed at Baltray where there is active accretion. Species present include Sand Couch (*Elymus farctus*), Lyme-grass (*Leymus arenarius*), Marram, Sea Sandwort and Prickly Saltwort. The embryonic dunes grade into a narrow band of shifting Marram dunes. Marram is dominant, though there are also such species as Cat's-ear (*Hypochoeris radicata*), Mouse-ear Hawkweed (*Hieracium pilosella*) and Dandelion (*Taraxacum* agg.). The areas of fixed dunes on the site have a typical diversity of species, including Marram, Red Fescue (*Festuca rubra*), Wild Carrot (*Daucus carota*), Common Bird's-foot-trefoil (*Lotus corniculatus*), Common Restharrow (*Ononis repens*), Wild Thyme (*Thymus praecox*), Lady's Bedstraw (*Galium verum*) and Wild Pansy (*Viola tricolor*). Vegetation dominated by bryophytes and lichens is limited, though such species as *Brachythecium albicans*, *Hypnum cupressiforme*, *Peltigera canina* and *Cladonia* spp. occur. Some dune slacks may still occur at the site. A number of scarce plants such as Viper's-bugloss (*Echium vulgare*), Adder's-tongue (*Ophioglossum vulgatum*), Variegated Horsetail (*Equisetum variegatum*) and Wild Clary/Sage (*Salvia verbenaca*) have been recorded from the site in the past. The last-named species is of particular note as it is a Red Data Book species at its most northerly known Irish station.

The Boyne is the second most important estuary for wintering birds on the Louth-Meath coastline. From a recent wetland survey carried out over 4 seasons (1994/95-97/98), it is known that this site supports nationally important numbers of Shelduck (176 individuals), Golden Plover (5,338), Lapwing (4,755), Knot (1,559), Black-tailed Godwit (414), Redshank (539), Turnstone (104), Oystercatcher (922), Grey Plover (112) and Sanderling (93).

Other species of regional or local importance include Brent Goose (142), Wigeon (485), Teal (185), Mallard (160), Dunlin (627), Curlew (352) and Ringed Plover (approx. 100). An area of shingle at Baltray Dunes is also an important breeding site for Little Tern, with 14 pairs recorded in 1995. Little Tern is the rarest Irish tern species, and is listed on Annex I of the E.U. Birds Directive. Part of the estuary is a Wildfowl Sanctuary and has been designated a Special Protection Area under the E.U. Birds Directive.

This site has been somewhat modified by human activities. The river is regularly dredged to accommodate cargo ships, which causes disturbance to the bird, fish and invertebrate communities in the estuary. Several factories operate upstream from the estuary and pollution and disturbance associated with them has had an impact on the ecology of the area. There is a proposal to create a deep water facility at the north end of Mornington Dunes on the mouth of the Boyne estuary.

The site is of considerable conservation interest as a coastal complex that supports good examples of eight habitats that are listed on Annex I of the E.U. Habitats Directive, including one which is listed with priority status, and for the important bird populations that it supports.