

**Site Name: The Murrough Wetlands SAC**

**Site Code: 002249**

The Murrough is a coastal wetland complex which stretches for 15 km from Ballygannon to north of Wicklow town, and in parts, extends inland for up to 1 km. A shingle ridge stretches the length of the site and carries the mainline Dublin-Wexford railway.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (\* = priority; numbers in brackets are Natura 2000 codes):

- |  |
|--|
| [1210] Annual Vegetation of Drift Lines    |
| [1220] Perennial Vegetation of Stony Banks |
| [1330] Atlantic Salt Meadows               |
| [1410] Mediterranean Salt Meadows          |
| [7210] <i>Cladium</i> Fens*                |
| [7230] Alkaline Fens                       |

On the seaward side of the shingle bank which runs along The Murrough Wetlands SAC site drift line vegetation includes species such as Sea Rocket (*Cakile maritima*), Sea Sandwort (*Honkenya peploides*), Sea-holly (*Eryngium maritimum*) and Yellow Horned-poppy (*Glaucium flavum*). The rare and legally protected Oysterplant (*Mertensia maritima*) (Flora (Protection) Order, 1999) has been recorded on the gravelly shore in the past but is now considered to be extinct from this locality.

Low sand hills occur at Kilcoole, with Marram (*Ammophila arenaria*) and Lyme-grass (*Leymus arenarius*). In other areas and further inland a rich grassy sward, which is most extensive at the south of the site, has developed. Typical species include Sweet Vernal-grass (*Anthoxanthum odoratum*), Crested Dog's-tail (*Cynosurus cristatus*), Common Bird's-foot-trefoil (*Lotus corniculatus*), Burnet Rose (*Rosa pimpinellifolia*) and Pyramidal Orchid (*Anacamptis pyramidalis*). A community dominated by Silverweed (*Potentilla anserina*) and Strawberry Clover (*Trifolium fragiferum*) occurs in some of the wetter, grassy areas. In some places, particularly at the south of the site, a gorse (*Ulex* sp.) heath has developed on the stony ridge.

Saltmarsh is present within the site in two distinct areas. At the southern end of the site is found Broad Lough. This is a brackish, partly tidal lake, and has a well developed saltmarsh community which includes Saltmarsh Rush (*Juncus gerardi*), Common Saltmarsh-grass (*Puccinellia maritima*), Sea Aster (*Aster tripolium*), Sea Purslane (*Halimione portulacoides*) and Common Scurvygrass (*Cochlearia officinalis*).

Common Reed (*Phragmites australis*) is abundant along the western shore, along with some Sea Club-rush (*Scirpus maritimus*). Saltmarsh is also present in the northern end of the site in the vicinity of The Breaches. Though this has been greatly affected by drainage in the late 1980s and early 1990s, localised Sea Couch (*Elymus pycnanthus*) still occurs. The grassland which was created and improved as a result of the drainage is now influenced by seepage and flooding of saline waters.

Fen vegetation is well developed in the Murrrough wetlands, with both alkaline and calcareous fen with Great Fen-sedge (*Cladium mariscus*) represented. The fens occur mostly between Five Mile Point and Six Mile Point, especially in the townland of Blackditch and also in the Leamore and Grange areas. The alkaline fen is dominated by Black Bog-rush (*Schoenus nigricans*), with Marsh Pennywort (*Hydrocotyle vulgaris*), Purple Moor-grass (*Molinia caerulea*), Devil's-bit Scabious (*Succisa pratensis*), Heather (*Calluna vulgaris*), Cross-leaved heath (*Erica tetralix*), and a wide variety of orchids also present. The rare, Narrow-leaved Marsh-orchid (*Dactylorhiza traunsteineri*) has also been recorded here. Great Fen-sedge occurs in mosaic with several vegetational elements but chiefly with alkaline fen. Its many forms can range from pure stands of Great Fen-sedge, through to occurring as a dominant with Greater Tussock-sedge (*Carex paniculata*) and Blunt-flowered Rush (*Juncus subnodulosus*). *Cladium* fen also occurs at Blackditch within stretches of swamp woodland or fen carr dominated by Rusty Willow (*Salix cinerea* subsp. *oleifolia*) and Downy Birch (*Betula pubescens*).

A fine wet woodland occurs at Blackditch. Downy Birch is the dominant species, with some Alder (*Alnus glutinosa*), willows (*Salix* spp.) and Ash (*Fraxinus excelsior*) also present. The ground flora of this wooded area is often quite dense. This wood also contains a rich invertebrate community with at least eight rare or notable species of fly (Order Diptera) occurring, including *Syntormon setosus*, a species unknown elsewhere in Britain or Ireland.

A wide range of freshwater and brackish marsh habitats occur within the site. These vary from reed-marsh dominated by reeds and rushes (*Juncus* spp.), to those of sedges (*Carex* spp.), with other areas supporting a mixture of sedges and Yellow Iris (*Iris pseudacorus*). A wide variety of grasses and herbs are also found. These include Meadowsweet (*Filipendula ulmaria*), Silverweed and Common Spike-rush (*Eleocharis palustris*). The scarce Red Data Book species Marsh Pea (*Lathyrus palustris*) occurs in one area. The marshes merge into wet grassland in many areas. Where grazing pressure is low, a herb-rich sward occurs with species such as Ragged-Robin (*Lychnis flos-cuculi*), Cuckooflower (*Cardamine pratensis*), Meadowsweet and Heath Spotted-orchid (*Dactylorhiza maculata*) occurring. Sedges are abundant in the wetter areas. Where drains have been cut, there are many other species such as Greater Spearwort (*Ranunculus lingua*), Bogbean (*Menyanthes trifoliata*) and the scarce Reed Sweet-grass (*Glyceria maxima*).

The Murrrough is an important site for wintering waterfowl and breeding birds. Species listed on Annex I of the E.U. Birds Directive include Little Egret, Whooper Swan, Greenland White-fronted Goose, Golden Plover, Kingfisher and Little Tern. Average peak winter counts from 1994/95 - 1997/98 showed the site to have an

internationally important population of Brent Goose (1,318, higher than in the early 1990s), nationally important populations of Wigeon (1,518), Teal (772) and Lapwing (3,140), and regionally or locally important populations of Whooper Swan (80), Little Grebe (22), Shelduck (95), Gadwall (9), Mallard (391), Shoveler (22), Golden Plover (615), Curlew (605) and Redshank (181). Greylag Goose numbers were nationally important in the early 1990s but these numbers have dropped off. The average peak is now 213.

Little Tern breed on the shingle beach near The Breaches and this is the largest colony on the east coast (approx. 50 pairs in 1993, an average of 37 pairs over the ten year period 1988-1998 ). Redshank, Oystercatcher, Ringed Plover and Water Rail also breed. The reedbeds at Broad Lough provide habitat for Reed Warbler and the rare Bearded Tit has bred here.

Otter has been reported regularly from the Murrough. This is a Red Data Book Species, and is also listed on Annex II of the Habitats Directive.

Recent farming and drainage practices and afforestation have greatly reduced the area and quality of the wetlands habitats - the area between Kilcoole and Newcastle is particularly affected. In 1997 there was some levelling of the sand hills below Killougher station. Pollution, reclamation and further drainage would adversely affect this site. A section of the wetlands at Blackditch, which includes alkaline and *Cladium* fen, has been acquired by BirdWatch Ireland and is being managed for nature conservation.

This site is of importance as it is the largest coastal wetland complex on the east coast of Ireland. Although much affected by drainage, it still contains a wide range of coastal and freshwater habitats, including six listed on Annex I of the E.U. Habitats Directive, some of which contain threatened plants. Areas on the site contain a rich invertebrate fauna, including several rarities. It is an important site for both wintering and breeding birds and supports a variety of species listed on Annex I of the E.U. Birds Directive.