



Site Name: Lough Swilly SAC

Site Code: 002287

This large site, situated in the northern part of Co. Donegal, comprises the inner part of Lough Swilly. It extends from below Letterkenny to just north of Buncrana. Lough Swilly is a long sea lough, cutting through a variety of metamorphic rocks on the west side of Inishowen. The main rivers flowing into the site are the Swilly, Lennan and Crana. At low tide, extensive sand and mudflats are exposed, especially at the mouths of the Swilly and Lennan rivers. The site is estuarine in character, with shallow water and intertidal sand and mudflats being the dominant habitats.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

- [1130] Estuaries
- [1150] Coastal Lagoons*
- [1330] Atlantic Salt Meadows
- [6410] *Molinia* Meadows
- [91A0] Old Oak Woodlands
- [1355] Otter (*Lutra lutra*)
- [1351] Harbour Porpoise (*Phocoena phocoena*)

Ecological communities present in the intertidal sediments at Lough Swilly SAC include fine sand community complexes, intertidal mixed sediment communities with polychaetes, subtidal mixed sediment communities with polychaetes and bivalves, muddy fine sand communities with *Thyasira flexuosa*, muddy community complexes and *Ostrea edulis* dominated communities. Bivalves and polychaete worms are well represented in the macro-invertebrate fauna, with species such as Cockles (*Cerastoderma edule*), Mussels (*Mytilus edulis*), Baltic Tellin (*Macoma balthica*), Ragworm (*Nereis diversicolor*) and Sand Mason (*Lanice conchilega*) being common. Common Cord-grass (*Spartina anglica*) is well established on parts of the intertidal flats. The shoreline above the flats varies from bedrock shore to shingle or cobbles, and here is found a scattering of salt tolerant plants such as Common Scurvygrass (*Cochlearia officinalis*), Sea-milkwort (*Glaux maritima*) and Red Fescue (*Festuca rubra*).

Saltmarshes are well represented in the inner sheltered areas of the site, with good examples in the Ramelton area. The marshes are the Atlantic salt meadow type, and are characterised by such species as Thrift (*Armeria maritima*), Sea-milkwort, Sea Aster (*Aster tripolium*), Sea Arrowgrass (*Triglochin maritima*) and Red Fescue.

Lakes which are lagoonal in character occur at Inch and Blanket Nook. Inch Lough is a good example of a large, shallow lagoon with very low salinity in most of the lagoon. Less information is available for Blanket Nook but it is of a higher salinity and adds to the richness of the habitat within the site as a whole. The vegetation in Inch is diverse and typically lagoonal, with well-developed charophyte communities, including a large population of *Chara canescens* (a Red Data Book species). It also supports Horned Pondweed (*Zannichellia palustris*) and a mixed Pondweed/Tassleweed community (*Potamogeton pectinatus/Ruppia maritima*). The green alga *Chaetomorpha linum* occurs at Blanket Nook. The aquatic fauna at Inch is rich and diverse and includes a range of lagoonal specialists and species that are apparently rare. These include *Lekanesphaera hookeri* (Order Isopoda), *Palaemonetes varians* (Order Decapoda), *Conopeum seurati* (a bryozoan), *Sigara stagnalis* (Order Hemiptera), *S. concinna* (Order Hemiptera), *Jaera nordmanni* (Order Isopoda), *Neomysis integer* (Order Mysida) and *Cordylophora caspia* (a hydrozoan). Less information is available on the aquatic fauna at Blanket Nook, though several lagoonal specialists have been recorded, including *Jaera ischiosetosa* (Order Isopoda), which appears to be a rare species in Ireland. Inch Lough is regarded as being of significant conservation value for ecotonal Coleoptera, with several species indicative of well-developed shoreline habitats including two that are apparently rare: *Bembidion bipunctatum*, a halo-tolerant shore species, and *Bembidion aeneum*, a stenotopic halobiont species. Although artificial in origin, the lagoon habitat in this site is one of the largest and best examples of a shallow, low salinity lagoon habitat in the country.

Over 11 hectares of *Molinia* Meadows, a habitat listed on Annex I of the E.U. Habitats Directive, are reported to occur at Inch Level, according to the Irish Semi-natural Grasslands Survey, 2010.

Two woodlands occur adjacent to the north-western shore of Lough Swilly. These are Rathmullen and Carradoan Woods, the former being a Nature Reserve. They are dominated by Sessile Oak (*Quercus petraea*) and Downy Birch (*Betula pubescens*), but many other species are present including exotics such as Beech (*Fagus sylvatica*). Alder (*Alnus glutinosa*) and willows (*Salix* spp.) occur in the wetter parts. An area of wet heath, dominated by Heather (*Calluna vulgaris*), occurs on the hill above Carradoran Wood. These woodlands display a generally intact structure and support a range of breeding birds, including Woodcock.

A further area of woodland, scrub and heath occurs above the north-east shore at Crockacashel and at Porthaw. Oak/Hazel (*Corylus avellana*) scrub is frequent, and there is a well-developed ground flora with species such as Wood-sorrel (*Oxalis acetosella*), Herb-Robert (*Geranium robertianum*), Lady-fern (*Athyrium filix-femina*) and Broad Buckler Fern (*Dryopteris dilatata*). The heath vegetation includes Heather, Gorse (*Ulex europaeus*), Wood Sage (*Teucrium scorodonia*) and Tormentil (*Potentilla erecta*). In the wetter areas, Purple Moor-grass (*Molinia caerulea*) and Cross-leaved Heath (*Erica tetralix*) occur.

The site supports a population of Otter, a species listed on Annex II of the E.U. Habitats Directive.

Lough Swilly is an important site for waterfowl in autumn and winter. The shallow waters provide suitable habitat for grebes and diving duck, while the intertidal flats are used by an excellent diversity of wildfowl and waders. At high tide, the duck and wader species roost on the saltmarshes and shorelines, with some species moving to the adjacent pasture and arable fields. In the three winters 1994/95 to 1996/97, 16 species occurred in nationally important numbers as follows (figures are average maximum counts for the 3 winters): Great Crested Grebe (274), Shelduck (646), Wigeon (1,673), Teal (1,381), Mallard (1,155), Shoveler (58), Scaup (143), Goldeneye (169), Red-breasted Merganser (103), Coot (335), Oystercatcher (1,459), Knot (327), Dunlin (7,995), Curlew (1,716), Redshank (1,080) and Greenshank (30). Other species which occur in regionally or locally important numbers, and at times may exceed the threshold for national importance, include Brent Goose, Pochard, Tufted Duck, Lapwing, Ringed Plover, Grey Plover, Bar-tailed Godwit and Turnstone. The site is also an important area for the Great Northern Diver and the rare Slavonian Grebe.

The adjacent pasture and arable polders at Inch, Big Isle and Blanket Nook support internationally important populations of Whooper Swans, Greenland White-fronted Goose and Greylag Goose. Inch Lough is an important roosting area for these birds and at times they utilise other parts of Lough Swilly for roosting.

This site is of conservation importance as it contains good examples of at least five habitats listed on Annex I of the E.U. Habitats Directive (estuaries, lagoons, Atlantic salt meadows, *Molinia* meadows, old oak woods) and supports a population of Otter. In addition, it is of high ornithological importance for wintering waterfowl, with 16 species occurring regularly in numbers of national importance, plus three species occurring within the site and on adjacent polders in numbers of international importance.