

Site Name: Brown Bog SAC

Site Code: 002346

Brown Bog NHA is located 5 km north-west of Longford town, mainly in the townlands of Tully, Lissanurlan and Cartronlebagh. The site comprises a raised bog that includes both areas of high bog and cutover bog. The bog margins are mainly surrounded by scrub/woodland.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (* = priority; numbers in brackets are Natura 2000 codes):

[7110] Raised Bog (Active)*
[7120] Degraded Raised Bog
[7150] Rhynchosporion Vegetation

Active raised bog comprises areas of high bog that are wet and actively peat-forming, where the percentage cover of bog mosses (*Sphagnum* spp.) is high, and where some or all of the following features occur: hummocks, pools, wet flats, *Sphagnum* lawns, flushes and soaks. Degraded raised bog corresponds to those areas of high bog whose hydrology has been adversely affected by peat cutting, drainage and other land use activities, but which are capable of regeneration. The Rhynchosporion habitat occurs in wet depressions, pool edges and erosion channels where the vegetation includes White Beak-sedge (*Rhynchospora alba*) and/or Brown Beak-sedge (*R. fusca*), and at least some of the following associated species, Bog Asphodel (*Narthecium ossifragum*), sundews (*Drosera* spp.), Deergrass (*Scirpus cespitosus*) and Carnation Sedge (*Carex panicea*).

This site is situated in a drumlin-filled valley and consists of a small raised bog characterised by a central wet depression with quaking mats of bog mosses and tear pools colonised by algae. Water flows through the pools and it is possible that there is a spring located in the bog centre. A flush area occurs in the north. Abandoned cutover is found around the northern, western and north-eastern bog margins. Remnant old deciduous woodland occurs to the north-west.

The site supports typical Midland Raised Bog communities, which include Heather (*Calluna vulgaris*), Carnation Sedge, Bog-rosemary (*Andromeda polifolia*) and occasional Cranberry (*Vaccinium oxycoccos*). The high bog supports extensive quaking carpets of bog mosses including *Sphagnum magellanicum*, *S. papillosum* and *S. capillifolium*. Pools occur frequently and support *Sphagnum auriculatum*, Bogbean (*Menyanthes trifoliata*) and Great Sundew (*Drosera anglica*). Bare pools and algal pools are also found. Hummocks of *Sphagnum imbricatum* and *S. fuscum* occur. The high

bog is drier around the margins, where Heather and lichens (*Cladonia* spp.) dominate. Scattered Downy Birch (*Betula pubescens*) occurs in association with the northern flush, along with Soft Rush (*Juncus effusus*). Quaking flats of Bog Asphodel and bog moss lawns dominate the inter-pool areas of the flush. One pool with obvious water flow supports Bog Pondweed (*Potamogeton polygonifolius*). Old cutover is mainly colonised by Gorse (*Ulex europaeus*), Downy Birch, Scots Pine (*Pinus sylvestris*) and Purple Moor-grass (*Molinia caerulea*). In the north-west, old deciduous woodland with Downy Birch, Scots Pine, Rowan (*Sorbus aucuparia*) and occasional the Beech (*Fagus sylvatica*) is found.

There are few land uses associated with this site. There are no high bog drains and only two sets of marginal drains are present in the cutover to the north-west. At present there is no active peat-cutting on the site. A large area of cutover to the east of the site has been recently afforested with Sitka Spruce (*Picea sitchensis*). The majority of the bog has not been burnt for some time, although recent localised burning has taken place along the southern margin. Overall there has been little damage to this bog, with only small areas of cutover present. Most of the extent of the original peat basin appears to be remaining. However, peat-cutting and burning are the two main threats to the site.

Brown Bog is a site of considerable conservation significance as it comprises a relatively little-damaged raised bog, a rare habitat in the E.U. and one that is becoming increasingly scarce and under threat in Ireland. Although the site is small it supports a good diversity of raised bog microhabitats including hummock/hollow complexes, pools and a flush system with surrounding tear pool complex, along with cutover which adds to the diversity and scientific value of the site. Active raised bog is listed as a priority habitat on Annex I of the E.U. Habitats Directive. Priority status is given to habitats and species that are threatened throughout the E.U. Ireland has a high proportion of the E.U. resource of this habitat type (over 60%) and so has a special responsibility for its conservation at an international level.