

**Hook Head SAC (site code: 764)
Conservation objectives supporting document
-coastal habitats**

Version 1

July 2011

Table of Contents

	Page No.
1 Introduction	2
2 Conservation objectives	2
3 Vegetated sea cliffs	2
3.1 Overall objective	4
3.2 Area	4
3.2.1 Habitat extent	4
3.3 Range	5
3.3.1 Habitat distribution	4
3.4 Structure and Functions	4
3.4.1 Functionality and hydrological regime	4
3.4.2 Vegetation structure: zonation	5
3.4.3 Vegetation structure: vegetation height	5
3.4.3 Vegetation composition: typical species & sub-communities	5
3.4.4 Vegetation composition: negative indicator species	6
3.4.5 Vegetation composition: bracken and woody species	6
4 References	7
Appendix I: Map of Hook Head showing the distribution of vegetated sea cliffs as identified by the Irish Sea Cliff Survey	8

1 Introduction

Achieving Favourable Conservation Status (FCS) is the overall objective to be reached for all Annex I habitat types and Annex II species of European Community interest listed in the Habitats Directive 92/43/EEC (Commission of the European Communities, 2003). It is defined in positive terms, such that a habitat type or species must be prospering and have good prospects of continuing to do so.

Hook Head SAC (site code: 764) is designated for a range of marine and coastal habitats including reefs and sea cliffs. The following coastal habitat is included as a qualifying interest for the site:

- Vegetated sea cliffs of the Atlantic and Baltic coasts (1230)

This backing document sets out the conservation objectives for the vegetated sea cliffs in Hook Head SAC, which is defined by a list of parameters, attributes and targets. The main parameters are (a) Area, (b) Range and (c) Structure and Functions, the latter of which is broken down into a number of attributes, including functionality, vegetation structure and vegetation composition.

The targets set for the vegetated sea cliffs is based in part on the findings of the Irish Sea Cliff Survey (ISCS) (Barron *et al.*, 2011) and this document should be read in conjunction with that report.

The Irish Sea Cliff Survey (ISCS) identified the following three sub-sites within Hook Head SAC:

1. Loftushall (west side of Hook Peninsula)
2. Hook Head (tip of Hook Peninsula)
3. Baginbun Head (northwest side of Hook Peninsula)

These sub-sites were not visited during the Irish Sea Cliff Survey (Barron *et al.*, 2011) but they were assessed using aerial photographs, as part of the desk study. Therefore, while their extent was measured, no information was obtained on the vegetation present. Information on the vegetation was obtained from the Natura 2000 assessment conducted for this site.

The conservation objective for the vegetated sea cliff habitat within the entire SAC is therefore quite generic and may be adjusted in the future in light of new information.

2 Conservation Objectives

The conservation objective aims to define the favourable conservation condition of a habitat or species at a particular site. Implementation of these objectives will help to ensure that the habitat or species achieves favourable conservation status at a national level.

3 Vegetated sea cliffs

Sea cliffs can be broadly divided into two categories: hard (or rocky) cliffs and soft (or sedimentary) cliffs, both of which are covered by the Annex I habitat 'vegetated sea cliffs of the Atlantic and Baltic coasts'. Hard cliffs are composed of rocks such as limestone, sandstone, granite or quartzite which are hard and relatively resistant to mechanical erosion. Soft cliffs are composed of softer rock, such as shale, or unconsolidated material, such as glacial till. Vegetation of hard sea cliffs in exposed situations exhibits a strong maritime influence and is relatively stable. Soft cliff habitats are more prone to slope failure, which results in the presence of fast-colonising pioneer species.

Defining the limits of what constitutes a sea cliff is problematic and a number of different interpretations have been used in the past (Fossitt, 2000; Commission of the European Communities, 2003; JNCC, 2004; Browne, 2005). In order to address any inconsistencies, the following definition for sea cliffs was developed and used during the Irish Sea Cliff Survey (Barron *et al.*, 2011):

“A sea cliff is a steep or vertical slope located on the coast, the base of which is in either the intertidal (littoral) or subtidal (sublittoral) zone. The cliff may be composed of hard rock such as basalt, or of softer substrate such as shale or boulder clay. Hard cliffs are at least 5m high, while soft cliffs are at least 3m high. The cliff top is generally defined by a change to an obvious less steep gradient. In some cases the cliff may grade into the slopes of a hillside located close to the coast. In these cases the cliff is defined as that part of the slope which was formed by processes of coastal erosion, while the cliff top is where there is the distinct break in slope. Both the cliff and the cliff top may be subject to maritime influence in the form of salt spray and exposure to coastal winds. A cliff can ascend in steps with ledges, and the top of the cliff is taken to occur where erosion from wave action is no longer considered to have been a factor in the development of the landform. The cliff base may be marked by a change in gradient at the bottom of the cliff. Where the base is exposed it can be characterised by scree, boulders, a wave-cut platform or sand, among other substrates. During this survey where cliffs occur within the subtidal zone the base was considered to be the high water mark. A cliff is considered to have reached its end point where it is no longer over 5m high (hard cliffs) or 3m high (soft cliffs), or no longer has a steep slope. To be considered in this study, a cliff had to be a minimum of 100m in length. Sea cliffs may support a range of plant communities such as grassland, heath, scrub and bare rock communities, among others.”

The Hook peninsula is a long, narrow, low-lying headland that protrudes into the sea in a south-southwest direction on the eastern side of Waterford Harbour. There are approximately 15km of sea cliffs composed of a mixture of clay and rock. In some places the clay overlies the rock. The geology of the area is of high interest as it represents a fine example of the junction between Devonian Old Red Sandstone and overlying Carboniferous limestone. The cliffs are mostly low (usually not more than 10m) but rise to 20m at Hook Head and 30m towards Baginbun Head. Below the rocky cliffs a well developed bedrock shore occurs, which is often wide and contains a series of platforms. Fossils are a feature of the limestone rock formations.

The cliffs are predominantly hard but in some places are overlain by soft rock. They support a typical maritime flora and habitat for a diversity of cliff nesting birds including peregrine falcon (*Falco peregrinus*) and chough (*Pyrrhocorax pyrrhocorax*), the latter at one of their most easterly stations.

3.1 Overall Objective

The overall objective for ‘vegetated sea cliffs of the Atlantic and Baltic coasts’ in Hook Head SAC is to ‘maintain the favourable conservation condition’. This objective is based on an assessment of the current condition of the habitat under a range of attributes and targets. The assessment is divided into three main headings (a) Area, (b) Range and (c) Structure and Functions.

3.2 Area

3.2.1 Habitat extent

Habitat extent is a basic attribute to be assessed when determining the condition of a particular habitat. The target is for the ‘*area to remain stable*’. Bearing in mind that coastal systems are naturally dynamic and subject to change even within a season, this target is assessed subject to natural processes, including erosion and succession.

The distribution of vegetated sea cliffs as identified during the Irish Sea Cliff Survey (ISCS) (Barron *et al.*, 2011) is shown on a map in Appendix I. Rocky sea cliffs run from Harrylock on the western side of Hook Head to Baginbun Head on the north-eastern side of the peninsula.

As cliffs are linear features on maps, their extent is measured in kilometres rather than hectares, as you would with other habitats. During the ISCS (Barron *et al.*, 2011), each cliff was divided into sections based on physical characteristics and vegetation cover. Breaks (i.e. non-cliff areas) of between 80m and 500m along a length of cliff were discounted from the calculations. The total length of the cliff sections within each sub-site in Hook Head as measured by Barron *et al.* (2011) is presented in the following table. The area of each cliff sub-site that is located within the SAC boundary is also presented.

There are a number of differences in the two sets of figures below. Most of the differences are explained by the fact that the ISCS mapped the total sea cliff resource at each site and not all of the sea cliff mapped is contained within the SAC boundary. In addition, the County boundary line was used to draw the line for the ISCS, while a different mapping dataset than was used to draw the SAC boundary. As a result the length of cliff inside the SAC boundary may be under-estimated. The total length of cliff sections recorded by the ISCS in the Hook Head area was 13.36km. However, when this dataset was clipped to the SAC boundary 12.11km was included within the boundary. However, in reality this figure is likely to be higher as a result of these mapping anomalies.

Site name	Total area/length (km) of sea cliff from ISCS	Total area/length (km) of sea cliff within SAC boundary
Loftushall	0.63	0.55
Hook Head	2.35	2.36
Baginbun Head	10.38	9.20
Totals	13.36	12.11

3.3 Range

3.3.1 Habitat distribution

The distribution of sea cliffs throughout Hook Head SAC as identified by the Irish Sea Cliff Survey (Barron *et al.*, 2011) is presented in Appendix I. The hard cliffs in Hook Head SAC are not likely to be redistributed through natural processes, unlike other more dynamic coastal systems such as sand dunes or saltmarshes.

3.4 Structure and Functions

A fundamental aim of sea cliff conservation is to facilitate some degree of natural mobility through slumping. Sea cliffs can be of geomorphological interest as well as ecological interest and erosion can also expose geological features of interest.

3.4.1 Functionality and hydrological regime

Coastal protection works can disrupt the natural integrity of a sea cliff. The health and on-going development of vegetated sea cliffs relies on natural processes such as erosion continuing without any impingement. This is generally a bigger issue for soft cliffs that require a degree of slumping and erosion to expose bare soil for pioneer species to colonise; otherwise, the vegetation is replaced by hardy grasses, and scrub of little conservation value can develop. In addition, cliff erosion provides an important sediment source to sites further along the coast (e.g. sand dunes). Preventing erosion at a cliff site can lead to beach starvation at another site.

The target is to maintain, or where necessary restore, the natural geomorphological processes without any physical obstructions and the local hydrological regime, including groundwater quality.

3.4.2 Vegetation structure: zonation

Ecological variation in this habitat type depends on a number of physical and biological factors, in particular climate, degree of exposure to sea-spray, geology and soil type, as well as the level of grazing and sea bird activity. The vegetation above most of the cliff length has been improved and transitions into semi-natural habitats such as maritime grassland are rare.

The target is to maintain the sea cliff habitat, as well as transitional zones, including those to terrestrial communities.

3.4.3 Vegetation structure: vegetation height

A varied vegetation structure is important for maintaining species diversity and is particularly important for invertebrates and birds. Grazing increases the species diversity and is particularly important for maritime grasslands and coastal heath, which are often associated with sea cliffs. The target is to maintain the structural variation in the sward height.

3.4.4 Vegetation composition: typical species & sub-communities

Different sea cliff communities develop in a number of habitat zones related to the degree of maritime influence (exposure to wind and sea spray), geology and soil type. In general Irish sea cliffs display a range of zones running in a series of horizontal bands up the cliff face, each of which has its own distinct sub-communities including:

- Splash zone
- Pioneer zone
- Rock crevice/cliff ledge zone
- Maritime grassland zone
- Maritime heath zone
- Maritime slope flush zone

There is considerable variation but the general pattern would be that the maritime influence is strongest near the base of the cliff and becomes gradually less dominant towards the cliff top. At the cliff base, the vegetation is naturally very open and the species present have a high tolerance to salinity. The splash zone generally has a well developed lichen flora dominated by species such as *Verrucaria maura*, *Ramalina* spp. And *Xanthoria* spp. These plant communities are dependent on rock crevices for rooting. Moving up the cliff, between the splash zone and the cliff top, vegetation on cliff ledges is less open and can support some species that are not exclusively associated with coastal conditions. Closer to the cliff top, maritime grasslands can occur. The plant communities and physical characteristics of maritime grasslands vary depending on the degree of exposure and whether or not grazing is a factor. Plant communities typical of sea bird cliffs and maritime therophyte communities are exceptions to this horizontal zonation and can occur as a mosaic with the other plant communities. The following table presents lists of species that are considered typical of the different zones associated with hard cliffs by Barron *et al.* (2011), such as those found at Hook Head.

Typical splash zone species			
<i>Ramalina</i> spp.	<i>Verrucaria maura</i>	<i>Xanthoria</i> spp.	
Typical crevice & ledge species			
<i>Anthyllis vulneraria</i>	<i>Asplenium marinum</i>	<i>Armeria maritima</i>	<i>Aster tripolium</i>
<i>Atriplex prostrata</i>	<i>Beta vulgaris</i> ssp. <i>maritima</i>	<i>Cerastium diffusum</i>	<i>Lavatera arborea</i>
<i>Catapodium marinum</i>	<i>Crithmum maritimum</i>	<i>Festuca rubra</i>	<i>Limonium</i> sp.
<i>Inula crithmoides</i>	<i>Ligusticum scoticum</i>	<i>Plantago coronopus</i>	<i>Plantago maritima</i>
<i>Sedum anglicum</i>	<i>Spergularia rupicola</i>	<i>Sedum rosea</i>	<i>Silene uniflora</i>
Typical coastal heath species			
<i>Calluna vulgaris</i>	<i>Daboecia cantabrica</i>	<i>Empetrum nigrum</i>	<i>Erica cinerea</i>
<i>Erica tetralix</i>	<i>Vaccinium myrtillus</i>	<i>Scilla verna</i>	<i>Ulex gallii</i>
Typical maritime grassland species			
<i>Anthyllis vulneraria</i>	<i>Armeria maritima</i>	<i>Daucus carota</i>	<i>Festuca rubra</i>
<i>Crithmum maritimum</i>	<i>Hyacinthoides non-scripta</i>	<i>Plantago coronopus</i>	<i>Plantago maritima</i>
<i>Sedum anglicum</i>	<i>Spergularia rupicola</i>	<i>Silene uniflora</i>	<i>Scilla verna</i>

The cliffs in Hook Head SAC are thought to support maritime vegetation associated with hard cliffs that is typical for the south-east of the country. Species present include sea pink (*Armeria maritima*), scurvy grass (*Cochlearia* spp.), red fescue (*Festuca rubra*), sea campion (*Silene maritima*), sea mayweed (*Tripleurospermum maritimum*), plantains (*Plantago maritima*, *P. coronopus*), sea samphire (*Crithmum maritimum*), rock sea spurry (*Spergularia rupicola*) and rock sea-lavenders (*Limonium binervosum*). Bramble (*Rubus fruticosus*) usually occurs at the top of the cliffs but there is mostly an abrupt transition to improved grassland in the fields above the cliffs.

The target for this attribute is to ensure that the typical flora of vegetated sea cliffs is maintained, as are the range of sub-communities within the different zones.

3.4.5 Vegetation composition: negative indicator species

Negative indicator species can include non-native species (e.g. *Hebe* spp., *Carpobrotus edulis*, *Gunnera tinctoria*); species indicative of changes in nutrient status (e.g. *Urtica dioica*) and species not considered to be typical of the habitat (e.g. *Pteridium aquilinum*).

The target for this attribute is that negative indicator species (including non-native species) should make up less than 5% of the vegetation cover.

3.4.6 Vegetation composition: bracken and woody species

Encroachment of bracken (*Pteridium aquilinum*) and woody/scrub species on cliffs, particularly the maritime grasslands and coastal heath leads to a reduction in species diversity.

The target for this attribute is that in the case of maritime grassland an/or heath, bracken should make up less than 10% of the vegetation cover, while woody species should make up no more than 20% of the vegetation cover.

4 References

Barron, S., Delaney, A., Perrin, P., Martin, J. and O'Neill, F. (2011). National survey and assessment of the conservation status of Irish sea cliffs. *Irish Wildlife Manuals*, No. 53. National Parks and Wildlife Service, Department of the Environment, Heritage and Local Government, Dublin, Ireland.

Browne, A. (2005). *National inventory of sea cliffs and coastal heaths*. Unpublished report to the National Parks and Wildlife Service, Dublin.

Commission of the European Communities (2003). *Interpretation Manual of European Union Habitats – EUR 25*. DG Environment-Nature and Biodiversity, Brussels.

Fossitt, J.A. (2000). *A guide to habitats in Ireland*. The Heritage Council, Kilkenny.

JNCC (2004). *Common standards and monitoring guidance for maritime cliff and slope habitats*. Joint Nature Conservation Committee, Peterborough, UK.

Appendix I – Map of Hook Head showing the distribution of vegetated sea cliffs as identified by the Irish Sea Cliff Survey (based on data from Barron *et al.*, 2011).

