

National Parks & Wildlife Service

Garriskil Bog SAC  
(site code 000679)

**Conservation objectives supporting document -  
raised bog habitats**

Version 1

October 2015

# Contents

<b>1</b>	<b>INTRODUCTION.....</b>	<b>2</b>
1.1	RAISED BOGS.....	2
1.1.1	<i>Raised Bogs Microtopography.....</i>	3
1.1.2	<i>Typical Flora of Irish Raised Bogs.....</i>	5
1.1.3	<i>Typical Fauna of Irish Raised Bogs.....</i>	6
1.2	HABITATS DIRECTIVE RAISED BOG HABITATS IN IRELAND.....	9
1.2.1	<i>Restoration of Active Raised Bog in Ireland.....</i>	10
1.3	GARRISKIL BOG SAC .....	10
1.3.1	<i>Flora of Garriskil Bog.....</i>	11
1.3.2	<i>Fauna of Garriskil Bog.....</i>	12
<b>2</b>	<b>CONSERVATION OBJECTIVES.....</b>	<b>13</b>
2.1	AREA .....	13
2.2	RANGE .....	14
2.3	STRUCTURE AND FUNCTIONS.....	15
2.3.1	<i>High bog area .....</i>	15
2.3.2	<i>Hydrological regime: water levels.....</i>	15
2.3.3	<i>Hydrological regime: flow patterns.....</i>	17
2.3.4	<i>Transitional areas between high bog and surrounding mineral soils (includes cutover areas).....</i>	17
2.3.5	<i>Vegetation quality: central ecotope, active flush, soaks.....</i>	18
2.3.6	<i>Vegetation quality: microtopographical features.....</i>	19
2.3.7	<i>Vegetation quality: bog moss (Sphagnum) species .....</i>	19
2.3.8	<i>Typical species: bog flora.....</i>	20
2.3.9	<i>Typical species: bog fauna.....</i>	20
2.3.10	<i>Elements of local distinctiveness .....</i>	21
2.3.10.1	Site features.....	21
2.3.10.2	Rare flora .....	21
2.3.10.3	Rare fauna.....	21
2.3.11	<i>Negative physical indicators.....</i>	21
2.3.12	<i>Vegetation composition: native negative indicator species.....</i>	22
2.3.13	<i>Vegetation composition: non-native invasive species .....</i>	22
2.3.14	<i>Air quality: nitrogen deposition .....</i>	22
2.3.15	<i>Water quality.....</i>	23
<b>3</b>	<b>REFERENCES .....</b>	<b>25</b>

**Map 1: Extent of potential active raised bog on Garriskil Bog.**

**Map 2: Distribution of raised bog ecotopes on Garriskil Bog.**

**Map 3: Digital elevation model and drainage patterns at Garriskil Bog.**

## 1 Introduction

This document presents a summary of the background information that has informed the process of setting the Site-Specific Conservation Objective in relation to the priority Annex I habitat 'active raised bog' (habitat code 7110) (hereafter referred to as Active Raised Bog (ARB)), for which Garriskil Bog Special Area of Conservation (SAC) has been designated.

Garriskil Bog SAC is also designated for two other related Annex I habitats, namely; 'degraded raised bogs still capable of natural regeneration' (habitat code 7120) (hereafter referred to as Degraded Raised Bog (DRB)) and 'depressions on peat substrates of the Rhynchosporion' (habitat code 7150). Based on the close ecological relationship between these three habitats types, it is not necessary to set specific Conservation Objectives for all three habitats individually. It is considered that should favourable conservation condition for ARB be achieved on the site, then, as a consequence, favourable conservation condition for the other two habitats would also be achieved.

### 1.1 Raised Bogs

Raised bogs are accumulations of deep peat (typically 3-12m) that originated in shallow lake basins or topographic depressions. The name is derived from the elevated surface, or dome, that develops as raised bogs grow upwards through the accumulation of peat; the domed effect is often exaggerated when the margins of a bog are damaged by turf cutting or drainage, and are drying out. Raised bogs are most abundant in the lowlands of central and mid-west Ireland.

Irish raised bogs are classified into two sub-types (Schouten 1984): 1. Western or intermediate raised bogs, and 2. True midland or eastern raised bogs, based on phytosociological and morphological characteristics. In terms of overall morphology, the main difference between these two raised bog types is that while eastern raised bogs tended to stay more confined to the depressions in which they were formed, western raised bogs tended to grow out beyond their original basin, presumably a result of the higher rainfall levels (Cross 1990). In terms of vegetation differences the most obvious difference between the two bog types is the presence of a number of oceanic plant species on western raised bogs which are absent from the true midland raised bogs. The liverwort species *Pleurozia purpurea*<sup>1</sup> and the moss species *Campylopus atrovirens* grow on western raised bogs but not on eastern raised bogs; similarly, *Carex panicea* is generally more common on the high bog surfaces of western raised bogs (Schouten 1984). All of these plant species are widespread in the low-level Atlantic blanket bogs and their presence in western raised bogs is presumed to be due to the higher rainfall levels and greater rain-derived nutrient fluxes.

Exploitation has been extensive and none of the remaining Irish raised bogs are completely intact (Cross 1990). It is estimated that less than 10% of the original raised bog habitat in Ireland is in a near intact state (uncut), with less than 0.5% continuing to support ARB (DAHG 2014). Excavated face banks, whether active or inactive, are a common feature around the margins. Any areas where part of the bog has been removed are termed cutover bog, with the remaining area referred to as high bog or intact bog. In a natural state, raised bogs are circled by a wetland fringe, known as the lagg zone, which is usually characterised by fen communities. In Ireland, most lags have been lost through drainage and land reclamation (Fossitt 2000).

---

<sup>1</sup> Note on species nomenclature: *In the case of plant species, only scientific names are used throughout the main text while common English names are included in tables. In the case of faunal species, common English names are used throughout the text (where known) together with scientific names.*

The surface of a relatively intact raised bog is typically wet, acid, deficient in plant nutrients, and supports specialised plant communities that are low in overall diversity and comprising species adapted to the biologically harsh conditions. The vegetation is open, treeless and bog mosses or *Sphagnum* species dominate the ground layer. Small-scale mosaics of plant communities are characteristic and reflect the complex microtopography of hummocks and hollows on the bog surface (see Section 1.1.1 below). Raised bogs are driest at the margins and wetness generally increases towards the centre of the peat mass where well-developed pool systems are most likely to occur.

Raised bogs may also contain soaks and flushes (wet 'active' or dry 'inactive') due to the increased supply of nutrients over time through concentrated surface flows, or where there are links with regional groundwater or the underlying mineral substratum. Slight mineral enrichment and / or constant through flow of water provide conditions suitable for a range of species that are not typically associated with other areas of raised bog.

When damaged by peat extraction or drainage, the water table in the peat drops and the bog surface becomes relatively dry; pools are rare or absent, cover of bog mosses is greatly reduced and *Calluna vulgaris* increases in abundance. The drop in water table causes the peat to compress under its own weight causing the bog surface to deform. Greater deformation occurs closest to areas where the water table has dropped. This increases the slope of the bog surface causing rain falling on the ground surface to flow off the bog more quickly. The effect is normally greatest around the margins and in a typical situation surface wetness increases towards the centre of the bog. Trees such as *Betula pubescens* and *Pinus sylvestris* frequently invade the drier cut margins, but may also occur in flushed areas.

In Ireland, the Annex I habitat ARB is currently considered to be in unfavourable bad conservation status principally as a result of marginal turf cutting, more recent semi-industrial peat extraction, and associated drainage effects caused by these activities (NPWS 2008; 2013). The lowering of regional groundwater levels is also known to have had an effect on some sites. Fires associated with turf cutting, dumping, or agricultural activities may also adversely affect the condition of the habitat.

### **1.1.1 Raised Bogs Microtopography**

Raised bogs are typically treeless and are characterised by a distinctive vegetation dominated by bog mosses (*Sphagnum*), sedges, and dwarf shrubs, all of which are adapted to waterlogged, acidic and exposed conditions. Bog mosses, which have unique properties, are the principal component of peat, and are largely responsible for the typical surface features of hummocks, hollows, lawns, and pools. The wettest bogs, which have extensive pool systems, have the greatest variety of plant and animal life and support a range of specialist species.

The following terms that describe microtopography are generally accepted in the study of mire ecology (Gore 1983). A schematic diagram showing the typical microtopographical divisions is presented in Figure 1.

#### **Pools**

Depressions in the bog surface where the water table remains above the surface level all year around or below surface level for only a very short period of time. They are characterised by the presence of aquatic plant species such as *Sphagnum cuspidatum*, *S. denticulatum*, and *Cladopodiella fluitans*. In more degraded scenarios or where high seasonal water fluctuation occurs, the pools contain open water and/or algae. Tear pools are found on bogs where internal tensions, due to mass movement of peat, has taken place within the high bog and

has caused the development of elongated pools. These are frequently found on western bogs and may be natural or anthropogenic in origin.

### Hollows

These are shallow depressions (less than 5cm deep) on the bog surface where surface water collects, or where the water table reaches or lies just above ground level, depending on seasonal conditions. They are often filled with *Sphagnum* species such as *S. papillosum* and *S. cuspidatum*. They take many forms but are often eye shaped. Marginal hollows tend to be elongated as they are focused points for surface water run-off. They are often dominated by *Narthecium ossifragum*.

### Lawns


These are shallow hollows or flat areas where one species dominates to form a lawn. This is frequently a *Sphagnum* species, such as *Sphagnum magellanicum*, or *S. papillosum* which can completely fill in a hollow to form a small lawn.

### Flats

These are more or less flat areas which are intermediate between hollow and hummock communities. They tend to be drier than the above situations.

### Hummocks

These are mounds on the bog surface which can range from a few centimetres to more than one metre in height. They are usually composed mainly of *Sphagnum* species, such as *Sphagnum magellanicum*, *S. capillifolium*, *S. austinii* and *S. fuscum* but other bryophyte species such as *Hypnum jutlandicum* and *Leucobryum glaucum* are also important, especially as the hummock grows taller and becomes drier. *Calluna vulgaris* is another important element, as it flourishes where the water table is not at surface level (Kelly & Schouten 2002).



**Figure 1** Raised bog microtopographical divisions on the high bog surface (reproduced from Kelly & Schouten 2002).

### 1.1.2 Typical Flora of Irish Raised Bogs

Raised bogs are characterised by a distinctive vegetation dominated by a variety of mosses (e.g. *Sphagnum* spp., *Hypnum* spp., *Racomitrium* spp.), sedges and grass-like species (e.g. *Eriophorum* spp., *Rhynchospora* spp., *Narthecium ossifragum*, *Molinia caerulea* and *Carex* spp.), and dwarf shrubs (e.g. *Calluna vulgaris*, *Erica tetralix*, *Vaccinium* spp. and *Empetrum nigrum*). In addition to these groups, a number of other species characterise raised bogs including carnivorous plants (e.g. *Drosera* spp., *Utricularia* spp.), lichens of both the bog surface and epiphytes on the stems of dwarf shrubs and the occasional trees on bogs (e.g. *Cladonia* spp., *Usnea* spp.). Herbaceous plants are not a significant element on raised bogs and include a few commonly occurring species such as *Menyanthes trifoliata*, *Pedicularis sylvatica*, and *Potentilla erecta* (Cross 1990).

Drier areas and hummocks usually support *Calluna vulgaris*, *Eriophorum vaginatum*, *Trichophorum germanicum*, *Erica tetralix*, lichens (*Cladonia* spp.), bog mosses (*Sphagnum capillifolium*, *S. austinii*, *S. fuscum*, *S. papillosum*), and other mosses (*Dicranum scoparium*, *Leucobryum glaucum*). Wet hollow areas and pools are characterised by *Eriophorum angustifolium*, *Rhynchospora alba*, *Narthecium ossifragum*, *Drosera* spp., *Menyanthes trifoliata*, bladderworts (*Utricularia* spp.), and bog mosses (*Sphagnum cuspidatum*, *S. denticulatum* and *S. magellanicum*).

A list of flora species that are regarded as being typical of ARB habitat in Ireland is presented in Table 1. A number of these typical species would have a restricted distribution and do not occur throughout the range of the habitat in Ireland (see above), therefore only a subset of these species would be expected to be present on any individual bog.

**Table 1** Flora species typically associated with active raised bog in Ireland (after NPWS 2013). *Species list is based on vegetation communities defined by Kelly (1993) and Kelly & Schouten (2002).*

Common name	Scientific Name
Bog rosemary	<i>Andromeda polifolia</i>
Bog bead moss	<i>Aulacomnium palustre</i>
Bristly Swan-neck moss*	<i>Campylopus atrovirens*</i>
Lichen	<i>Cladonia ciliata</i>
Lichen	<i>Cladonia portentosa</i>
Long leaved sundew	<i>Drosera anglica</i>
Intermediate leaved sundew*	<i>Drosera intermedia*</i>
Round leaved sundew	<i>Drosera rotundifolia</i>
Common cotton grass	<i>Eriophorum angustifolium</i>
Hare's tail cotton grass	<i>Eriophorum vaginatum</i>
Large white moss	<i>Leucobryum glaucum</i>
Bogbean	<i>Menyanthes trifoliata</i>
Bog asphodel	<i>Narthecium ossifragum</i>
Purple spoonwort*	<i>Pleurozia purpurea*</i>
Woolly fringe moss*	<i>Racomitrium lanuginosum*</i>
White beak-sedge	<i>Rhynchospora alba</i>
Austin's bog moss	<i>Sphagnum austinii</i>
Red bog moss	<i>Sphagnum capillifolium</i>
Feathery bog moss	<i>Sphagnum cuspidatum</i>
Cow-horn bog moss*	<i>Sphagnum denticulatum*</i>
Rusty bog moss	<i>Sphagnum fuscum</i>
Magellanic bog moss	<i>Sphagnum magellanicum</i>
Papillose bog moss	<i>Sphagnum papillosum</i>
Golden bog moss*	<i>Sphagnum pulchrum*</i>
Lustrous bog moss	<i>Sphagnum subnitens</i>
Bladderwort	<i>Utricularia minor</i>
Cranberry	<i>Vaccinium oxycoccos</i>

Notes: \* Species more typical of western raised bog sites.

### 1.1.3 Typical Fauna of Irish Raised Bogs

Raised bogs are extremely nutrient poor ecosystems. Acidic, waterlogged and exposed conditions make them an unattractive habitat for animal life. As a consequence they are relatively poor both in terms of species diversity and population densities. Many species are opportunists, vagrant or temporary rather than specialists, but nonetheless may have an important impact on the ecosystem through nutrient imports and exports or other interactions (Cross 1990). A list of fauna species that would be typically associated with raised bog habitat in Ireland is presented in Table 2. These species are not confined to ARB and most, if not all, will use other areas of the bog and surrounding habitats.

Raised bog is unsuitable habitat for many vertebrates due to the lack of available foraging and suitable breeding places. The Irish hare is the only mammal commonly occurring. The common frog is the most common vertebrate predator.

Although 18 species of birds have been reported breeding on raised bogs (Wilson 1990) many of these species utilise the bog as a nesting habitat only. They are dependent on other neighbouring habitats such as open water bodies, callows and wet grassland particularly for

feeding. Just a few species of bird, including meadow pipit (*Anthus pratensis*), skylark (*Alauda arvensis*) and curlew (*Numenius arquata*) complete their full breeding cycle on the bog and the first two species are the commonest species occurring (Bracken *et al.* 2008). Red grouse (*Lagopus lagopus*) must also be included as a typical bog species, occurring year round as a resident. Red grouse and curlew have declined significantly on across raised bogs in recent times. BirdWatch Ireland have published an Action Plan for Raised Bog Birds in Ireland which lists 13 species of conservation concern that are associated with Raised Bogs (O'Connell 2011). A recent review of birds of conservation concern in Ireland has since added meadow pipit (*Anthus pratensis*) to the red list of birds of conservation concern (Colhoun & Cummins 2013).

Our knowledge of the invertebrate assemblages associated with Irish raised bogs remains incomplete (particularly micro-invertebrate species) with few studies undertaken (Reynolds 1984a; Reynolds 1984b; Reynolds 1985; De Leeuw 1986; O Connor *et al.* 2001; Crushell *et al.* 2008; Hannigan & Kelly-Quinn 2011; Wisdom & Bolger 2011; Nolan 2013). Van Duinen (2013) highlights the importance of structural diversity at various spatial scales (e.g. micro-scale of hummock hollow topography to macro-scale which would include the landscape setting of the bog, see Schouten (2002)) as a prerequisite for hosting the full species diversity of raised bog landscapes.

A recent study of Lepidoptera associated with raised bogs identified two species that appear to be characteristic of higher quality raised bog habitat, namely bordered grey (*Selidosema brunnearia* (Villers, 1789)) and light knot grass (*Acronicta menyanthidis* (Esper, 1789)) (Ciara Flynn pers. comm.).

Recent research on spiders has revealed that a number of species are known to occur in Ireland only on raised bog habitats, all of which are considered local/uncommon or rare across Europe (Myles Nolan pers. comm.). Five of these species that can be considered useful indicators of ARB include: *Glyphesis cottonae* (La Touche 1945), *Walckenaeria alticeps* (Denis 1952), *Satlatlas britteni* (Jackson 1913), *Pirata piscatorius* (Clerck 1757), and *Minicia marginella* (Wider 1834) (Myles Nolan pers. comm.).

The information currently available on other invertebrate groups of peatland systems in Ireland is not sufficient to allow a determination of many species that are typically associated with or may be characteristic of higher quality ARB. A selection of invertebrate species and species groups that are known to be typically associated with raised bogs are included in Table 2.

**Table 2** Fauna species typically associated with raised bog ecosystems in Ireland (after O’Connell 1987; Cross 1990; Renou-Wilson *et al.* 2011; Bracken & Smiddy 2012).

Common name	Scientific name
<b>Mammal species</b>	
Irish hare	<i>Lepus timidus hibernicus</i>
Otter	<i>Lutra lutra</i>
Pygmy shrew	<i>Sorex minutes</i>
Fox	<i>Vulpes vulpes</i>
<b>Bird species</b>	
Skylark	<i>Alauda arvensis</i>
Mallard	<i>Anas platyrhynchos</i>
Greenland white-fronted goose	<i>Anser albifrons flavirostris</i>
Meadow pipit	<i>Anthus pratensis</i>
Hen harrier	<i>Circus cyaneus</i>
Cuckoo	<i>Cuculus canorus</i>
Merlin	<i>Falco columbarius</i>
Kestrel	<i>Falco tinnunculus</i>
Snipe	<i>Gallinago gallinago</i>
Red grouse	<i>Lagopus lagopus</i>
Curlew	<i>Numenius arquata</i>
Golden plover	<i>Pluvialis apricaria</i>
Lapwing	<i>Vanellus vanellus</i>
<b>Reptiles and amphibians</b>	
Common lizard	<i>Lacerta vivipara</i>
Common frog	<i>Rana temporaria</i>
<b>Typical invertebrates</b>	
Black slug	<i>Arion ater</i>
Large heath butterfly	<i>Coenonympha tullia</i>
Marsh fritillary butterfly	<i>Euphydryas aurinia</i>
Bog-pool spider	<i>Dolomedes fimbriatus</i>
Water striders	<i>Gerris</i> and <i>Velia</i> species
Oak eggar moth	<i>Lasiocampa quercus</i>
Four-spotted chaser dragonfly	<i>Libellula quadrimaculata</i>
Fox moth	<i>Macrothylacia rubi</i>
Ant	<i>Myrmica ruginodis</i>
Emperor moth	<i>Saturnia pavonia</i>
Great green bog grasshopper	<i>Stethophyma grossa</i>
Other species groups that are well represented on raised bogs include:	Araneae (spiders and mites)
	Ceratopogonidae (biting-midges)
	Chironomids (non-biting midges)
	Coleoptera (beetles)
	Collembola (springtails)
	Diptera (true flies)
	Dytiscidae (water beetles)
	Hemiptera (true bugs)
	Hymenoptera (bees, wasps, ants and sawflies)
	Lepidoptera (butterflies and moths)
	Odonta (dragonflies and damselflies)
	Orthoptera (grasshoppers)
	Syrphidae (hoverflies)
Tipulidae (crane-flies)	
Tabanidae (horseflies)	

## 1.2 Habitats Directive Raised Bog Habitats in Ireland

Four habitat types listed on Annex I of the EU Habitats Directive are typically associated with raised bogs in Ireland, two of which are priority habitats (\*):

- 7110 Active raised bogs (ARB)\*
- 7120 Degraded raised bogs still capable of natural regeneration (DRB)
- 7150 Depressions on peat substrates of the *Rhynchosporion*
- 91D0 Bog woodland\*

The interpretation manual of EU habitats gives the following description for 'active raised bogs': "*Acid bogs, ombrotrophic, poor in mineral nutrients, sustained mainly by rainwater, with a water level generally higher than the surrounding water table, with perennial vegetation dominated by colourful Sphagna hummocks allowing for the growth of the bog (Erico-Sphagnetalia magellanici, Scheuchzerietalia palustris p., Utricularietalia intermedio-minoris p., Caricetalia fuscae p.). The term "active" must be taken to mean still supporting a significant area of vegetation that is normally peat forming, but bogs where active peat formation is temporarily at a standstill, such as after a fire or during a natural climatic cycle e.g., a period of drought, are also included.*" (CEC 2007).

DRB should be, according to the interpretation manual capable of regeneration to 'Active Raised Bog' in 30 years if appropriate measures are put in place (i.e. no major impacting activities are present and any necessary restoration works are implemented).

In Ireland, the identification of ARB is made at ecotope level based on the vegetation classification developed by Kelly (1993) and Kelly & Schouten (2002).

Raised bog vegetation communities are grouped into a series of community complexes and these complexes are then amalgamated into a series of ecotopes characterised by different physical characteristics using the approach outlined by Kelly & Schouten (2002).

The main ecotopes that community complexes are grouped into include:

- Central ecotope
- Sub-central ecotope
- Active flushes and soaks
- Sub-marginal ecotope
- Marginal ecotope
- Inactive flushes
- Face-bank ecotope

Actively accumulating peat conditions occur within the sub-central and central ecotopes, which are the wettest on the bog and an indication of good quality ARB. Active flushes and soaks are also dominated by *Sphagnum* mosses and typically have wet conditions. These features are associated with ARB and contribute to the overall diversity of the habitat.

The adjacent surrounding marginal, sub-marginal, and face-bank bog areas typically have a supporting function for the central and sub-central communities but are not peat accumulating. These drier ecotopes may or may not correspond to the Annex I habitat DRB, as it depends on whether they are capable of regeneration to ARB. Other drier ecotopes recorded on the high bog that do not correspond to ARB include 'inactive flushes' which typically have a low *Sphagnum* cover.

The Annex I habitat Rhynchosporion depressions (7150) typically occurs along pool edges and on flats underlain by deep, wet and quaking peat. Typical plant species include *Rhynchospora alba*, *Drosera anglica*, *Narthecium ossifragum*, *Sphagnum cuspidatum*, *S. denticulatum*, *S. magellanicum*, *S. papillosum*, *Menyanthes trifoliata*, and *Eriophorum angustifolium*.

The priority Annex I habitat bog woodland is also actively peat-forming and overlaps with the ARB habitat. Such woodlands are usually dominated by *Betula pubescens* with a characteristic ground cover dominated by *Sphagnum* moss species, which often form deep carpets, and other mosses including species of *Polytrichum*. Woodland areas are occasionally found on raised bogs that have an absence of the characteristic moss layer and are not regarded as peat forming. Such areas do not correspond to the Annex I habitat.

### **1.2.1 Restoration of Active Raised Bog in Ireland**

As already mentioned in the section 1.1, ARB is currently considered to be in unfavourable bad conservation status in Ireland. In addition, according to its definition, DRB should be capable of regeneration to ARB in a 30-year timescale. Thus, it follows that restoration measures are required in order to halt further losses and increase the area of ARB as well as to improve the condition of existing areas of the Annex I habitat.

Most of the restoration works undertaken so far in Ireland have concentrated on the high bog (e.g. Clara Bog, Mongan Bog, Sharavogue Bog and Raheenmore Bog) to prevent further losses as well as to restore areas to ARB. Nevertheless, some restoration works have also been undertaken on cutover areas such as at Ballykenny and Fisherstown Bogs and Killyconny Bog. Such work aims to do one or more of the following (depending on the bog in question): restore ARB on the high bog; reduce further ARB and DRB loss on the high bog; restore peat forming habitats (such as ARB, bog woodland, poor fen) on the cutover.

Works undertaken by the NPWS have indicated that there are significant differences, both ecological and economic, when comparing the effectiveness of works carried out on the cutover with those carried out on the high bog. Positive and significant results (i.e. expansion or development of ARB) can be achieved over a relatively short timeframe (10 years) on favourable areas of the high bog by blocking high bog drains. In contrast, a longer time period (30 years+) is required to achieve active peat formation on cutover areas, and even then the results are generally confined to smaller areas; i.e. flat areas ( $\leq 0.3\%$  surface slope) or enclosed depressions that have sufficient water flow (minimum catchment 0.5ha) to maintain wet conditions throughout the year. A longer time period (minimum 50-100 years) is likely to be required for high quality ARB habitat (vegetation structure and species diversity) to develop on such cutover areas. In addition, costs of restoration measures on cutover areas are typically significantly higher than those on high bog areas.

## **1.3 Garriskil Bog SAC**

The SAC includes the raised bog, known as Garriskil Bog. The SAC has been selected for the following three Annex I habitats:

- [7110] Active raised bogs\*
- [7120] Degraded raised bogs still capable of natural regeneration
- [7150] Depressions on peat substrates of the *Rhynchosporion*

This raised bog lies 3km west of Lough Derravaragh and 3km east of Rathowen in County Westmeath. It is bounded to the south-east and south-west by the rivers Inny and Riffey. It is considered an excellent example of a midland raised bog of the ridge river morphological type (Kelly *et al.* 1995).

The bog contains a large, wet central core of ARB which comprises a relatively large area of

the remaining high bog. There is a well developed system of pools and hummocks within this area.

Peat cutting has not been reported from the site since the 1990s (Kelly *et al.* 1995; Fernandez *et al.* 2005, 2014a, b). Nevertheless, old face banks and cutover drainage associated with cutting along the entire southern half of the site are likely to continue to be causing negative impacts on the high bog habitats. Past drainage of the bog associated with peat cutting has unfavourably impacted on the bog (by causing a lowering of the water table), although many of these drains have now been infilled and blocked. Arterial drainage of the River Inny (undertaken by the OPW in 1996 and initially during the 1950s) is likely to have impacted on the site and may continue to pose a threat to the hydrological integrity of the site.

Burning in the past has caused damage, though there has been no severe fire events reported in recent years with no burning reported since 1997 (Fernandez *et al.* 2014a, b).

Notably, a large proportion of the uncut high bog comprises very wet ARB, which is unusual for raised bogs in the eastern half of the country. Because of its relatively good condition the site is considered to be one of the best remaining examples of a raised bog ecosystem in the eastern half of the country.

### 1.3.1 Flora of Garriskil Bog

The largest ARB area in the central part of the high bog occurs around the well-developed pool complexes found in this area. This area has a bog surface that is wet and quaking in places. These pools are mainly inter-connecting and are more typical of western raised bogs with islands containing *Racomitrium lanuginosum* (though not frequent), *Sphagnum denticulatum* in the pools and *Campylopus atrovirens* occurring around the pool edges. The pools contain *Sphagnum cuspidatum*, *S. denticulatum*, and *S. magellanicum* with frequent *S. magellanicum* and *S. papillosum* around the pool edges. The pools also contain *Menyanthes trifoliata*, *Rhynchospora alba*, and *Eriophorum angustifolium* in places. The inter-pool areas are generally dominated by *Narthecium ossifragum* in flats and hollows with *Calluna vulgaris* occurring on hummocks. There are occasional hummocks of *Sphagnum fuscum* and more frequent *S. papillosum* and *S. capillifolium*. *Sphagnum austinii* is relatively frequent in some parts of the ARB with older typical *S. austinii* hummocks and low more recently developed hummocks both present, especially in some of the best quality ARB areas. The *Cladonia* spp. cover is not as extensive as on other sites suggesting a more recent burning history (Fernandez *et al.* 2005).

A second central ecotope complex occurs in a small part of the south-west corner of this area, which has smaller pools infilled with *Sphagnum*, which are more typical of midlands raised bogs. The sub-central ecotope areas usually contain poorer quality pools with less frequent *Sphagnum* and more frequent algae. Around the edges of the active bog area the pools are less frequent, not interconnecting and of lower quality.

The ARB area located in the north-west of the high bog close to the high bog margin includes active flush, central, and sub-central ecotope. The flush and central ecotope area contains a very wet and quaking extensive *Sphagnum* cover dominated by lawns of *Sphagnum magellanicum*, *S. cuspidatum* with *Rhynchospora alba* and occasional small pools. The flush occurs in a slight depression and some parts contain *Carex rostrata* and *Aulacomnium palustre*, which defines the flushed area, otherwise there is no distinct boundary between the flush and central ecotope.

A small ARB area occurs on the south side of the bog with frequent *Eriophorum vaginatum* and *Eriophorum angustifolium* occurring, along with *Sphagnum* cover, in a slight depression. Drain blocking has been relatively successful here with re-wetting occurring along the drains

and *Sphagnum* cover developing, dominated by lawns and low hummocks of *S. capillifolium* and *S. papillosum*.

The inactive areas on Garriskil Bog surface are generally firm or firm to soft around the margins. These complexes occur on flat and gently sloped parts of the bog with steeper slopes along the margins. The sub-marginal and marginal ecotope areas generally occur around the margins of the high bog. These areas generally have few well-developed pools. However, a large section of sub-marginal ecotope occurs which has frequent pools of poor quality and the interpool flats dominated by *Narthecium ossifragum* and a low *Sphagnum* cover. Generally the sub-marginal areas are dominated by hummocks of *Calluna vulgaris* and frequent *Eriophorum vaginatum* and hollows and flats with frequent *Narthecium ossifragum*, *Eriophorum vaginatum*, and *E. angustifolium*. In some areas the hummocks become less frequent. The marginal vegetation is similar to the sub-marginal vegetation although the *Sphagnum* cover is poorer and this ecotope generally occurs on steeper slopes. *Trichophorum germanicum* occurs more frequently in some parts of the margin and there are more frequent *Narthecium ossifragum* dominated areas. The area within the drain complex contains well-developed ridges of *Calluna vulgaris* on old spoil dug out of the drains (Fernandez *et al.* 2005).

There are several small flushes scattered around the high bog characterised by small groups of *Pinus sp.*, *Betula pubescens*, and *Salix sp.* trees and saplings. Often the areas with *Pinus sp.* are associated with small flushes dominated by species such as *Molinia caerulea*, *Juncus effusus*, *Rubus fruticosus* agg., and *Calluna vulgaris*.

A swallowhole occurs on the north-western margin of the high bog. This area is quite poached from cattle encroaching onto the bog and using the swallow hole as a drinking place.

*Rhynchospora alba*-dominated depressions occur relatively frequently on the high bog and occur within both the active and degraded raised bog in mosaic with these habitats. Relatively good quality examples of this habitat occur within the central ecotope areas.

Along the north-east margin of the high bog a narrow band of fen-grassland occurs.

Areas of old cutover bog provides an additional habitat where *Molinia caerulea* and *Calluna vulgaris* dominate, along with *Eriophorum sp.*, while in some parts *Betula pubescens* woodland is developing.

### 1.3.2 Fauna of Garriskil Bog

The following is based on information contained in NPWS (2004). A flock of Greenland white-fronted geese (*Anser albifrons flavirostris*) (listed in Annex I of the Birds Directive) ranges over the midland lakes, including those close to the site (e.g. Lough Derravaragh). In the winter of 1993/94 this internationally important flock consisted of 346 birds (Fox *et al.* 1994). Garriskil Bog formerly provided an important feeding and/or refuge area for the geese, but Fox *et al.* (1994) report that the birds now feed mostly on intensively managed grassland and take refuge on the lakes.

Merlin (*Falco columbarius*) (Annex I, Birds Directive) is reported to hunt within the site and may breed in a nearby conifer plantation

Other birds using the site include hunting barn owl (*Tyto alba*) and small numbers of breeding redshank (*Tringa totanus*), curlew (*Numenius arquata*), lapwing (*Vanellus vanellus*) and snipe (*Gallinago gallinago*), with meadow pipit (*Anthus pratensis*) also occurring during the summer months.

## 2 Conservation objectives

A site-specific conservation objective aims to define the favourable conservation condition of a habitat or species at site level. The maintenance of habitats and species within sites at favourable condition will contribute to the maintenance of favourable conservation status of those habitats and species at a national level.

Conservation objectives for habitats are defined using attributes and targets that are based on parameters as set out in the Habitats Directive for defining favourable status, namely area, range, and structure and functions. Attributes and targets may change or become more refined as further information becomes available.

National Conservation Objectives for raised bog SACs have recently been published in the Draft National Raised Bog SAC Management Plan (DAHG 2014). The various attributes and the justification of appropriate targets used to define favourable conservation condition for ARB relevant to Garriskil Bog SAC are discussed in the following sections.

### 2.1 Area

NPWS has commissioned a number of raised bog surveys between 1993 and the present - Kelly *et al.* (1995); Derwin & MacGowan (2000); Fernandez *et al.* (2005); Fernandez *et al.* (2006); Fernandez *et al.* (2014). Mapping from these surveys has been used to derive the area of ARB for each bog as shown in Table 3. More recent surveys have been able to employ more precise and detailed mapping techniques and more standardised ecotope descriptions. NPWS undertook a review of data from earlier surveys in 2014 taking into account these improved techniques with the aim of providing more accurate figures for ARB. This in some cases has resulted in a change in ARB area for these earlier time periods (NPWS, unpublished data).

The national SAC target for the attribute 'habitat area' has been set at 2,590ha (DAHG 2014). This target is based on the estimated area of ARB (1,940ha) and DRB (650ha) present within the SAC network in 1994 (when the Habitats Directive came into effect).

The area of ARB at Garriskil Bog in 1994 is estimated to have been 71.2ha, while the area of DRB is estimated to have been 11.2ha at that time (see Table 3). Using the same approach that has been adopted in setting the national SAC target, the site-specific target for Garriskil Bog would equate to 82.4ha (sum of ARB and DRB in 1994). However, in setting the site-specific target the current hydro-ecological conditions on the bog (including cutover) have been considered in order to ensure that the target being set is based on a realistic appraisal of what is achievable as set out below.

The most recent monitoring surveys of the bog estimated the area of ARB to be 50.9ha (Fernandez *et al.* 2014a, b). This represents a decline of 20.3ha (28.5%) during the period 1994 – 2012, as a result of drying out of areas in the north-east caused by peat cutting activities, and cracking and slumping of the bog in the east (see Table 3) (Fernandez *et al.* 2014).

The current extent of DRB as estimated using a recently developed hydrological modelling technique, based largely on Light Detection And Ranging (LiDAR)<sup>2</sup> data, is 31.6ha (see DAHG 2014 for further details of the technique). This represents the area of the high bog, which does not currently contain ARB but has topographical conditions deemed suitable to support ARB (see Map 1 which shows the total area of current and modelled potential ARB). Following

---

<sup>2</sup> LiDAR is a remote sensing technology that measures vertical surface elevation by illuminating a target with a laser and analysing the reflected light. This provides much more detailed topographical maps than can be collected by traditional surveying techniques.

a review of the current drainage on the high bog, it was considered that this entire area could potentially be restored by the continued positive effects of past drain blocking along with additional blocking of remaining drains on the high bog (see DAHG 2014).

Based on the current assessment of the bog above, it is therefore concluded that the maximum achievable target for ARB on the high bog is 82.5ha, which is 0.1ha less than the estimated area at the time of designation. However, it is important to note that this assumes no further decline of ARB due to losses of high bog caused by turf cutting and drainage activities associated with same (Fernandez *et al.* 2014ab). Similarly, should the bog be significantly dependent on regional groundwater levels then any deepening of drains in the cutover could further impact the potential restoration of ARB on the high bog.

**Table 3** Area of ARB and DRB recorded on the high bog at Garriskil Bog in 1994, 2004, and 2012 (Source: Fernandez *et al.* 2014a, b).

1994		2004		2011	
ARB (ha)	DRB (ha)	ARB (ha)	DRB (ha)	ARB (ha)	DRB (ha)
71.2	11.2	45.1	Unknown	50.9	31.6

A recent eco-hydrological assessment of the cutover surrounding the high bog undertaken as part of the restoration planning process estimates that, by implementing appropriate management, an additional 2.4ha of ARB could be restored in this area. The long term achievable target for ARB on Garriskil Bog is therefore set at 84.9ha which exceeds the estimated area of ARB and DRB in 1994 by 2.5ha.

In conclusion, the site-specific target for the attribute habitat area is: **Restore area of active raised bog to 84.9ha, subject to natural processes.**

## 2.2 Range

At a national scale, range represents the geographic range that encompasses all significant ecological variations of the ARB habitat. The national SAC target for the attribute ‘range’ has been set as ‘not less than current range subject to natural processes’.

However, range, in the form of habitat distribution, may also be important at the site level, particularly within larger SACs, including those containing a number of individual bogs (i.e. complexes). The attribute therefore under the parameter of range is ‘Habitat distribution’. At the local level, it is important to conserve the variability and distribution of ARB across a raised bog SAC. This will help to ensure the diversity of the habitat is maintained while lessening the impact of localised damaging activities such as fire.

The conservation of ARB within Garriskil Bog as set out in Section 2.1 above will contribute to safeguarding the national range of the habitat.

The ARB habitat at Garriskil Bog includes central and sub-central ecotope, as well as active flush. A map showing the most recent distribution of ecotopes throughout Garriskil Bog is presented in Map 2. Drain blocking undertaken in Garriskil Bog during the 1990s is having a significant positive effect on the distribution of ARB (Fernandez *et al.* 2014a, b).

The site-specific target for the attribute habitat distribution is: **Restore the distribution and variability of active raised bog across the SAC.**

## 2.3 Structure and functions

Structure and functions relates to the physical components of a habitat (“structure”) and the ecological processes that drive it (“functions”). For ARB these include attributes such as the hydrological regime, water quality, habitat quality, species occurrence, elements of local distinctiveness, marginal habitats, negative physical indicators, and negative species occurrence. As several of these attributes are inter-connected, they are all included in order to better define habitat quality in a meaningful way. In some cases, attribute targets are not quantified; however, as more detailed information becomes available (for example through further research), more measurable site-specific targets may be developed. Structure and functions attributes are expanded on in the sections below.

### 2.3.1 High bog area

On individual raised bogs adequate high bog is required to support the development and maintenance of ARB. Raised bog habitat that is classified as neither ARB nor DRB is still important particularly as a supporting habitat for those listed in Annex I of the Habitats Directive. It is an essential part of the hydrological unit which supports the ARB and DRB habitats. High bog is of value in its own right as a refuge for species characteristic of drier bog conditions as well as for providing a transitional zone between the Annex I habitats of the high bog and surrounding areas. Additional values for the maintenance of high bog include the preservation of its record of past environmental conditions and carbon storage. The area of high bog in the entire SAC network in 1994 was 10,740ha. The corresponding area in 2012 is 10,515ha – indicating that there has been a 225ha loss of high bog since 1994.

The national target for the attribute ‘high bog’ habitat is to ensure no decline in extent of high bog to support the development and maintenance of ARB.

The area of high bog within Garriskil Bog SAC in 1994 was mapped as 170.3ha, and the corresponding area in 2012 is also 170.3ha (based on interpretation of LiDAR and aerial photography flown in 2012), indicating there has been no loss of high bog (DAHG 2014). The extent of high bog within the SAC in 2012 is illustrated on Map 1.

The site-specific target for the attribute high bog is: **No decline in extent of high bog necessary to support the development and maintenance of active raised bog.**

### 2.3.2 Hydrological regime: water levels

Hydrological processes are key drivers of raised bog ecology. The different raised bog communities, assemblages and species are affected by various hydrological attributes. For ARB, mean water levels need to be near or above the surface of bog lawns for most of the year. Seasonal fluctuations should not exceed 20cm, and water level should be within 10cm of the surface, except for very short periods of time (Kelly & Schouten 2002). Gentle slopes that limit intermittent lateral losses of water (through surface runoff) and encourage sustained waterlogging are the most favourable to achieve these conditions. These conditions may be maintained on steeper slopes in areas of focused flow (flushes).

The traditional view of water flowing across the bog laterally has been recently refined to also consider that water flows vertically through peat into the underlying substrate. Water loss, by this route, depends on the permeability of the material through which the water must flow and the difference in head (water level elevation) in the bog and underlying mineral substrate; larger differences encountered in higher permeability materials will result in greater losses. Although the proportion of water lost in this manner may be small, the sustained loss during prolonged dry periods may be sufficient to impact bog ecotopes. Drains extending into the mineral substrate in marginal areas surrounding the bog can lead to an

increased gradient between the head in the peat and the head in the underlying substrate resulting in increased vertical water losses from the bog.

The most recent description of drainage at Garriskil Bog is presented in Fernandez *et al.* (2014a, b) who noted few unblocked drains remaining on the high bog surface (0.4km), of which 0.2km are considered to be reduced functional and 0.2km non-functional. Most of the drains on the high bog were blocked in 1998 and 12.0km of blocked drains are reported to be reduced functional. Positive results from drain blocking activities have been recorded during the last two monitoring surveys, with on-going infilling of drains and expansion of active raised bog in these areas. Cutover drains occur around most of the high bog, although drainage associated with agriculture is more common along the north/north-east.

Garriskil Bog is bounded by three permanently flowing water courses; the River Inny along the south-eastern boundary, River Riffey along the southern boundary and a small tributary of the River Riffey along the west of the site. Kelly *et al.* (1995) noted that the drain/stream to the west of the site had recently been deepened and re-dredged, cutting directly into a stony till mineral substrate. Kelly *et al.* (1995) also noted that the River Inny and River Riffey had been dredged and suggested that the subsidence, suspected in the southern part of the high bog, is likely to be linked to this activity. The southern side of the Inny River was dredged by the Office of Public Works (OPW) in 1996, under their statutory obligation to maintain the channel. The river had previously been dredged in the 1950s. It was not confirmed whether this activity has taken place again in the 2004/5-2011 reporting period. Other issues which may be causing impacts to water levels include invasive species; *Rhododendron* in particular has been recorded as spreading on the bog and may have a comparable effect to trees in maintaining water levels at depths that prevent *Sphagnum* growth.

Much of the knowledge regarding the hydrological requirements of raised bog communities in Ireland stems from the extensive ecological and hydrological work undertaken on Clara Bog since the early 1990s. The only available hydrological study for Garriskil Bog is the work carried out by Kelly *et al.* (1995)

The hydrochemistry survey carried out as part of this study indicated that generally cutover drains had limited groundwater influence since electrical conductivities were typically in the region of 100 $\mu$ s/cm. However, very few sample points were noted within this study, therefore it is not possible to definitely state that there was not upwelling of mineralised groundwater in the cutover. As indicated previously, river channels surrounding the bog were dredged and deepened which is likely to have impacted upon regional groundwater heads. This may have impacted upon pore water pressure within the peat and resulted in subsidence as suggested by Kelly *et al.* (1995). The extent of any impact would depend on the permeability of the substrate. Geological mapping indicates that the bog is underlain by dark limestone and shale, while a SW/NE trending fault lies along the south-eastern boundary of the site. This unit is classified as a locally important aquifer, which is moderately productive only in local zones. The dominant subsoil type in the surrounding areas is sandstone till (Lower Palaeozoic/Devonian) along with chert till north-west of the site. Some esker deposits occur towards the western side of the bog, while limestone till is dominant east of the River Inny. This would suggest that there may be some relatively permeable material underlying some parts of the bog. Therefore there is a risk that further dredging or deepening of drains could lower water levels within the peat.

The site-specific target for the attribute hydrological regime – water levels is: **Restore appropriate water levels throughout the site.**

### 2.3.3 Hydrological regime: flow patterns

As outlined above, ARB depends on water levels being near or above the surface of bog lawns for most of the year. Long and gentle slopes are the most favourable to achieve these conditions. Changes to flow directions due to subsidence of bogs can radically change water regimes and cause drying out of high quality ARB areas and soak systems.

A map illustrating the slopes and drainage patterns on Garriskil Bog based on a digital elevation model generated from LiDAR imagery flown in 2012 is presented in Map 3.

This map shows that Garriskil Bog has a prominent domed topography, as would be typically associated with raised bogs, with a flat central dome that slopes more steeply towards the margins. In general, water flow patterns are radial from the centre of the bog towards the margins; however, there is also evidence that the topography in some areas may have been affected by peat cutting, marginal drainage, and dredging of the adjacent rivers. Steep slopes extending into the central dome and areas of focused flow, such as along the southern margin of the bog, suggest this bog has been impacted by hydrogeological impacts. Further impacts to regional groundwater heads as a result of dredging or drainage in the surrounding areas have the potential to result in further subsidence on Garriskil Bog which could impact upon flow patterns.

The site-specific target for the attribute hydrological regime – flow patterns is: **Restore, where possible, appropriate high bog topography, flow directions and slopes.**

### 2.3.4 Transitional areas between high bog and surrounding mineral soils (includes cutover areas)

Transitional zones between raised bogs and surrounding mineral soils are typically cutover bog and drained lagg zones. The maintenance / restoration of these areas will help to maintain hydrological integrity of ARB and DRB and support a diversity of other wetland habitats (e.g. wet woodland, swamp and fen) as well as species that they sustain. In some cases, these areas may assist in reducing further losses of ARB / DRB on the high bog and in time could develop into active peat forming habitats (including ARB - see Section 2.1 above). These transitional zones, once restored, can provide ecosystem services through flood attenuation and water purification to downstream areas and potentially increase the carbon storage / sink function of the bog. The estimated extent of such transitional areas within the SAC network is 3,000ha (DAHG 2014). The national target for these transitional areas is to maintain / restore semi-natural habitats with high water levels around as much of the bog margins as necessary.

The transitional areas at Garriskil Bog include a range of different habitat types (e.g. improved grassland, wet grassland, cutover bog, scrub, rivers, and railway embankments). The total area of cutover bog within the Garriskil Bog SAC is estimated to be circa 109ha. The development of habitats within cutover areas depends on a number of factors including prevailing land-use, topography, upwelling regional groundwater, and drainage.

At Garriskil Bog, wet grassland mainly occurs along the Inny River and near the railway and is generally used for grazing cattle. Species present include *Juncus articulatus*, *J. acutiflorus*, *J. effusus*, *Agrostis stolonifera*, sedges (*Carex* spp.), *Ranunculus acris*, and *Succisa pratensis* as described by O'Sullivan & Ryan (1994). Sections of the wet grassland are species-rich and Douglas & Grogan (1986) noted an area that they described as wet fen-grassland. There are also small fragments of marsh, with species such as *Potentilla palustris*, *Valeriana officinalis* and *Mentha* sp.

The Inny River is a medium-flowing, marl-bottomed river with a broad channel approximately

50m wide. In places, the channel is colonised by *Typha latifolia* and *Schoenoplectus lacustris*. Other species present include *Potamogeton natans* and starworts (*Callitriche* spp.). The small, gravel-bottomed stream along the north-western boundary of the site supports species such as *Glyceria fluitans* and *Iris pseudacorus*.

In some areas of the cutover, *Ulex europaeus* and/or *Betula pubescens* dominated scrub is established. In several areas woodland is beginning to form with a canopy reaching 15m. The dominant species is *Betula pubescens*, with a few other species such as *Sorbus aucuparia*, willows (*Salix* spp.) and *Ilex aquifolium* occurring. Grazing has led to a sparse ground flora dominated by grasses and *Rubus fruticosus* agg.

A few reclaimed fields occur in the northern part of the bog, located adjacent to the railway, which are quite intensively managed and fertilised (O'Sullivan & Ryan 1994).

The site-specific target for the attribute transitional areas is: **Restore adequate transitional areas to support / protect active raised bog and the services it provides.**

### **2.3.5 Vegetation quality: central ecotope, active flush, soaks, bog woodland**

A diverse good quality microtopography on raised bogs consists of *Sphagnum* dominated pools, hollows, lawns and hummocks, which support the highest diversity of species including hummock indicators: *Sphagnum fuscum* and *S. austinii*; pool indicators: *S. cuspidatum*, *S. denticulatum*, and indicators of lack of burning events e.g. some lichen species (*Cladonia* spp.) (Cross 1990).

The national target for the attribute vegetation quality has been set as “to maintain / restore sufficient high quality bog vegetation (i.e. central ecotope and / or flushes / soaks). At least 50% of ARB habitat should be central ecotope and / or flush / soaks.” Bog woodland is also regarded as a desirable variant of ARB as it adds species and structural diversity to the habitat and therefore, where relevant, also contributes to the 50% target at a site level.

A summary description of the vegetation of Garriskil Bog is presented in Section 1.3.1 above. The vegetation and habitats of the bog have been described in more detail by Kelly *et al.* (1995), Fernandez *et al.* (2005, 2014a, b).

The extent of the different ecotopes that correspond with ARB based on the most recent surveys is presented in Table 4 and on Map 2. It can be seen that the proportion of ARB that comprises central ecotope and active flush is currently 31.9%. Comparing this to results of surveys undertaken in 2004 indicates that although the total area of ARB increased across the site, the proportion of each of the component ecotopes remained relatively static. The examples of both ecotopes that are present on the bog equate to high quality ARB.

The target for this attribute is 42.5ha of central ecotope, active flush and/or bog woodland (50% of ARB target area (84.9ha)). This requires an increase from the current area of central ecotope, active flush and/or bog woodland from 14.7ha to 42.5ha.

**Table 4** Extent of ecotopes classified as ARB in 2004 and 2011 (modified from Fernandez *et al.* 2014).

Ecotope	2004		2011	
	ha	% of total ARB	ha	% of total ARB
Sub-central ecotope	30.7	68.0	36.2	71.1
Central ecotope	14.0	31.0	14.3	28.1
Soaks / active flush	0.4	0.9	0.4	0.8
<b>Total ARB</b>	<b>45.12</b>		<b>50.9</b>	

The site-specific target for the attribute vegetation quality is: **Restore 42.5ha of central ecotope/active flush/soaks/bog woodland as appropriate.**

### 2.3.6 Vegetation quality: microtopographical features

The characteristic microtopographical features of raised bogs are described in Section 1.1.1 above.

Hummock and hollow microtopography, and pool systems are well developed in the central part of Garriskil Bog. Previous drainage efforts on the north-eastern part of the bog and possibly the effects of arterial drainage have had a negative effect on the surface microtopography (Kelly *et al.* 1995; Fernandez *et al.* 2005, 2014a, b), although the effects of this may be declining following drain blocking works on the bog in 1998.

The site-specific target for the attribute microtopographical features is: **Restore adequate cover of high quality microtopographical features.**

### 2.3.7 Vegetation quality: bog moss (*Sphagnum*) species

Bog mosses, which have unique properties, are the principal component of peat, and are largely responsible for the typical microtopographical features as described in Section 2.3.6 above.

The vegetation of a typical raised bog that is still hydrologically intact is characterised by the dominance of several species of *Sphagna* and dwarf ericoid shrubs. The most abundant species are *Sphagnum capillifolium*, *S. austinii* and *S. papillosum* which form hummocks or low ridges. *Sphagnum fuscum* may also form hummocks (Cross 1990). On the flats *Sphagnum magellanicum*, *S. papillosum*, *S. tenellum*, and *S. subnitens* are the key species. *Sphagnum pulchrum* may also be dominant in flats on western raised bogs. In permanently waterlogged hollows *Sphagnum cuspidatum* and *S. denticulatum* (western bogs) occur. *Sphagnum fallax* is common where there is slight flushing (Cross 1990). The most commonly occurring *Sphagnum* moss species that occur on raised bogs in Ireland are presented in Table 5 along with a summary of their ecology and typical contribution to peat formation.

Kelly *et al.* (1995) and Fernandez *et al.* (2005 & 2014a, b) provide further information on the occurrence of *Sphagnum* species throughout Garriskil Bog.

**Table 5** *Sphagnum* species typically associated with raised bog ecosystems in Ireland. Ecology as described by Laine *et al.* (2009) with minor modifications.

Species	Ecology	Peat forming capacity
<i>Sphagnum austinii</i>	Hummock species	High
<i>Sphagnum capillifolium</i>	Forms small hummocks and carpets	Moderate
<i>Sphagnum cuspidatum</i>	Pool and hollow species	Low
<i>Sphagnum denticulatum</i>	Pool and hollow species	Low
<i>Sphagnum fallax</i>	Occurs in lawns and carpets, shade tolerant. Indicative of some nutrient enrichment (soaks and active flushes)	Low
<i>Sphagnum fuscum</i>	Forms dense low and wide, and occasionally high hummocks	High
<i>Sphagnum magellanicum</i>	Lawn species forming carpets and low hummocks	Moderate
<i>Sphagnum palustre</i>	Forms hummocks and dense carpets, often in shaded conditions. Indicative of nutrient enrichment (soaks and active flushes)	Low
<i>Sphagnum papillosum</i>	Lawn, hollow, and low hummock species	Moderate
<i>Sphagnum pulchrum</i>	Grows in lawns and hollows, more typical of western bogs	Moderate
<i>Sphagnum squarrosum</i>	Forms carpets and small mounds. Indicative of nutrient enrichment (soaks and active flushes)	Low
<i>Sphagnum subnitens</i>	Occurs as individual shoots or small cushions and lawns. Tolerant of minerotrophic conditions	Moderate
<i>Sphagnum tenellum</i>	Occurs as single shoots or weak cushions, typically in disturbed patches of the bog surface	Low

The site-specific target for the attribute bog moss (*Sphagnum*) species is: **Restore adequate cover of bog moss (*Sphagnum*) species to ensure peat-forming capacity.**

### 2.3.8 Typical ARB species: flora

Garriskil Bog supports the full complement of plant species typically associated with a true midland raised bog (see Section 1.1.1 above).

The key typical species that are indicative of high quality raised bog include *Sphagnum fuscum* and *S. austinii* which are associated with hummocks and *S. cuspidatum* and *S. denticulatum* associated with pools and hollows. All of these species have been reported from Garriskil Bog (Fernandez *et al.* 2005, 2014a, b).

The site-specific target for the attribute typical bog flora is: **Restore, where appropriate, typical active raised bog flora.**

### 2.3.9 Typical ARB species: fauna

As mentioned in section 1.1.3, a list of typical fauna specific to ARB has not been developed and the table contains species that use the wider raised bog habitat. This may be refined as more information becomes available.

Site specific information on the faunal assemblages associated with Garriskil Bog is currently limited. It is likely that most species groups referred to in section 1.1.3 occur on the bog.

The site-specific target for the attribute typical bog fauna is: **Restore, where appropriate, typical active raised bog fauna.**

### 2.3.10 Elements of local distinctiveness

A range of features may be associated with raised bogs which add to the scientific, historical, or conservation value of a bog. These can include geological, topographical, archaeological and hydrological features (e.g. soaks, lakes, flushes) and noteworthy species of flora and fauna (Cross 1990). Notable species of flora and fauna include those listed in the Habitats and Birds Directives, Red-listed species and other rare or localised species. For this attribute, features that are particularly associated with ARB are relevant.

#### 2.3.10.1 Site features

The most striking feature of the ARB areas on Garriskil Bog is the large area of inter-connecting pools, which cover approximately 25% of the bog dome. The pools are more typical of western bog sites that are intermediate between raised and blanket bogs. However, it is likely that the pools are a secondary feature, i.e. tears in the bog surface which have developed through subsidence caused by peat cutting and drainage around the margins of the bog.

There are a few flushes on the high bog, typified by the presence of *Molinia caerulea* and *Betula pubescens*. In one location, a large *Sphagnum* lawn flush occurs with *S. magellanicum* and *Menyanthes trifoliata* present.

#### 2.3.10.2 Rare flora

A relatively scarce lichen, *Cladonia rangiferina*, is found in abundance on and around hummocks in the ARB on the site (Douglas & Grogan 1986).

#### 2.3.10.3 Rare fauna

As mentioned above, there is a lack of current documented site-specific data relating to species that are particularly associated with ARB, including rare species.

In conclusion, the site-specific target for the attribute elements of local distinctiveness is: **Maintain features of local distinctiveness, subject to natural processes.**

### 2.3.11 Negative physical indicators

Raised bogs that have been damaged by marginal cutting and drainage, reclamation for agriculture, forestry activities, fire, surface drainage, or the lowering of regional water tables show a range of negative physical indicators (Cross 1990). Such negative physical features of ARB include: bare peat, algae dominated pools and hollows, marginal cracks, tear patterns, subsidence features such as dry peat and / or mineral mounds / ridges emerging or expanding, and burning evidence.

The NPWS conservation plan for Garriskil Bog noted that the most recent burn at the site was in 1997 (NPWS 2004). No fire damage was recorded during a survey undertaken in 2011 (Fernandez *et al.* 2014a, b).

Cattle grazing has been reported in the north-western part of the site. Severe poaching and bare peat areas were noted at this location by Kelly *et al.* (1995) and Fernandez *et al.* (2005, 2014a, b) and around the margin at other locations.

A block of conifer forestry occurs adjacent to the southern boundary outside the SAC. The River Riffey marks the boundary of the SAC and separates the conifer plantation from the cutover bog and other habitats in the SAC. The Sligo-Dublin railway runs along the northern and north-eastern boundary and is likely to have impacted on the high bog in the past.

No peat cutting currently occurs within the site, although 49% of the bog has been cutaway

since the 1840s. Subsidence is evident in the southern part of the high bog as described by Kelly *et al.* (1995).

The site-specific target for the attribute negative physical indicators is: **Negative physical features absent or insignificant.**

### **2.3.12 Vegetation composition: native negative indicator species**

Indicators of disturbance on a raised bog include species indicative of drying out conditions such as abundant *Narthecium ossifragum* and *Trichophorum germanicum*; *Eriophorum vaginatum* forming tussocks; abundant *Sphagnum magellanicum* in pools previously dominated by species typical of very wet conditions (e.g. *Sphagnum cuspidatum*). Indicators of frequent burning events include abundant *Cladonia floerkeana* and high cover of *Carex panicea* (particularly in the true midlands raised bog type).

The site-specific target for the attribute native negative indicator species is: **Native negative indicator species at insignificant levels.**

### **2.3.13 Vegetation composition: non-native invasive species**

Non-native invasive species that can commonly occur on raised bog habitats include: *Pinus contorta*, *Rhododendron ponticum*, and *Sarracenia purpurea* (Cross 1990).

Fernandez *et al.* (2005) mentioned a single large *Rhododendron ponticum* bush (2m high) to the west of flush Z surrounded by numerous seedlings and small plants (c.100 plants less than 1m high) covering a large area (50 m<sup>2</sup>). *Rhododendron ponticum* is spreading in this area. This bush and numerous seedlings were already noted by Kelly *et al.* (1995). Fernandez *et al.* (2014) survey in 2011 confirmed the spreading of *Rhododendron* (100 x 25m containing up to 100 plants and many seedlings).

Individual and groups of *Pinus* sp. trees are frequently found on the southern slopes of the high bog mainly in the marginal ecotope (Fernandez *et al.* 2005).

Invasive species are considered to have low intensity impact on the high bog habitats (Fernandez *et al.* 2014a, b).

The site-specific target for the attribute non-native invasive species: **Non-native invasive species at insignificant levels and not more than 1% cover.**

### **2.3.14 Air quality: nitrogen deposition**

Peatlands are highly sensitive to air pollution, particularly nitrogen deposition. Reactive nitrogen from fossil fuel combustion or intensive agriculture can contaminate rain and snow, causing soil acidification, nutrient enrichment, and a decline in species that are sensitive to these conditions. There is evidence that the combined impact of elevated nitrogen deposition and a warming climate could exceed the sum of the individual stressors and lead to a dramatic decline in the biodiversity of mosses, sensitive vascular plants, and microbes, potentially leading to catastrophic peat loss (PEATBOG project - <http://www.sste.mmu.ac.uk>).

Air pollution can change both the species composition and the functioning of peatlands. The primary atmospheric pollutant from the Industrial Revolution to the mid 1970s was sulphur deposition, but levels have since greatly declined. Reactive nitrogen (N) deposition (primarily NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup>), which can both acidify and eutrophy, became significantly elevated over a widespread area in the early to mid-20th century and is now the major pollutant in atmospheric deposition across most of Europe (Fowler *et al.* 2005).

Nitrogen is commonly a limiting terrestrial nutrient and in un-impacted peatlands it is tightly cycled. With long-term elevated N deposition, vegetation composition typically shifts toward species adapted to higher nutrient levels, with an overall loss of diversity (Malmer & Wallén 2005). In peatlands, field experiments with N additions within the current European range have shown significant declines in bryophyte species-richness and productivity, and shifts in composition toward vascular plants (Bobbink *et al.* 1998; Bubier *et al.* 2007). Community shifts toward more nitrophilous bryophytes in N-enriched regions such as parts of the Netherlands are also well documented (Greven 1992). In the UK, both a general survey of peatlands across the country (Smart *et al.* 2003), and a targeted study of *Calluna* moorland (Caporn *et al.* 2007) showed significant inverse relationships between levels of nitrogen deposition and species richness, with bryophytes particularly impacted. Changes in the vegetation also impact below-ground communities and biogeochemical processes.

Moderate increases in N deposition from a low level may increase *Sphagnum* and vascular plant productivity without an equal increase in decomposition rates, leading to enhanced carbon accumulation (Turunen *et al.* 2004). However, shifts in species composition from bryophytes to vascular plants may increase the production of easily-decomposable plant material, leading to higher rates of decomposition, and reduced carbon accumulation (Lamers *et al.* 2000; Bubier *et al.* 2007).

The particular sensitivity of nutrient-poor ombrotrophic peatlands to nitrogen enrichment is reflected in the low critical load threshold of between 5 and 10kg N/ha/yr for these ecosystems (Bobbink & Hettelingh 2011), a level which is exceeded over a significant portion of their range. An Irish study during the late 1990s undertaken by Aherne & Farrell (2000) concluded that total N deposition shows a strong east-west gradient, with lowest deposition in the west at 2kg N/ha/yr and highest in the east and south-east at 20kg N/ha/yr. Average N deposition over the Republic of Ireland was estimated to be approximately 12kg N/ha/yr. The study also concluded that the Critical Load Threshold for N was exceeded in at least 15% of ecosystems studied. The critical load applied to peatland ecosystems by Aherne & Farrell (2000) was 10kg N/ha/yr. This is in line with the recommendation by Bobbink & Hettelingh (2011) that the critical load should be set at the high end of the range in areas of high precipitation and at the low end of the range in areas of low precipitation assuming that Ireland represents a high precipitation area.

It is recommended in the case of Garriskil Bog that the level of N deposition should not exceed the low end of the range i.e. 5kg N/ha/yr. This recommendation is based on a precautionary approach, as the evidential basis for setting a higher level is not particularly strong as alluded to by Payne (2014). Total N deposition in the vicinity of Garriskil Bog as reported by Henry & Ahern (2014) is 19.3kg N/ha/yr.

The site-specific target for the attribute air quality is: **Air quality surrounding bog close to natural reference conditions. The level of N deposition should not exceed 5kg N/ha/yr.**

### 2.3.15 Water quality

Ombrotrophic peat waters found on the surface of raised bogs are characterised by low pH values (pH < 4.5) (Moore & Bellamy 1974) and also have low values of electrical conductivity. This is due to the fact that the raised bog system derives its mineral supply from precipitation, which is usually acidic and low in nutrients. Raised bog vegetation exchanges cations with protons to further reduce the pH.

Hydrochemistry varies in the areas surrounding a raised bog. Locally, conditions may be similar to the high bog due to a dominance of water originating from the bog. However, elsewhere in the marginal areas, there may be increased mineral and nutrient content of the

water due to regional groundwater influences, runoff from surrounding mineral soils, and the release of nutrients through oxidation of peat resulting from reduced water levels.

A brief hydrochemistry survey carried out as part by Kelly *et al.* (1995) indicated that generally cutover drains had limited groundwater influence since electrical conductivities were typical in the region of 100 $\mu$ s/cm. However, very few sample points were noted within this study, therefore it is not possible to definitely state that there was not upwelling of mineralised groundwater in the cutover.

The site-specific target for the attribute water quality is: **Water quality on the high bog and in transitional areas close to natural reference conditions.**

### 3 References

- Aherne, J., & Farrell, E.P. (2000) Final Report: Determination and mapping of critical loads for sulphur and nitrogen and critical levels for ozone in Ireland. Environmental Protection Agency, Dublin, 212pp.
- Bobbink, R., Hornung, M. & Roelofs, J.G.M. (1998) The effects of air—borne nitrogen pollutants on species diversity and semi—natural European vegetation. *Journal of Ecology* 86: 717–738.
- Bobbink, R. & Hettelingh, J.P. (2011) Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010. RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM).
- Bracken, F., McMahon, B. & Whelan, J. (2008) Breeding bird populations of Irish Peatlands: capsule peatlands are very important habitats for birds despite low species diversity. *Bird Study* 55 (2): 169-178.
- Bracken, F. & Smiddy, P. (2012) Lowland bogs, fens and reedswamps, pp. 73-89. In: Nairn, R., and O'Halloran, J. (eds.) *Bird Habitats in Ireland*. The Collins Press, Cork.
- Bubier, J., Moore, T. & Bledzki, L.A. (2007) Effects of nutrient addition on vegetation and carbon cycling in an ombrotrophic bog. *Global Change Biology* 13: 1168–1186.
- Caporn, S.J.M., Edmondson, J., Carroll, J.A., Pilkington, M. & Ray, N. (2007) Long-term impacts of enhanced and reduced nitrogen deposition on semi-natural vegetation. Report to Defra. Terrestrial Umbrella. Work Package 2: Impacts, Recovery and Processes. Task 4. Defra London.
- CEC (2007) Interpretation manual of European Union Habitats. Version EUR 27. European Commission, DG Environment, Brussels. Nature and Biodiversity.
- Colhoun, K. & Cummins, S. (2013) Birds of Conservation Concern in Ireland 2014–2019. *Irish Birds* 9: 523-544.
- Cross, J. (1990) The Raised Bogs of Ireland, their ecology, status and conservation. Report to the Minister of State at the Department of Finance. The Stationery Office, Dublin.
- Crushell, P.H., Schouten, M.G.C., Robroak, B.J.M. & van Duinen, G-J. (2008) The contribution of soak lakes to macroinvertebrate diversity of raised bogs in Ireland. In: Crushell, P.H. (2008). *Soak Systems of an Irish Raised Bog: a multidisciplinary study of their origin, ecology, conservation and restoration*. PhD thesis, Wageningen University, with a summary in Dutch and Irish.
- DAHG (2014) National Raised Bog SAC Management Plan. Draft for Consultation. Main report and appendices. Department of Arts, Heritage and the Gaeltacht.
- De Leeuw, J.P.M. (1986) Een onderzoek naar het voorkomen en de verspreiding van aquatische macro- en mirofauna in de Ierse hoogvenen. Deel 1: Macrofauna. *Aquatische Oecologie*, Katholieke Universiteit Nijmegen, Nijmegen, The Netherlands.
- Derwin, J. & MacGowan, F. (2000) Raised Bog Restoration Project: A Continuation of the Investigation into the Conservation and Restoration of Selected Raised Bog Sites in Ireland. Unpublished report, Dúchas the Heritage Service, Dublin
- Douglas, C. & Grogan, H. (1986). Survey to locate raised bogs of scientific interest in Counties Longford, Westmeath and Mayo. Unpublished report to the Forest and Wildlife Service, Dublin.
- Fernandez Valverde, F., Fanning, M., McCorry, M. & Crowley, W. (2005) Raised bog

monitoring project 2004-2005. Document 3: Site Reports and Maps Volume 1-5. Unpublished Report. National Parks and Wildlife Service, Dublin.

Fernandez, F., Connolly, K., Crowley, W., Denyer, J., Duff, K. & Smith, G. (2014a) Raised bog monitoring and assessment survey 2013. Irish Wildlife Manuals, No. 81. National Parks and Wildlife Service, Department of Arts, Heritage and Gaeltacht, Dublin, Ireland.

Fernandez, F., Connolly, K., Crowley, W., Denyer, J., Duff, K. & Smith, G. (2014b) Raised Bog Monitoring and Assessment Survey 2013 - Garriskil Bog (SAC 000679), Co. Westmeath - site report. National Parks and Wildlife Service, Department of Arts, Heritage and Gaeltacht, Dublin, Ireland.

Fossitt, J. (2000) A Guide to Habitats in Ireland. The Heritage Council, Ireland.

Fowler, D., Smith, R.I., Muller, J.B.A., Hayman, G. & Vincent, K.J. (2005) Changes in the atmospheric deposition of acidifying compounds in the UK between 1986 and 2001. *Environmental Pollution*, 137: 15-25.

Fox, A.D., Norriss, D.W., Stroud, D.A. & Wilson, H.J. (1994) Greenland White-fronted Geese in Ireland and Britain 1982/83-1993/94. Greenland White-fronted Goose Study Research Report No. 8.

Gore, A.J.P. (ed.) (1983) *Ecosystems of the world 4A. Mires: Swamp, bog, fen and moor. General studies.* Elsevier Scientific Publishing Company, Amsterdam.

Greven, H.C. (1992) Changes in the moss flora of the Netherlands. *Biological Conservation* 59: 133-137.

Hannigan, E., and Kelly-Quinn, M. (2011) Chapter 2.6 - Aquatic macro-invertebrate diversity. pp. 140-157 In: Renou-Wilson, F. (ed.) *BOGLAND: Sustainable Management of Peatlands in Ireland.* Environmental Protection Agency, Wexford.

Kelly, M.L. (1993) Hydrology, hydrochemistry and vegetation of two raised bogs in county Offaly. PhD thesis, Trinity College Dublin.

Kelly, L., Doak, M. & Dromey, M. (1995) Raised Bog Restoration Project: An Investigation into the Conservation and Restoration of Selected Raised Bog Sites in Ireland. Part 1 Summary Reports. National Parks & Wildlife Service, Department of Environment, Heritage and Local Government, Dublin.

Kelly, L. & Schouten, M.G.C. (2002) Vegetation. In: Schouten, M.G.C. (ed.), *Conservation and restoration of raised bogs: geological, hydrological and ecological Studies.* Dúchas – The Heritage Service of the Department of the Environment and Local Government, Ireland; Staatsbosbeheer, the Netherlands; Geological Survey of Ireland, Dublin. pp. 110-169.

Laine, J., Harju, P., Timonen, T., Laine, A., Tuittila, E.S, Minkkinen, K. & Vasander, H. (2009) The intricate beauty of Sphagnum mosses - A Finnish guide to identification. University of Helsinki Department of Forest Ecology Publications, 39: 1–190.

Lamers, L. P. M., Bobbink, R. & Roelofs, J. G. M. (2000) Natural nitrogen filter fails in polluted raised bogs. *Global Change Biology*, 6: 583–586.

Malmer, N. & Wallén, B. (2005) Nitrogen and phosphorus in mire plants: variation during 50 years in relation to supply rate and vegetation type. *Oikos*, 109: 539–554.

Moore, P.D. & Bellamy, D.J. (1974) *Peatlands.* Elek Science. London.


Nolan, M. (2013) Spiders (Araneae) of Irish raised bogs: Clara bog, Co. Offaly and Carrowbehy bog, Co. Roscommon. *Bulletin of the Irish Biogeographical Society* 37: 172-203.

- NPWS (2004). Conservation Plan 2004-2009: Garriskil Bog cSAC & SPA. Draft 2. Department of the Environment, Heritage, and Local Government.
- NPWS (2008) The Status of EU Protected Habitats and Species in Ireland. National Parks and Wildlife Service, Ireland.
- NPWS (2013) The Status of EU Protected Habitats and Species in Ireland. Version 1.0. Unpublished Report, National Parks and Wildlife Services. Department of Arts, Heritage and the Gaeltacht, Dublin, Ireland.
- O'Connell, C. (ed.) (1987) The IPCC Guide to Irish Peatlands. Irish Peatland Conservation Council, Dublin.
- O'Connell, P. (2011) Action Plan for Raised Bog Birds in Ireland 2011-2020. BirdWatch Ireland, Kilcoole, Co. Wicklow.
- O'Connor, Á., Reynolds, J.D. & Kavanagh, B. (2001) Aquatic macroinvertebrate colonisation of artificial water bodies in cutaway oceanic raised bog in Ireland. In: Rochfort, L. and Daigle, J.Y. (eds.), Proceedings of the 11<sup>th</sup> International Peat Congress. pp. 742-750.
- O'Sullivan, A. & Ryan, C. (1994). Garriskil Bog (Site Code: 679) National ASI Survey Site Card. NHA database, Dúchas, Dublin.
- Renou-Wilson, F., Bolger, T., Bullock, C., Convery, F., Curry, J., Ward, S., Wilson, D. & Müller, C. (2011) BOGLAND: Sustainable Management of Peatlands in Ireland. STRIVE Report Series No.75. Prepared for the Environmental Protection Agency. pp. 181.
- Reynolds, J.D. (1984a) Invertebrate survey of Irish midlands raised bogs. Bulletin of the British Ecological Society 15: 81-82.
- Reynolds, J.D. (1984b) Invertebrate fauna of Irish raised bogs. Part II: Odonata, aquatic Hemiptera and Trichoptera. Bulletin of the Irish Biogeographical Society 8: 98-102.
- Reynolds, J.D. (1985) Invertebrates of Lough Roe, Co. Offaly; a rare and endangered bogland habitat. Bulletin of the Irish Biogeographical Society 9: 41-45.
- Schouten, M.G.C. (1984) Some aspects of the ecogeographical gradient in the Irish ombrotrophic bogs, paper presented to 7th Int. Peat Congress, Dublin, vol. 1, pp. 414-432, The International Peat Society, Helsinki.
- Schouten, M.G.C. (ed.) (2002) Conservation and Restoration of Raised Bogs – geological, hydrological and ecological studies. Dúchas – The Heritage Service of the Department of the Environment and Local Government, Ireland; Staatsbosbeheer, The Netherlands; and The Geological Survey of Ireland. pp. 220.
- Smart, S.M., Robertson, J., Shield, E.J. & van de Poll, M.H. (2003) Locating eutrophication effects across British vegetation between 1990 and 1998. Global Change Biology 9: 1763-1774.
- Tomassen H.B.M., Smolders A.J.P., Limpens J., Lamers, L.P.M. & Roelofs, J.G.M. (2004) Expansion of invasive species on ombrotrophic bogs: desiccation or high N deposition? Journal of Applied Ecology. 41: 139–150.
- Turunen, J., Roulet, N.T., Moore, T.R. & Richard, P.J.H. (2004) Nitrogen deposition and increased carbon accumulation in ombrotrophic peatlands in eastern Canada. Global Biogeochemical Cycles. 18 (3): GB3002.
- Van Duinen G.A. (2013) Rehabilitation of aquatic invertebrate communities in raised bog landscapes. PhD thesis, Radboud University Nijmegen, the Netherlands.
- Wilson, H.J. (1990) Birds of raised bogs. pp. 29-36. In: Cross, J. (ed.) The Raised Bogs of


Ireland, their ecology, status and conservation. Report to the Minister of State at the Department of Finance. The Stationery Office, Dublin.

Wisdom, R. & Bolger, T. (2011) Chapter 2.4 - Terrestrial invertebrate biodiversity. pp. 103-121  
In: Renou-Wilson, F. (ed.) BOGLAND: Sustainable Management of Peatlands in Ireland. Environmental Protection Agency, Wexford.

Map 1: Extent of potential active raised bog on Garriskil Bog.



Map 2: Distribution of raised bog ecotopes on Garriskil Bog.



Map 3: Digital elevation model and drainage patterns at Garriskil Bog.

