

NPWS

Lisnageeragh Bog and Ballinastack
Turlough SAC
(site code 296)

**Conservation objectives supporting document-
Turloughs**

Version 1

March 2016

Contents

Contents.....	1
1. Introduction	3
1.1 Turloughs.....	3
1.2 Lisnageeragh Bog and Ballinastack Turlough SAC	3
1.3 Conservation objectives	3
2. Area.....	4
3. Range	4
4. Structure and functions	4
4.1 Hydrological regime	5
4.1.1 Hydrological regime: groundwater contribution	5
4.1.2 Hydrological regime: flood duration	5
4.1.3 Hydrological regime: flood frequency.....	6
4.1.4 Hydrological regime: flood area	6
4.1.5 Hydrological regime: flood depth	6
4.1.6 Hydrological regime: permanently flooded/wet areas.....	6
4.2 Soil type.....	7
4.3 Soil nutrient status.....	7
4.4 Physical structure: Bare ground	7
4.5 Chemical processes: Calcium carbonate deposition.....	7
4.6 Water quality	8
4.6.1 Water quality: nutrients	8
4.6.2 Water quality: colour	9
4.6.3 Water quality: phytoplankton biomass.....	9
4.6.4 Water quality: epiphyton biomass.....	10
4.7 Active peat formation	10
4.8 Vegetation composition: vegetation communities	10
4.9 Vegetation composition: vegetation zonation	11
4.10 Vegetation structure: sward height.....	11
4.11 Typical species	12

4.11.1 Typical species: terrestrial, wetland and aquatic plants	12
4.11.2 Typical species: aquatic invertebrates	14
4.11.3 Typical species: other invertebrates	15
4.11.4 Typical species: birds.....	16
4.12 Fringing habitats	16
4.13 Vegetation structure: turlough woodland	17
5. References	17
Appendix 1 Map of Ballinastack Turlough	24
Appendix 2 Orthophotograph of Ballinastack Turlough.....	25

1. Introduction

1.1 Turloughs

“Turloughs” (habitat code 3180) is a priority habitat in Annex I of the Habitats Directive. A turlough is a topographic depression in karst which is intermittently inundated on an annual basis, mainly from groundwater, and which has a substrate and/or ecological communities characteristic of wetlands (Working Group on Groundwater, 2005; Tynan *et al.*, 2007). Turloughs are semi-natural habitats that are virtually unique to Ireland. They generally flood in winter and dry out in summer, but there may be other sporadic rises in response to high rainfall. Turloughs are considered to be of high conservation value for their plant, invertebrate (both terrestrial and aquatic) and bird communities (Goodwillie, 1992, 2003; NPWS, 2007, 2008; Sheehy Skeffington, 2006). They are grass- or sedge-dominated habitats, often with areas of fen, marsh or permanent pond. Many turloughs show a distinctive zonation of herbaceous perennials which relates to the depth and duration of flooding. Turloughs intergrade into many other wetland types, from marl lakes and ponds to swamp, to fen and even to coastal lagoons and marine habitats.

The interpretation manual of EU habitats gives the following description: *“Temporary lakes principally filled by subterranean waters and particular to karstic limestone areas. Most flood in the autumn and then dry up between April and July. However, some may flood at any time of the year after heavy rainfall and dry out again in a few days; others, close to the sea, may be affected by the tide in summer. These lakes fill and empty at particular places. The soils are quite variable, including limestone bedrock, marls, peat, clay and humus, while aquatic conditions range from ultra oligotrophic to eutrophic. The vegetation mainly belongs to the alliance Lolio-Potentillion anserinae Tx. 1947, but also to the Caricion davallianae Klika 1934.”* (CEC, 2013).

In Ireland, turloughs are considered to be in poor/inadequate conservation status as a result of nutrient enrichment, inappropriate grazing and drainage pressures (NPWS, 2007, 2008, 2013).

1.2 Lisnageeragh Bog and Ballinastack Turlough SAC

Lisnageeragh Bog and Ballinastack Turlough SAC comprises a large raised bog and a small turlough, situated about 3 km north-east of Glenamaddy in north-east Co. Galway. It is selected for four habitats listed on Annex I of the EU Habitats Directive, two of which are priority habitats (turloughs* and active raised bogs*).

Ballinastack Turlough is located at the north-west of the SAC. The eastern side of the turlough merges with the raised bog. There are peat deposits within the turlough and the vegetation is sedge dominated. It was site 54 in Coxon (1986, 1987 a and b) and number 18 in Goodwillie (1992).

1.3 Conservation objectives

A site-specific conservation objective aims to define the favourable conservation condition of a habitat or species at site level. The maintenance of habitats and species within sites at favourable condition will contribute to the maintenance of favourable conservation status of those habitats and species at a national level.

Conservation objectives are defined using attributes and targets that are based on parameters as set out in the Habitats Directive for defining favourable status, namely area, range, and structure and functions.

Grazing is integral to the ecology of turloughs and it is important that appropriate grazing levels are maintained. Over-grazing and, in particular, under-grazing are considered significant threats to turlough conservation status. Turlough diversity will benefit from within-site spatial variations in grazing regime. Both the timing (post flood-recession) and intensity of grazing are important. Various attributes set out below and various authors (e.g. Goodwillie, 1992, 2003; Bond, 1997; Ní Bhriain *et al.*, 2002; Moran, 2005; Regan, 2005; Ryder *et al.*, 2005; Sheehy Skeffington *et al.*, 2006; Moran *et al.*, 2008; Kimberley *et al.*, 2012) provide further information on the importance of grazing.

Attributes and targets may change/become more refined as further information becomes available.

2. Area

Goodwillie (1992) calculated the area of Ballinastack turlough at 24.9 ha, but could not determine the eastern boundary where it opens into the raised bog. The area covered by the turlough as mapped in the conservation objective is 23.7 ha (see Appendix 1). This map was created by digitising the outer boundary of the Goodwillie (1992) vegetation units.

The natural transition between Ballinastack turlough and the bog was altered by historic turf-cutting and agricultural land use. Restoration of the hydrological regime to support the active raised bog habitat should also aim to restore the vegetation transition between the two priority habitats.

It is possible that the full extent of flooding and turlough vegetation is not contained within the SAC boundary.

The **target** for habitat area is: stable or increasing, subject to natural processes.

3. Range

The distribution of turloughs in Lisnageeragh Bog and Ballinastack Turlough SAC is shown in Appendix 1. This map was created by digitising the outer boundary of the turlough vegetation as mapped in Goodwillie (1992).

The **target** for the habitat distribution attribute is: no decline, subject to natural processes.

4. Structure and functions

Structure and functions relates to the physical components of a habitat (“structure”) and the ecological processes that drive it (“functions”). For turloughs these include attributes such as hydrology, soils and various water quality attributes.

4.1 Hydrological regime

Hydrology is the key driver of turlough ecology. The different turlough communities, assemblages and species are affected by various hydrological attributes. The most important of these are described below.

Goodwillie (1992), recorded two small streams entering Ballinastack turlough from the south and a ditch from the eastern side, all of which discharge to the swallow holes towards the north of the turlough (marked on six inch map and in GSI karst database). It is possible that he overlooked some drains, or that more drainage has occurred within the turlough since his survey.

In restoring the hydrological regime of the raised bog, consideration should be given to restoration of the turlough hydrological regime to support the turlough communities and transitional vegetation between these two priority habitats.

4.1.1 Hydrological regime: groundwater contribution

Turloughs flood mainly as a result of rising groundwater levels. The groundwater contribution exerts a strong influence on turlough ecology, notably through mineral contributions (e.g. calcium carbonate). It is, therefore, important that groundwater makes a high percentage contribution to turlough floodwaters. At Ballinastack turlough, there may be a significant contribution of base-poor surface run-off from areas of raised bog, however it is uncertain to what extent the natural groundwater : surface-water ratio has been altered by drainage of the peatland.

The **target** for the attribute hydrological regime, groundwater contribution is: Maintain appropriate groundwater contribution necessary for the natural functioning of the habitat.

4.1.2 Hydrological regime: flood duration

Flood duration is considered to be the dominant hydrological driver of turlough vegetation. Flood duration is also important for turlough invertebrates. Both terrestrial and aquatic species and communities must be considered when investigating the influence of flood duration. A number of hydrological indicators of flood duration have been used in turlough studies. The recession constant (k) characterises the rate at which floodwaters recede, with a low k indicating a more gradual recession and, hence, longer flood duration (Paul Johnston, pers. comm., Tynan, *et al.*, 2007; Naughton, 2011). The usefulness of other hydrological functions, such as gamma function, for characterising turlough hydrology is also being investigated (Paul Johnston, pers. comm.; Naughton, 2011). It should also be possible to use plant species as indicators of the extent and duration of flooding in turloughs (see Goodwillie, 1992; Goodwillie *et al.*, 1997; Waldren, 2015). Consequently, both hydrological and vegetation indicators should be considered when measuring this attribute. Ballinastack turlough is thought to hold some water for much of the winter and to drain rapidly in dry periods. Drainage activities in the area may have affected the natural flood duration.

The **target** for the attribute hydrological regime, flood duration is: Maintain appropriate hydrological regime necessary for the natural functioning of the habitat.

4.1.3 Hydrological regime: flood frequency

Hydrological regime is highly variable among turloughs. Some turloughs demonstrate a characteristically low flood frequency (flooding just once most years for six months or more), whilst others have more dynamic water levels with higher flood frequencies (Naughton, 2011).

The **target** for the attribute hydrological regime, flood frequency is: Maintain natural annual temporal patterns in flood frequency.

4.1.4 Hydrological regime: flood area

The extent of flooding determines the overall area of wetland vegetation, as well as the area of each characteristic vegetation community. Flood area is also important to aquatic invertebrate communities, determining the extent of the available habitat and influencing competitive and predation interactions. A number of hydrological indicators might be used to measure the flood area, such as the average annual maximum flood area, the stage (level):area relationship, or the areal (flood) reduction rate (Porst, 2009; Naughton, 2011; Naughton *et al.*, 2012). Drainage activities in the area may have affected the natural flood area.

The **target** for the attribute hydrological regime, flood area is: Maintain natural temporal pattern in flood area.

4.1.5 Hydrological regime: flood depth

Flood depth is directly related to the other measures of hydrological regime (duration, area etc.) and basin morphology. The depth (and area/duration) of flooding could be impacted by increases or decreases in the discharge to the turlough (resulting in changes to the water level/stage) or by re-contouring the land (changing ground elevation and topography). Changes to flood depth will affect flood duration and, therefore, the vegetation communities. A number of hydrological indicators might be used to measure the flood depth, such as average annual maximum flood depth, the stage/depth:time series, or the average depth.

The **target** for the attribute hydrological regime, flood depth is: Maintain natural temporal and spatial patterns in flood depths.

4.1.6 Hydrological regime: permanently flooded/wet areas

Most turloughs have areas of permanent or semi-permanent flooding or saturated soils (i.e. areas that do not dry out every year). These may reflect the presence of aquitards (low permeability deposits such as peat or marl) rather than the local groundwater level during the turlough's 'dry phase'. These ponds and wet areas act as refuges for aquatic invertebrate species during the dry phase. They are particularly important to high conservation value species and assemblages, such as the moss-edge dwelling aquatic coleopteran community (Bilton, 1988; Bilton and Lott, 1991; Foster *et al.*, 1992; Bradish *et al.*, 2002). These areas also support important aquatic macrophyte communities, notably charophytes. Much of Ballinastack has peat soil with wet *Carex nigra* vegetation, and permanently wet areas also appear to be associated with the streams.

The **target** for the attribute hydrological regime, permanently flooded/wet areas is: Maintain any areas of permanent or semi-permanent flooding or water-logging.

4.2 Soil type

Turloughs are characterised by a broad range of wetland soils from mineral alluvium to peat and marl (Coxon, 1987b; Goodwillie, 1992; Kimberley, 2008; Waldren, 2015). Turlough soil type is largely determined by geology, morphology and hydrology (MacGowran, 1985; Coxon, 1986). Soil type is an important driver of vegetation and influences grazing and other land management practices (Moran *et al.*, 2008; Kimberley *et al.*, 2012). Ballinastack turlough is dominated by peat, with some sandy silt in the north-west (Goodwillie, 1992).

The **target** for the attribute soil type is: Maintain variety, area and extent of soil types necessary to support turlough vegetation and other biota.

4.3 Soil nutrient status

Nutrient concentrations are typically low in turlough soils, but are highly variable (temporally, among soil types, and with flood duration and land-use) and exert an influence on vegetation communities (Sarah Kimberley, pers. comm.; Waldren *et al.*, 2002; Kimberley, 2008; Kimberley *et al.*, 2012; Kimberley and Waldren, 2012). Maintenance of the natural nutrient status is important, as increases in soil nutrient status can significantly alter the vegetation communities and impact on sensitive aquatic assemblages.

The **target** for the attribute soil nutrient status is: Maintain nutrient status (phosphorus and nitrogen concentrations) appropriate to soil types and vegetation communities.

4.4 Physical structure: Bare ground

Bare ground in turloughs occurs naturally in areas with long flood duration and late drying and where fine sediments (alluvium) are deposited by the floodwaters. Trampling by grazers can also result in areas of bare ground. Bare ground is important for certain vegetation communities, such as *Chenopodium rubri* (Habitats Directive Annex I habitat 3270), which is dominated by short-lived annuals, including the rare *Callitriche palustris*, *Limosella aquatica* and *Rorippa islandica*. It is also important to invertebrate communities, notably ground beetles (Regan, 2005a; Sheehy Skeffington *et al.*, 2006).

The **target** for the attribute Physical structure, bare ground is: Maintain sufficient wet bare ground, as appropriate.

4.5 Chemical processes: Calcium carbonate deposition

Calcium carbonate (CaCO₃) deposition is a feature of very many turloughs (Coxon, 1994; Goodwillie, 2003). Both physical and biological processes clearly play a part in precipitating CaCO₃ from saturated/ super-saturated groundwater, as evidenced by deposits on turlough vegetation after flood recession (Goodwillie, 2003; Jim Ryan, pers. comm.). While it may appear difficult to change CaCO₃ precipitation, it could be affected by drainage activities in the turlough or the zone of

contribution (groundwater catchment) impacting the CaCO₃ concentration in the floodwater, or by changes to biological communities, impacting the precipitation processes. Coxon (1986), Goodwillie (1992) and Coxon and Coxon (1994) found no evidence of marl in Ballinastack turlough. Á. O Connor however observed marl underlying peat near the swallow hole and it is likely that historic precipitation of marl facilitated peat formation. While marl may underlie much of the peat in Ballinastack, recent CaCO₃ precipitation may, naturally, be limited at the site, or may have been impacted by increased run-off of base-poor water from surrounding raised bog.

The **target** for the attribute chemical processes, Calcium carbonate deposition is: Maintain CaCO₃ deposition rate and/or soil concentration.

4.6 Water quality

Turloughs, being groundwater fed, are typically associated with high water quality. This is demonstrated by naturally low dissolved nutrients, clear water and low algal growth. Water quality can be measured by the following attributes:

4.6.1 Water quality: nutrients

The WFD Working Group on Groundwater developed a methodology to assess the risk to turloughs from phosphorus pollution, which established provisional phosphorus thresholds for turlough flood waters and the up-gradient groundwater (Working Group on Groundwater, 2005). Two total phosphorus (TP) thresholds were used for the turlough water, depending on the site's fertility and sensitivity to phosphorus enrichment. A threshold of ≤10µg/l TP was set for the least fertile ('extremely sensitive') turloughs as defined by the vegetation communities, and a threshold of ≤20µg/l TP for all other turloughs. These values were based on the Phosphorus Regulations' standards for TP in Irish lakes, which indicate that when mean TP is ≤10µg/l a lake is oligotrophic and >10 to ≤20µg/l mesotrophic (McGarrigle *et al.*, 2002, Appendix I).

Work undertaken by TCD clearly demonstrated that turloughs behave like lakes in terms of their phytoplankton biomass and their TP:phytoplankton biomass (chlorophyll *a*) relationship (Norman Allott, pers. comm.; Cunha Pereira *et al.*, 2010; NPWS, 2013; Waldren 2015). The TCD team considered that the natural background levels for all turloughs were likely to be <10µg/l (Norman Allott and Catherine Coxon, pers. comm.). Waldren (2015) concluded that <10µg/l TP indicates 'Very good' quality, 10-20µg/l TP indicates 'Good' quality, 20-50µg/l TP 'Intermediate' quality, and >50µg P/l indicates 'Bad' quality.

While it may ultimately be necessary to set site-specific TP targets for turloughs, a target of ≤10µg/l TP is used here for more oligotrophic sites dominated by fen and other sedge-rich vegetation of low-fertility and high species diversity, and a target of ≤20µg/l TP for less oligotrophic sites and there should be no deterioration in mean TP values.

The Working Group on Groundwater (2005) categorised Ballinastack turlough as naturally highly sensitive to nutrient enrichment, and considered that it is likely to have been impacted by groundwater enrichment. Based on this assessment, the appropriate target for Ballinastack turlough is ≤20µg/l TP, however this should be confirmed through investigation of the phosphorus concentration of the flood waters and evidence of eutrophication responses in the biota. Where the

average annual TP concentration is $\leq 10\mu\text{g/l}$ TP, there should be no deterioration, i.e. the target should be revised to $\leq 10\mu\text{g/l}$ TP.

The **target** for the attribute water quality, nutrients is: Maintain average annual TP concentration of $\leq 10\mu\text{g/l}$ TP, or $\leq 20\mu\text{g/l}$ TP, as appropriate.

4.6.2 Water quality: colour

Colour is typically low in turlough floodwaters, however drainage of peatland in the Slieve Aughty Mountains is believed to have increased the colour in Blackrock, Lough Coy, Garryland and Caherglassaun turloughs and to have altered the nutrient-algal biomass relationship and phytoplankton species composition (Norman Allott, pers. comm.; Cunha Pereira *et al.*, 2010). Reduced light penetration is the most likely causative factor, although sequestration of ions can also affect phytoplankton growth (Norman Allott, pers. comm; Cunha Pereira *et al.*, 2010). Increased water colour could also impact on the primary productivity of rooted plants, either the truly aquatic plants or those wetland species that can grow even when flooded. All TCD study turloughs, other than those on the Gort-series (72-85 mg/L PtCo), had colour of < 48 mg/L PtCo, with the Burren turloughs all ≤ 14 mg/L PtCo (Cunha Pereira *et al.*, 2010; Waldren, 2015). Drainage of surrounding raised bog may have artificially increased water colour at Ballinastack turlough.

The **target** for the attribute water quality, colour is: Maintain/restore appropriate water colour.

4.6.3 Water quality: phytoplankton biomass

Phytoplankton biomass as chlorophyll *a* is not significantly lower in turloughs than that found in permanent lakes (Cunha Pereira *et al.* 2010). Peak chlorophyll *a* concentration was recorded in winter in the TCD study turloughs (Norman Allott, pers. comm; Cunha Pereira *et al.*, 2010). It is probable that the decline in phytoplankton biomass in spring in turloughs is, at least in part, the result of grazing by zooplankton, whose development is likely dependent on temperature triggers.

The targets below are based on the OECD fixed boundary system for annual mean and annual peak chlorophyll *a* (OECD, 1982). Lower thresholds apply for the more oligotrophic and sensitive sites. Waldren (2015) used maximum recorded Chlorophyll *a* $> 10 \mu\text{g/l}$ as a negative indicator for conservation condition in turloughs. See also Section 4.6.1 above, water quality: nutrients and the Working Group on Groundwater (2005) for more information on categorising the sensitivity of turloughs.

The chlorophyll *a* standards in the European Communities Environmental Objectives (Surface Water) Regulations (S.I. 272 of 2009) could not be adopted as they are based on “growing season (March to October) mean values¹”, which includes a turlough’s ‘dry phase’. In addition S.I. 272 of 2009 states that the EQR boundary conditions are not yet developed for shallow calcareous lakes, which is the lake type most similar to turloughs.

¹ a minimum of four samples is required, distributed throughout the growing season.

The **target** for the attribute water quality, phytoplankton biomass is: Maintain appropriate chlorophyll *a* concentrations (annual mean <2.5 or <8µg/l, annual maximum values ≤8 and <25µg/l, as appropriate).

4.6.4 Water quality: epiphyton biomass

Patches of filamentous algae are a common occurrence in turloughs, however extensive algal mats are only found in turloughs with high average TP concentrations (i.e. ≥20µg/l) (Normal Allott, pers. comm.). The filamentous forms recorded include *Oedogonium*, *Spirogyra* and *Mougeotia* species (Normal Allott, pers. comm.). A few turloughs with high TP did not develop extensive algal mats, at least partly because of high water colour (Normal Allott, pers. comm.). Waldren (2015) used the 'Presence of filamentous algal mats covering at least 2% of turlough area on at least one occasion over three years of observation' as a negative indicator for turlough conservation condition

As a general rule, therefore, higher TP concentrations increase epiphyton production in turloughs. Owing to the fact that epiphyton is typically deposited onto the vegetation as the floodwaters recede, the decomposition of these algae is a pathway for nutrients from water to soil and rooted vegetation.

The **target** for the attribute water quality, epiphyton biomass is: Maintain trace/ absent epiphyton as algal mats (<2% cover).

4.7 Active peat formation

Peat formation is not a feature of all turloughs, but is associated with sedge-rich, fen-type vegetation communities. The duration of flooding is the key determinant of peat formation in turloughs. The water level needs to be slightly below to above the soil surface for approximately 90% of the time for peat to form (Jim Ryan, pers. comm.). Active peat formation can be threatened by drainage and other earth movements, as well as changes in hydrological regime. Note: peat formation in turloughs can be affected by natural changes in hydrology (Coxon and Coxon, 1994).

Peat is the dominant substratum in Ballinastack turlough (Goodwillie, 1992). It is likely that drainage activities have impacted on peat formation in the turlough and restoration measures for active raised bog should consider the potential for restoring active peat formation in the turlough.

The **target** for the attribute active peat formation is: Restore active peat formation, where appropriate.

4.8 Vegetation composition: vegetation communities

The vegetation of turloughs has been described and classified a number of times, notably by Goodwillie (1992, 2003), Goodwillie *et al.* (1997), O'Connell *et al.* (1994), MacGowran (1985), Regan *et al.* (2007) and Sharkey (2012). The conservation value of the described vegetation communities was considered by most of these authors (e.g. Goodwillie, 1992; O'Connell *et al.*, 1994; Regan, 2005a; Sharkey, 2012; Waldren, 2015). The WFD Working Group on Groundwater classified sensitive vegetation communities by assigning Ellenberg N (fertility) values (Hill *et al.*, 1999) to data from Goodwillie (1992), Goodwillie *et al.* (1997) and NPWS surveys (Working Group on Groundwater,

2005). Waldren (2015) developed positive and negative indicator communities for turloughs. The characteristic turlough communities identified as positive indicators (turlough-type dependent) included the *Eleocharis acicularis* community, *Carex* fen (*Molinia caerulea*-*Carex panicea* community), *Schoenus nigricans* fen and flooded woodland and pavement communities (Waldren, 2015).

Turlough vegetation communities also support invertebrate species and assemblages. The association between the turlough moss-edge water beetle assemblage and moss-rich fen vegetation communities is particularly notable.

The vegetation of Ballinastack turlough is described in Goodwillie (1992). It is dominated by 'wet *Carex nigra*' (6B) (10.1 ha) and 'peat grassland' (2D) (7.0 ha), and also has areas of 'poor grassland' (2B) (2.3 ha), 'sedge fen' (5D) (2.1 ha), 'peaty *Carex nigra*' (6D) (1.9 ha), 'species rich *Potentilla reptans*' (4B) (0.4 ha) and '*Lolium* grassland' (2A) (0.3 ha). Goodwillie (1992) notes that the *Carex* is ungrazed, and was the only substantial area of ungrazed wet *Carex nigra* he encountered. As the it has not been surveyed since 1992, however, there may have since been changes to the land use and the vegetation at Ballinastack.

The **target** for the attribute vegetation composition, vegetation communities is: Maintain area of sensitive and high conservation value vegetation communities/units.

4.9 Vegetation composition: vegetation zonation

Sheehy Skeffington *et al.* (2006) provide a good commentary and summary of the studies on turlough vegetation zonation. Praeger (1932) was one of the first to record the zonation from dry soil species at the turlough margins to amphibious and aquatic species in the areas with the longest flood duration. This zonation has been recorded by many authors (Ivimey-Cook and Proctor, 1966; O'Connell *et al.*, 1984; Goodwillie, 1992; Goodwillie *et al.*, 1997). Goodwillie (2003) discusses the influence of flood duration, the timing of flooding and other hydrological characteristics on vegetation zonation.

Goodwillie (1992) noted "At Ballinastack a nice transect exists from cutover raised bog, through Peat Grassland (2D), Peaty *Carex nigra* (6D), Wet *Carex nigra* (6B) to Poor Grassland (2B) around the swallow hole".

Maintaining a diversity of vegetation zones (community patches) also helps support invertebrate diversity.

The **target** for the attribute vegetation composition, vegetation zonation is: Maintain the vegetation zonation/mosaic characteristic of the site.

4.10 Vegetation structure: sward height

Vegetation height varies over time in turloughs, as a result of the timing of flood recession and the plants' growth periods, as well as the grazing regime. Vegetation diversity can be maximised by employing a variable grazing regime across the turlough. Variation in vegetation height is also important to invertebrate communities, with some species being dependent on bare soil, some

associated with grazed, short vegetation, whilst others, notably Diptera and Lepidoptera, require taller herbaceous vegetation and scrub, respectively (Bond, 1997; Good and Butler, 2001; Ní Bhriain *et al.*, 2002; Moran, 2005; Regan, 2005a; Ryder *et al.*, 2005; Sheehy Skeffington *et al.*, 2006).

Moran *et al.* (2008) state “Given that different species of both plants and animals will respond differently to differing management conditions, maintenance of heterogeneity is vital for the maintenance of maximum biodiversity”.

The **target** for the attribute vegetation structure, sward height is: Maintain sward heights appropriate to the vegetation unit, and a variety of sward heights across the turlough.

4.11 Typical species

As wetlands with distinct terrestrial and aquatic phases, turloughs have a range of typical species that can broadly be divided into wetland and aquatic species. In listing the typical species for the various groups below, strong emphasis has been placed on those that are indicative of good condition in turloughs (positive indicator species) and/or are known to be restricted to or have most occurrences in turloughs (characteristic species). The species lists highlight the rare and threatened species found in turloughs.

4.11.1 Typical species: terrestrial, wetland and aquatic plants

Table 1 lists the typical turlough plant species (NPWS, 2013). Various botanical and review studies list characteristic turlough plants (e.g. Goodwillie, 1992, 2003; Goodwillie *et al.*, 1997; Sheehy Skeffington *et al.*, 2006; Sharkey, 2012; Waldren, 2015). Table 2 lists other more widespread vascular plant species that are commonly encountered in turloughs, often at high cover abundance.

The TCD integrated turlough project identified the following as positive turlough indicator species: *Potentilla fruticosa*, *Viola persicifolia*, *Teucrium scordium*, *Limosella aquatica*, *Plantago maritima*, *Rorippa islandica* and *Frangula alnus* (Steve Waldren, pers. comm.; Waldren, 2015).

The plant list in Table 1 includes species typical of the Annex I habitat ‘Rivers with muddy banks with *Chenopodium rubri* p.p. and *Bidention* p.p. vegetation’ (3270), which is a community of short-lived annual species associated with turloughs in Ireland (NPWS, 2013). Other species associated with habitat 3270 include *Atriplex prostrata*, *Gnaphalium uliginosum*, *Rorippa palustris*, *Persicaria hydropiper*, *Chenopodium rubrum*, *Juncus bufonius* and, *Bidens tripartita*. It should be noted that habitat 3270 was not recorded in Ballinastack turlough.

Waldren (2015) said “many of the characteristic plant species encountered in turloughs also occur in other wetlands, or indeed in well-drained calcareous habitats; what is unique about turlough vegetation is the juxtaposition of ecologically different species along short but strong ecological gradients.” Rather than recommending a revised list of typical species, the authors identified plant indicator species that are indicators for flood duration and nutrient status (Waldren, 2015).

Lockhart *et al.* (2012) noted that *Cinclidotus fontinaloides* is characteristic of turloughs and that the rare *Pseudocalliergon lycopodioides* and *Drepanocladus sendtneri* can be locally abundant in turloughs. Some of the best, fen-vegetation-rich turloughs support *Pseudocalliergon trifarium* (Lockhart *et al.*, 2012).

Table 1 The typical plant species of turloughs. Species indicated by * are considered turlough specialists, though not necessarily restricted to turloughs. † indicates typical species of habitat 3270. Regional Red List status is given, where relevant, as a superscript (bryophytes only).

Angiosperms:
Dicots: <i>Callitriche palustris</i> *†, <i>Frangula alnus</i> (prostrate form*), <i>Galium boreale</i> , <i>Limosella aquatica</i> *†, <i>Persicaria minor</i> †, <i>Plantago maritima</i> , <i>Potentilla fruticosa</i> *, <i>Ranunculus repens</i> (form with highly dissected leaves*), <i>Rhamnus cathartica</i> , <i>Rorippa islandica</i> *†, <i>Teucrium scordium</i> *, <i>Viola persicifolia</i> *
Monocots: <i>Alopecurus aequalis</i> †, <i>Carex viridula</i> agg., <i>Eleocharis acicularis</i> †, <i>Schoenus nigricans</i>
Bryophytes:
Mosses: <i>Cinclidotus fontinaloides</i> , <i>Drepanocladus sendtneri</i> ^{NT} , <i>Pseudocalliergon lycopodioides</i> ^{VU} , <i>Pseudocalliergon trifarium</i> ^{VU}
Liverworts: <i>Riccia cavernosa</i> †
Pteridophytes:
<i>Ophioglossum vulgatum</i>

Table 2 Widespread plant species commonly found in turloughs, frequently at high abundance. Species indicative of oligotrophic conditions are given a ‡.

Angiosperms:
Dicots: <i>Baldellia ranunculoides</i> ‡, <i>Filipendula ulmaria</i> , <i>Hydrocotyle vulgaris</i> , <i>Leontodon hispidus</i> , <i>Littorella uniflora</i> ‡, <i>Mentha aquatica</i> , <i>Myosotis scorpioides</i> , <i>Persicaria amphibia</i> , <i>Persicaria hydropiper</i> , <i>Potamogeton gramineus</i> , <i>Potamogeton polygonifolius</i> ‡, <i>Potentilla anserina</i> , <i>Potentilla reptans</i> , <i>Prunus spinosa</i> , <i>Ranunculus flammula</i> , <i>Ranunculus trichophyllus</i> , <i>Rumex acetosa</i> , <i>Scorzonerooides autumnalis</i> , <i>Thymus polytrichus</i> , <i>Trifolium repens</i> , <i>Viola canina</i>
Monocots: <i>Agrostis stolonifera</i> , <i>Carex hirta</i> , <i>Carex hostiana</i> , <i>Carex nigra</i> , <i>Carex panicea</i> , <i>Deschampsia caespitosa</i> , <i>Eleocharis multicaulis</i> ‡, <i>Eleocharis palustris</i> , <i>Eleogiton fluitans</i> ‡, <i>Festuca arundinacea</i> , <i>Festuca rubra</i> , <i>Glyceria fluitans</i> , <i>Juncus articulatus</i> , <i>Juncus bulbosus</i> ‡, <i>Molinia caerulea</i> , <i>Phalaris arundinacea</i> , <i>Poa annua</i>
Bryophytes:
<i>Fontinalis antipyretica</i> , <i>Scorpidium revolvens</i>

Further investigation into the charophyte flora and other aquatic macrophytes of turloughs is required.

Goodwillie (1992) recorded a variety of typical turlough species at Ballinastack turlough, but no rare plants.

The **target** for the attribute typical species, terrestrial, wetland and aquatic plants is: Maintain typical species within and across sites

4.11.2 Typical species: aquatic invertebrates

NPWS (2013) provided an updated list of typical aquatic invertebrate species of turloughs (see Table 3), and the available information on this group is reviewed by Sheehy Skeffington *et al.* (2006) and Reynolds (2016). The species of greatest conservation importance in turloughs include the rare chydorid, *Eurycercus glacialis* (see Duigan and Frey, 1987a, b; Reynolds, 1997, 2000; Reynolds and Marnell, 1999; Reynolds *et al.*, 2004), the copepod *Diaptomus castor* (see Ali *et al.*, 1987; Reynolds, 1997), the near threatened odonate *Lestes dryas* (see Nelson and Thompson, 2004; Nelson *et al.*, 2011), and the characteristic water beetle species *Agabus labiatus* (nt), *Graptodytes bilineatus* (nt), *Berosus signaticollis* (EN) and *Dryops similis* (nt) (see Bilton, 1988; Bilton and Lott, 1991; Foster *et al.*, 1992; Bradish *et al.*, 2002; Foster *et al.*, 2009). As well as these rare and threatened water beetle species, typical aquatic coleopteran species include: *Haliphus obliquus*, *Agabus nebulosus*, *Rhantus frontalis*, *Hygrotus impressopunctatus*, *Helophorus minutus*, *Laccobius colon*, *L. minutus*, and *Ochthebius minimus* (Garth Foster, pers. comm.). *Bagous brevis* is a critically endangered (CR) aquatic weevil that feeds on *Ranunculus flammula*. The only recent records for this species are from Knockaunroe turlough (Morris, 1985; Bilton and Lott, 1991; Foster *et al.*, 2009). *Siphonurus armatus* is a critically endangered (CR) mayfly, the only recent record of which is from a turlough (Kelly-Quinn and Regan, 2012). Both these species and their potential association with turloughs require further study.

Table 3 Typical aquatic invertebrate species of turloughs. The species included are characteristic and/or indicative of good quality in turloughs. Those species indicated by * have a strong association with turloughs, with most/all records for that species in Ireland coming from turloughs. Regional Red List status is given, where relevant, as a superscript.

<p>Aquatic invertebrates:</p> <p>Plathyhelminthes, Turbellaria: <i>Polycelis nigra</i>; Crustacea, Cladocera: <i>Alonella excisa</i>, <i>Alona rustica</i>, <i>Alonopsis elongate</i>, <i>Alona affinis</i>, <i>Eurycercus glacialis</i>*, Crustacea, Copepoda: <i>Diaptomus castor</i>;</p> <p>Insecta, Odonata: <i>Lestes dryas</i>*, <i>Sympetrum sanguineum</i>; Insecta, Coleoptera: <i>Agabus labiatus</i>*^{NT}, <i>Agabus nebulosus</i>, <i>Bagous limosus</i>^{CR}, <i>Berosus signaticollis</i>*^{EN}, <i>Dryops similis</i>*^{NT}, <i>Graptodytes bilineatus</i>*^{NT}, <i>Haliphus obliquus</i>, <i>Haliphus variegates</i>^{VU}, <i>Helophorus minutus</i>, <i>Helophorus nanus</i>^{VU}, <i>Hygrotus impressopunctatus</i>, <i>Laccobius colon</i>, <i>Laccobius minutus</i>, <i>Ochthebius minimus</i>, <i>Rhantus frontalis</i>.</p>
--

Characteristic turlough molluscan species include: *Stagnicola fuscus*, *Galba truncatula*, *Radix balthica*, *Anisus leucostoma*, *Oxyloma elegans* and *Pisidium personatum* (Evelyn Moorkens, pers.

comm.). High quality turloughs contain a good mixture of these species in relatively even abundance (i.e. no one species dominating) (Evelyn Moorkens, pers. comm.). *Valvata cristata* and *Armiger crista* may also be present but restricted to the more stable areas of water (Evelyn Moorkens, pers. comm.).

A survey of water beetles at Ballinastack in 2001/2002 did not yield rare or typical turlough species.

The **target** for the attribute typical species, aquatic invertebrates is: Maintain typical species within and across sites.

4.11.3 Typical species: other invertebrates

NPWS (2007, 2013) provides a list of typical terrestrial invertebrate species of turloughs (see Table 4). The EU Interpretation Manual (CEC, 2007) lists the following terrestrial invertebrate species as characteristic of turloughs: *Agonum lugens*, *A. livens*, *Badister meridionalis*, *Blethisa multipunctata* and *Pelophila borealis* (dry phase).

The available information on terrestrial invertebrates was reviewed by Sheehy Skeffington *et al.* (2006). They state “The terrestrial invertebrate fauna of turloughs includes several rare species and communities. Yet to date only some faunal groups have received much attention. The beetles and butterflies have been described to some extent, but the remaining terrestrial orders are poorly documented.”

Terrestrial Coleoptera have been recorded by a number of authors (e.g. Speight, 1976, 1977; Lott and Foster, 1990; Lott and Bilton 1991; Anderson, 1997; Owen, 1997; Good and Butler, 2001; Ní Bhriain *et al.*, 2002; Lott, 2003; Moran *et al.*, 2003; Regan and Anderson, 2004; Moran, 2005; Regan, 2005 a and b; Regan and Moran, 2005).

Bond (1997) noted that no lepidopteran species were exclusively associated with turloughs, but that the upper turlough zones and, especially, fringing scrub had high diversity and hosted some scarce and rare species. The frequency and longevity of flooding is probably a limiting factor for many of the terrestrial species and many of the claimed associations of species with turloughs remain unverified. For example, several species of Lepidoptera feed on *Rhamnus*, which tends to grow on pavement near and at the upper margins of turloughs, and so have been associated with the habitat. There is no evidence, however, that these species have any requirement for the turlough habitat itself. The Lepidoptera species listed in Table 4 are those which have been associated with turloughs and which feed on wetland species of plant. A possibly un-described species of micromoth of the genus *Elachista* (Elachistidae) was recently found at Coolorta Turlough in the Burren, the caterpillars of which were feeding on *Cladium mariscus* (Phil Sterling, pers. comm.).

Insufficient and lack of comprehensive survey of other groups limits selection of other typical species to a few. Morris (1974) provides an account of the Auchenorrhyncha (Hemiptera) of the Burren which refers to species associated with wetlands. However, work such as this needs to be repeated before definitive turlough associations can be made.

Table 4 The typical terrestrial invertebrate species of turloughs (NPWS, 2007, 2013). Species indicated by * have a strong association with turloughs, i.e. most/all records for that species in Ireland are from turloughs.

Terrestrial invertebrates:
<p>Diptera, Sciomyzidae: <i>Pherbellia nana</i>, <i>Colobaea distincta</i>, <i>Ilione albiceta</i>, <i>Pherbina coryleti</i>; Lepidoptera, Crambidae: <i>Paraponyx stratiotata</i>; Lepidoptera, Tortricidae: <i>Bactra furfurana</i>; Lepidoptera, Gelechiidae <i>Monochroa lutulentella</i>; Lepidoptera, Noctuidae: <i>Deltote uncula</i>; Coleoptera, Carabidae: <i>Blethisa multipunctata</i>, <i>Chlaenius nigricornis</i>, <i>Pelophila borealis</i>, <i>Agonum piceum</i>, <i>Carabus granulatus</i>, <i>Loricera pilicornis</i>, <i>Pterostichus nigrita</i>, <i>Bembidion clarkii</i>, <i>Agonum muelleri</i>, <i>Bembidion aeneum</i>, *<i>Agonum lugens</i>, *<i>Platynus livens</i>, *<i>Badister meridionalis</i>, *<i>Badister peltatus</i>; Coleoptera, Staphylinidae: <i>Philonthus furcifer</i>; Coleoptera, Silphidae: <i>Thanatophilus dispar</i>. Orthoptera, Tetrigidae <i>Tetrix subulata</i>; Orthoptera, Acrididae <i>Chorthippus albomarginatus</i> Heteroptera, Saldidae <i>Saldula opacula</i></p>

The **target** for the attribute typical species, other invertebrates is: Maintain typical species within and across sites.

4.11.4 Typical species: birds

The aquatic phase of turloughs has long been recognised as of importance for wintering waterbirds (Buckley and McCarthy, 1987; Buckley, 1993; Madden and Heery, 1997; Crowe, 2005). Turloughs can provide rich feeding grounds, and can also be important roosting areas. Waterbird usage may also contribute to the colonisation or re-colonisation of turloughs by certain invertebrate species (see Frisch *et al.*, 2007). Turloughs near the coast and those near large lakes, generally have higher numbers of waterbirds than the Burren or land-locked turloughs. Coastal turloughs can be particularly important for feeding and roosting during high tides and stormy seas. The more oligotrophic turloughs, however, tend to host fewer birds.

NPWS site files say that Ballinastack turlough attracts wintering waterfowl, which move between this site and other wetlands in the Glenamaddy area according to water levels and disturbance. The files say that three species listed on Annex I of the E.U. Birds Directive occur: Greenland White-fronted Goose, Whooper Swan and Golden Plover. The Greenland White-fronted sub-flock that uses the turlough is part of the River Suck population. Wigeon is also regular in winter. Ballinastack is not surveyed as part of the Irish Wetland Bird Survey (I-WeBS).

The **target** for the attribute typical species, birds is: Maintain typical species within and across sites.

4.12 Fringing habitats

Marginal woodland and scrub, as well as fringing limestone pavement and semi-natural grasslands are important for terrestrial invertebrates, particularly Lepidoptera (see also Section 4.11.3 above). *Rhamnus cathartica* (Purging Buckthorn) appeared to be of special significance (Bond, 1997). The Irish Annulet (*Odontognophos dumeteta*) is one such rare species that could have an association

with the fringing habitats of turloughs, as it feeds on *Rhamnus cathartica*. These fringing habitats are also important over-wintering sites for many terrestrial coleopteran species that occupy the turlough during the dry phase (Good and Butler, 2001; Lott, 2003).

Semi-natural and natural fringing habitats are also likely to act as seed/propagule source areas for turlough plant species.

Fringing grasslands may also be important feeding areas for wintering waterbirds.

Raised bog is an important and less common fringing habitat for some turloughs, such as Ballinastack, in east Galway and Roscommon. The co-occurrence of and the maintenance/restoration of transitional vegetation between these two priority habitats is of high conservation value. Further investigation is required of the use of these habitats where they co-occur by invertebrate, bird and mammal species.

The **target** for the attribute fringing habitats is: Maintain marginal fringing habitats that support turlough vegetation, invertebrate, mammal and/or bird populations.

4.13 Vegetation structure: turlough woodland

The natural climax vegetation for at least the upper margins of turloughs is wet-woodland. Goodwillie (2003) notes “Without grazing, a ‘wet’ turlough would have a central area of wetland vegetation made up of aquatic or amphibious plants that could start development when still water-covered, and be surrounded by small trees and shrubs, especially willows.” “A ‘dry’ turlough would be totally covered by trees though the trees in the centre would be relatively small because of periodic kills caused by spring or summer flooding.” Owing to traditional grazing of turloughs, turlough woodland is now quite rare. Some turloughs, however, notably those in the Coole-Garryland complex, have areas of mature, native woodland that floods most years. Perrin *et al.* (2008) identified a variant of their *Alnus glutinosa* – *Filipendula ulmaria* group, *Crataegus monogyna* – *Geranium robertianum* woodland (3 d) from the upper margins of turlough basins. They described this as a very rare type of stand of low scrub woodland dominated by *Crataegus monogyna* and *Rhamnus cathartica* (Perrin *et al.*, 2008). This, and possibly other not yet described turlough woodland sub-types, are important elements of the natural variation of turloughs in Ireland.

The **target** for the attribute vegetation structure, woodland structure is: Maintain appropriate turlough woodland diversity and structure.

5. References

Ali, T.H., Holmes, J.M.C. and Grainger, J.N.R., (1987) *Diaptomus cyaneus* Gurney, a freshwater copepod new to Britain and Ireland. *Irish Naturalists' Journal* 22 (6): 240–241.

Anderson, R. (1997) A second Irish locality for *Agonum livens* (Gyllenhal) (Carabidae). *The Coleopterist* 5 (3): 76.

Bilton, D.T. (1988) A survey of aquatic Coleoptera in central Ireland and the Burren. *Bulletin of the Irish Biogeographical Society* 11: 77-94.

- Bilton, D.T. and Lott, D.A. (1991) Further records of aquatic Coleoptera from Ireland. *Irish Naturalists' Journal* 23 (10): 389-397.
- Boland, H., Crowe, O. and Walsh, A. (2008) Irish Wetland Bird Survey: Results of waterbird monitoring in Ireland in 2006/07. *Irish Birds* 8: 341-350.
- Boland, H. and Crowe, O. (2012) Irish Wetland Bird Survey: Waterbird Status and Distribution 2001/02 – 2008/09. BirdWatch Ireland, Kilcoole, Co. Wicklow.
- Bond, K.G.M. (1997) Insect survey, Lepidoptera. In: An Investigation of the Flooding Problems in the Gort–Ardrahan Area of South Galway. Ecology Baseline Study Vol. II. Southern Water Global and Jennings O'Donovan and Partners (eds.). The Office of Public Works, Dublin. pp. 1–86.
- Bradish, S., O Connor Á. and Reed, T. (2002) New records of the water beetles *Berosus signaticollis* (Charpentier) and *Graptodytes bilineatus* (Sturm) in turloughs. *Irish Naturalists' Journal* 27: 83-84.
- Buckley, P. (1993) The Bird Communities and General Ecology of Rahasane Turlough and the Dunkellin/Lavally River System. Unpublished Ph.D. Thesis, Department of Zoology, University College, Galway.
- Buckley, P. and McCarthy, T.K. (1987) Bird Communities in the Dunkellin/Lavally Catchments. A Pre-drainage Survey and Environmental Impact Assessment. Unpublished report to Forest and Wildlife Service, Dublin.
- Byrne, A., Moorkens, E.A., Anderson, R., Killeen, I.J. and Regan, E.C. (2009) Ireland Red List No. 2 – Non-Marine Molluscs. National Parks and Wildlife Service, Department of the Environment, Heritage and Local Government, Dublin, Ireland.
- Commission of the European Communities (2013) Interpretation manual of European Union habitats. Eur 28. April 2013. European Commission DG Environment.
- Coxon, C. (1986) A study of the geology, hydrology and geomorphology of turloughs. Unpublished Ph.D. Thesis, University of Dublin, Trinity College, Ireland.
- Coxon, C.E. (1987a) The spatial distribution of turloughs. *Irish Geography* 20: 11-23.
- Coxon, C.E. (1987b) An examination of the characteristics of turloughs using multivariate statistical techniques. *Irish Geography* 20: 24-42.
- Coxon, C. (1994) Carbonate deposition in turloughs (seasonal lakes) on the western limestone lowlands of Ireland I: present day processes. *Irish Geography* 27: 14-27.
- Coxon, C. and Coxon, P.(1994) Carbonate deposition in turloughs (seasonal lakes) on the western limestone lowlands of Ireland II: the sedimentary record. *Irish Geography* 27: 28-35.
- Crowe, O. (2005) *Ireland's Wetlands and their Waterbirds: Status and Distribution*. BirdWatch Ireland, Dublin.
- Crowe, O., Austin, G. E., Colhoun, K., Cranswick, P. Kershaw, M. and Musgrove A. J. (2008) Estimates and trends of waterbird numbers wintering in Ireland, 1994/95-2003/04. *Bird Study* 55: 66–77.

- Cunha Pereira, H., Allott, N. and Coxon, C. (2010) Are seasonal lakes as productive as permanent lakes? A case study from Ireland. *Canadian Journal of Fisheries and Aquatic Sciences* 67: 1-13.
- Cunha Pereira, H., Allott, N., Coxon, C., Naughton O., Johnston P.M. and Gill L.W. (2011) Phytoplankton of turloughs (seasonal karstic Irish lakes). *Journal of Plankton Research* 33: 385-403.
- Curtis, T.G.F. and McGough, H.N. (1981) A study of the wetlands of the Fergus catchment and adjoining areas. Unpublished Report to the National Parks and Wildlife Service, Dublin.
- Duigan, C.A. and Frey, D.G. (1987a) *Eurycercus glacialis*, a chydorid cladoceran new to Ireland. *Irish Naturalists' Journal* 22 (5): 180–183.
- Duigan, C.A. and Frey, D.G. (1987b) *Eurycercus glacialis* in Ireland (Cladocera, Chydoridae). *Internationale Revue der Gesamten Hydrobiologie* 72 (2): 235–249.
- Foster, G.N., Nelson, B.H., Bilton, D.T., Lott, D.A., Merrit, R., Weyl, R.S. and Eyre, M.D., (1992) A classification and evaluation of Irish water beetle assemblages. *Aquatic Conservation: Marine and Freshwater Ecosystems* 2: 185-208.
- Foster, G. N., Nelson, B. H. and O Connor, Á. (2009) Ireland Red List No. 1 – Water beetles. National Parks and Wildlife Service, Department of Environment, Heritage and Local Government, Dublin, Ireland.
- Frisch, D., Green, A.J. and Figuerola, J. (2007) High dispersal capacity of a broad spectrum of aquatic invertebrates via waterbirds. *Aquat. Sci.* 69, 568–574.
- Gill, L.W. (2010). Modelling a network of turloughs. Unpublished Ph.D. Thesis, Department of Civil, Structural and Environmental Engineering, University of Dublin, Trinity College, Ireland.
- Good, J.A. and Butler, F.T. (2001) Turlough pastures as a habitat for Staphylinidae and Carabidae (Coleoptera) in south-east Galway and north Clare, Ireland. *Bulletin of the Irish Biogeographical Society* 25: 74–88.
- Goodwillie, R. (1992) Turloughs over 10 hectares: Vegetation survey and evaluation. Unpublished report to the National Parks and Wildlife Service, Dublin.
- Goodwillie, R. (2003) Vegetation of turloughs. In: *Wetlands of Ireland: Distribution, ecology, uses and economic value* (M. L. Otte, Ed.). University College Dublin Press, Dublin. pp. 135-144.
- Goodwillie, R., Heery, S. and Keane, S. (1997) Wetland vegetation on the Gort lowlands. In: *An Investigation of the Flooding Problems in the Gort–Ardrahan Area of South Galway. Ecology Baseline Study Vol. I.* Southern Water Global and Jennings O’Donovan and Partners (eds.). The Office of Public Works, Dublin. pp. 1–131.
- Hill, M.O., Mountford, J. O., Roy, D.B. and Bunce R.G.H. (1999) Ellenberg’s indicator values for British plants ECOFACT Volume 2, Technical Annex. Department of Environment, Transport and the Regions for HMSO, Norwich.
- Ivimey-Cook, R.B. and Proctor, M.C.F. (1966) The plant communities of the Burren Co. Clare. *Proceedings of the Royal Irish Academy* 64 (B), 211–301.

- Kelly-Quinn, M. and Regan, E.C. (2012) *Ireland Red List No. 7: Mayflies (Ephemeroptera)*. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht, Dublin, Ireland.
- Kimberley, S. (2008) Spatial and temporal fluxes of plant nutrients in turlough soils. Unpublished Ph.D. Thesis, University of Dublin, Trinity College, Ireland.
- Kimberley, S. and Coxon, C.E. (2013) *Evaluating the Influence of Groundwater Pressures on Groundwater-Dependent Wetlands*. STRIVE Report series 100. Environmental Protection Agency, Wexford.
- Kimberley, S., Naughton, O., Gill, L., Johnston, P. and Waldren, S. (2012) The influence of flood duration on the surface soil properties and grazing management of karst wetlands (turloughs) in Ireland. *Hydrobiologia* 696: 29-41.
- Kimberley, S. and Waldren, S. (2012) Examinations of turlough soil property spatial variation in a conservation assessment context. *Biology and Environment: Proceedings of the Royal Irish Academy*, 112B (2): 1-13.
- Kimberley, S., Naughton, O., Johnston, P., Gill, L. and Waldren, S. (2012) The influence of flood duration on the surface soil properties and grazing management of karst wetlands (turloughs) in Ireland. *Hydrobiologia* 692 (1): 29-40.
- Lockhart, N., Hodgetts, N. and Holyoak, D. (2012) *Rare and threatened Bryophytes of Ireland*. National Museums Northern Ireland Publication No. 028, Holywood, Co. Down.
- Lockhart, N., Hodgetts, N. and Holyoak, D. (2012) *Ireland Red List No.8: Bryophytes*. National Parks and Wildlife Service, Department of Arts, Heritage and the Gaeltacht, Dublin, Ireland.
- Lott, D.A. (2003) An annotated List of Wetland Ground Beetles (Carabidae) and Rove Beetles (Staphylinidae) found in the British Isles including a literature review of their ecology. Report No. 488. English Nature, Peterborough.
- Lott, D.A. and Bilton, D.T. (1991) Records of Coleoptera from Irish wetland sites in 1989. *Bulletin of the Irish Biogeographical Society* 14: 60–72.
- Lott, D.A. and Foster, G.N. (1990) Records of terrestrial Coleoptera from wetland sites in 1987, including *Stenus glabellus* Thomson (Staphylinidae) new to the British Isles. *Irish Naturalists' Journal* 23: 280–282.
- MacGowran, B. (1985) Phytosociological and ecological studies on turloughs in the west of Ireland. Unpublished Ph.D. Thesis, Department of Botany, University College, Galway, Ireland.
- Madden, B. and Heery, S. (1997) Breeding Waterfowl in South Galway Area. In: An Investigation of the Flooding Problems in the Gort–Ardrahan Area of South Galway. Ecology Baseline Study Vol. I. Southern Water Global and Jennings O'Donovan and Partners (eds.). The Office of Public Works, Dublin. Report September 1996. Report to the Office of Public Works. pp. 132–208.

- Moran, J., Gormally, M.J. and Sheehy Skeffington, M. (2003) Records of *Panagaeus crux-major* (Linnaeus) and *Pterostichus aterrimus* Herbst recorded on turloughs in Counties Clare and Mayo. *Irish Naturalists Journal* 27 (8): 309–311.
- Moran, J. (2005) Skealaghan Turlough, County Mayo: Implications of grazing and flooding regimes for plant and carabid beetle communities with reference to turlough farming systems in the region. Unpublished Ph.D. Thesis, National University of Ireland, Galway.
- Moran, J., Sheehy Skeffington, M. and Gormally, M. (2008) The influence of hydrological regime and grazing management on the plant communities of a karst wetland (Skealaghan turlough) in Ireland. *Applied Vegetation Science* 11: 13-24.
- Morris, M.G. 1974. Auchenorrhyncha (Hemiptera) of the Burren, with special reference to species-associations of the grasslands. *Proceedings of the Royal Irish Academy*. 74B: 7-30.
- Morris, M.G. (1985) *Bagous brevis* Gyllenhal new to Ireland from the Burren, Co. Clare, with a brief review of the Irish Bagoiini (Coleoptera: Curculionidae). *Irish Naturalists' Journal* 21: 400-403.
- Naughton, O. (2011). *The Hydrology and Hydroecology of Turloughs*. Unpublished PhD thesis, University of Dublin.
- Naughton, O. and Johnston, P. (2009) Hydrology and hydrogeology of turloughs as wetlands. In *Proceedings of the 29th IAH Irish Group Groundwater Conference*, Tullamore, 21-22 April 2009, International Association of Hydrogeologists (Irish Group), Dublin, Ireland. 99-106.
- Nelson, B. and Thompson, R. (2004) The Natural History of Ireland's Dragonflies. The National Museums and Galleries of Northern Ireland, Ulster Museum, Belfast.
- Nelson, B., Ronayne, C. and Thompson, R. (2011) *Ireland Red List No.6: Damselflies & Dragonflies (Odonata)*. National Parks and Wildlife Service, Department of the Environment, Heritage and Local Government, Dublin, Ireland.
- NPWS (2007) Turloughs, Draft Backing Document. In: The Status of EU Protected Habitats and Species in Ireland. Backing Documents, Article 17 Forms, Maps. Volume 2. Unpublished National Parks and Wildlife Service Report, Dublin. pp. 510-550.
- NPWS (2008) The Status of EU Protected Habitats and Species in Ireland. Conservation Status in Ireland of Habitats and Species listed in the European Council Directive on the Conservation of Habitats, Flora and Fauna 92/43/EEC. Unpublished National Parks and Wildlife Service Report, Dublin.
- NPWS (2013) The Status of EU Protected Habitats and Species in Ireland. Habitat Assessments Volume 2, Version 1.1. Unpublished Report, National Parks and Wildlife Services, Department of Arts, Heritage and the Gaeltacht, Dublin, Ireland. 390-408.
- Naughton, O. (2011) The Hydrology and Hydroecology of Turloughs. Unpublished Ph.D. Thesis. University of Dublin, Trinity College, Ireland.

Naughton, O., Johnston, P.M and Gill, L.W. (2012) Groundwater flooding in Irish karst: The hydrological characterisation of ephemeral lakes (turloughs). *J. Hydrol.* (2012), <http://dx.doi.org/10.1016/j.jhydrol.2012.08.012>

Ní Bhriain, B., (1999) A study of a turlough in the Burren Co. Clare in its agricultural context. Unpublished B.Sc. (Hons) Thesis, NUI, Galway.

Ní Bhriain, B., Sheehy Skeffington, M. and Gormally, M. (2002) Conservation implications of the land use practices on the plant and carabid beetle communities of two turloughs in Co. Galway Ireland. *Biological Conservation* 1, 81–92.

O’Connell, M., Ryan, J.B and MacGowran, B.A. (1984) Wetland communities in Ireland: a phytosociological review. In: *European Mires* (P. D. Moore, Ed.). Academic Press, London. pp. 303-364.

OECD (Organization for Economic Cooperation and Development). (1982) Eutrophication of waters. Monitoring, assessment and control. Final report, OECD cooperative programme on monitoring of inland waters (eutrophication control), Environment Directorate, OECD, Paris. 154 pp.

Perrin, P. Martin, J., Barron, S., O’Neill, F., McNutt, K. and Delaney, A. (2008) National Survey of Native Woodlands, 2003-2008. Volume II, Woodland Classification. Unpublished Report by BEC Environmental Consultants to the National Parks and Wildlife Service, Dublin.

Porst, G. (1999) The Effects of Season, Habitat, Hydroperiod and Water Chemistry on the Distribution of Turlough Aquatic Invertebrate Communities. Unpublished Ph.D. Thesis. University of Dublin, Trinity College, Ireland.

Regan, E.C. (2005a) An investigation of the plant, carabid, and staphylinid communities of turloughs in southeast Galway/north Clare, Ireland. Unpublished Ph.D. Thesis, National University of Ireland, Galway.

Regan, E.C. (2005b) Further records of carabid beetles from turloughs. *Irish Naturalists’ Journal* 28 (2): 59–61.

Regan, E.C. and Anderson, R. (2004) Terrestrial Coleoptera recorded in Ireland, May 2003. *Bulletin of the Irish Biogeographical Society* 28: 85–132.


Regan, E.C. and Moran, J. (2005) *Thanatophilus dispar* (Herbst) (Silphidae) in turloughs in the west of Ireland. *The Coleopterist* 14 (2): 89–91.

Regan, E.C., Sheehy Skeffington, M. and Gormally, M.J. (2007) Wetland plant communities of turloughs in southeast Galway/north Clare, Ireland in relation to environmental factors. *Aquatic Botany* 27: 22-30.

Reynolds, J.D. (1997) Invertebrate surveys of S.E. Galway turloughs. Baseline Report. In: An Investigation of the Flooding Problems in the Gort–Ardrahan Area of South Galway. Ecology Baseline Study Vol. II. Southern Water Global and Jennings O’Donovan and Partners (eds.). The Office of Public Works, Dublin. pp. 126–141.

- Reynolds, J.D. (2000) Invertebrate communities of turloughs (temporary lakes) in south-east Galway Ireland. *Verhandlungen Internationale Vereinigung für Theoretische und Angewandte Limnologie* 27 (3): 1679–1684.
- Reynolds, J.D. (2016) Invertebrates of Irish Turloughs (Chapter 6). In: D. Batzer, D. Boix (eds.), *Invertebrates in Freshwater Wetlands*. Springer International Publishing Switzerland 2016. 191-217.
- Reynolds, J.D. and Marnell, F. (1999) New records of *Eurycerus glacialis* (Cladocera: Chydoridae) in turloughs in south-east Galway. *Irish Naturalists' Journal* 26: 177–180.
- Reynolds, J.D., Murphy, M. and O Connor, Á. (2004) Early season cladoceran diversity of Atlantic temporary ponds (turloughs). *Archives des Sciences* 57: 97–104.
- Ryder, C., Moran, J., McDonnell, R. and Gormally, M. (2005) Conservation implications of grazing practices on the plant and dipteran communities of a turlough in Co. Mayo, Ireland. *Biodiversity and Conservation* 14: 186–187.
- Sharkey, N. (2012) Turlough vegetation communities – links with hydrology, hydrochemistry, soils and management. Unpublished Ph.D. Thesis. University of Dublin, Trinity College, Ireland.
- Sheehy Skeffington, M., Moran, J., O Connor, Á., Regan, E., Coxon, C.E., Scott, N.E. and Gormally, M., (2006) Turloughs – Ireland's unique wetland habitat. *Biological Conservation* 133: 265-290.
- Speight, M.C.D. (1976) *Agonum livens*, *Asemum striatum* and *Xylota coeruleiventris*: insects new to Ireland. *Irish Naturalists' Journal* 18: 274–275.
- Speight, M.C.D. (1977) Notes on three ground beetles (Coleoptera: Carabidae) *Dyschirius luedersi* new to Ireland, *Badister peltatus* and *Chlaenius tristis* reinstated as Irish. *Irish Naturalists' Journal* 19 (4): 116–118.
- Tynan, S., Gill, M. and Johnston, P. (2007). Development of a methodology for the characterisation of a karstic groundwater body with particular emphasis on the linkage with associated ecosystems such as turlough ecosystems. 2002-W-DS-8-M1. Final Report. Environmental RDTI programme 2000-2006, Water Framework Directive. Environmental Protection Agency, Wexford.
- Waldren, S. (Ed.) (2015) Turlough Hydrology, Ecology and Conservation. Unpublished Report, National Parks and Wildlife Services, Department of Arts, Heritage and the Gaeltacht, Dublin, Ireland.
- Working Group on Groundwater (Turlough sub-committee) (2005) Risk Assessment Sheet GWDTERA2a - Turloughs. Guidance Document No. GW9. Guidance on the Pressures and Impacts on Groundwater Dependent Terrestrial Ecosystems. WFD Pressures and Impact Assessment Methodology. <http://www.wfdireland.net/wfd-charreport.html>
- Waldren, S., Lynn, D.E. and Murphy, S. (2002) The Effects of Management Practices on the Dynamics and Conservation Status of Turlough Plant Communities. Final Scientific and Technical Report to Enterprise Ireland.

Appendix 1 Map of Ballinastack Turlough



Appendix 2 Orthophotograph of Ballinastack Turlough

